

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



let

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/06, A61K 31/415, 31/435		A1	(11) International Publication Number: WO 98/08840 (43) International Publication Date: 5 March 1998 (05.03.98)
(21) International Application Number: PCT/US97/14912 (22) International Filing Date: 25 August 1997 (25.08.97) (30) Priority Data: 60/025,123 29 August 1996 (29.08.96) US 60/033,579 19 December 1996 (19.12.96) US 9703015.9 13 February 1997 (13.02.97) GB 60/047,177 20 May 1997 (20.05.97) US (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DUGGAN, Mark, E. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HARTMAN, George, D. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HOFFMAN, William, F. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MEISSNER, Robert, S. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PERKINS, James, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). ASKEW, Ben, C. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). COLEMAN, Paul, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HUTCHIN-		SON, John, H. [CA/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). NAYLOR-OLSEN, Adel, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: INTEGRIN ANTAGONISTS			
(57) Abstract This invention relates to certain novel compounds and derivatives thereof, their synthesis, and their use as vitronectin receptor antagonists. The vitronectin receptor antagonist compounds of the present invention are $\alpha v\beta 3$ antagonists, $\alpha v\beta 5$ antagonists or dual $\alpha v\beta 3/\alpha v\beta 5$ antagonists useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation and tumor growth.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	R	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

TITLE OF THE INVENTION INTEGRIN ANTAGONISTS

FIELD OF THE INVENTION

5 The present invention is related to U.S. provisional applications Serial Nos. 60/047,177, filed May 20, 1997, 60/033,579, filed December 19, 1996, and 60/025,123, filed August 29, 1996, the contents of which are hereby incorporated by reference.

10 The present invention provides novel compounds and derivatives thereof, their synthesis, and their use as vitronectin receptor ligands. More particularly, the compounds of the present invention are $\alpha v \beta 3$ antagonists, $\alpha v \beta 5$ antagonists or dual $\alpha v \beta 3 / \alpha v \beta 5$ antagonists useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy,
15 macular degeneration, angiogenesis, atherosclerosis, inflammation and tumor growth.

BACKGROUND OF THE INVENTION

20 This invention relates to compounds for inhibiting bone resorption that is mediated by the action of a class of cells known as osteoclasts.

 Osteoclasts are multinucleated cells of up to 400 μm in diameter that resorb mineralized tissue, chiefly calcium carbonate and calcium phosphate, in vertebrates. They are actively motile cells that
25 migrate along the surface of bone. They can bind to bone, secrete necessary acids and proteases and thereby cause the actual resorption of mineralized tissue from the bone.

 More specifically, osteoclasts are believed to exist in at least two physiological states. In the secretory state, osteoclasts are flat,
30 attach to the bone matrix via a tight attachment zone (sealing zone), become highly polarized, form a ruffled border, and secrete lysosomal enzymes and protons to resorb bone. The adhesion of osteoclasts to bone surfaces is an important initial step in bone resorption. In the

- 2 -

migratory or motile state, the osteoclasts migrate across bone matrix and do not take part in resorption until they attach again to bone.

Integrins are transmembrane, heterodimeric, glycoproteins which interact with extracellular matrix and are involved in osteoclast attachment, activation and migration. The most abundant integrin in osteoclasts (rat, chicken, mouse and human) is the vitronectin receptor, or $\alpha v \beta 3$, thought to interact in bone with matrix proteins that contain the RGD sequence. Antibodies to $\alpha v \beta 3$ block bone resorption in vitro indicating that this integrin plays a key role in the resorptive process.

There is increasing evidence to suggest that $\alpha v \beta 3$ ligands can be used effectively to inhibit osteoclast mediated bone resorption in vivo in mammals.

The current major bone diseases of public concern are osteoporosis, hypercalcemia of malignancy, osteopenia due to bone metastases, periodontal disease, hyperparathyroidism, periarticular erosions in rheumatoid arthritis, Paget's disease, immobilization-induced osteopenia, and glucocorticoid treatment.

All these conditions are characterized by bone loss, resulting from an imbalance between bone resorption (breakdown) and bone formation, which continues throughout life at the rate of about 14% per year on the average. However, the rate of bone turnover differs from site to site, for example, it is higher in the trabecular bone of the vertebrae and the alveolar bone in the jaws than in the cortices of the long bones. The potential for bone loss is directly related to turnover and can amount to over 5% per year in vertebrae immediately following menopause, a condition which leads to increased fracture risk.

There are currently 20 million people with detectable fractures of the vertebrae due to osteoporosis in the United States. In addition, there are 250,000 hip fractures per year attributed to osteoporosis. This clinical situation is associated with a 12% mortality rate within the first two years, while 30% of the patients require nursing home care after the fracture.

Individuals suffering from all the conditions listed above would benefit from treatment with agents which inhibit bone resorption.

- 3 -

Additionally, $\alpha v \beta 3$ ligands have been found to be useful in treating and/or inhibiting restenosis (recurrence of stenosis after corrective surgery on the heart valve), atherosclerosis, diabetic retinopathy, macular degeneration and angiogenesis (formation of new blood vessels). Moreover, it has been postulated that the growth of tumors depends on an adequate blood supply, which in turn is dependent on the growth of new vessels into the tumor; thus, inhibition of angiogenesis can cause tumor regression in animal models. (See, Harrison's Principles of Internal Medicine, 12th ed., 1991). $\alpha v \beta 3$ antagonists, which inhibit angiogenesis, are therefore useful in the treatment of cancer for inhibiting tumor growth. (See e.g., Brooks et al., *Cell*, 79:1157-1164 (1994)).

Moreover, compounds of this invention can also inhibit neovascularization by acting as antagonists of the integrin receptor $\alpha v \beta 5$. A monoclonal antibody for $\alpha v \beta 5$ has been shown to inhibit VEGF-induced angiogenesis in rabbit cornea and the chick chorioallantoic membrane model; M.C. Friedlander, et.al., *Science* 270, 1500-1502, 1995. Thus, compounds that antagonize $\alpha v \beta 5$ are useful for treating and preventing macular degeneration, diabetic retinopathy, and tumor growth.

In addition, certain compounds of this invention antagonize both the $\alpha v \beta 3$ and $\alpha v \beta 5$ receptors. These compounds, referred to as "dual $\alpha v \beta 3 / \alpha v \beta 5$ antagonists," are useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation and tumor growth.

It is an object of the present invention to identify compounds which bind to the $\alpha v \beta 3$ receptor, $\alpha v \beta 5$ receptor or both the $\alpha v \beta 3$ and $\alpha v \beta 5$ receptors.

It is a further object of the invention to identify compounds which act as antagonists of the $\alpha v \beta 3$ receptor. It is another object of the invention to identify $\alpha v \beta 3$ antagonist compounds which are useful agents for inhibiting: bone resorption mediated by osteoclast cells, restenosis, atherosclerosis, inflammation, diabetic retinopathy, macular

- 4 -


degeneration and angiogenesis in animals, preferably mammals, especially humans. Still another object of the invention is to identify $\alpha v \beta 3$ antagonists which cause tumor regression and/or inhibit tumor growth in animals.

- 5 A further object of the invention is to identify $\alpha v \beta 3$ antagonists useful for preventing or treating osteoporosis. An additional object of the invention is to identify $\alpha v \beta 3$ antagonists useful for treating cancer.


- 10 It has now been found that the compounds of the present invention, $\alpha v \beta 3$ ligands, are useful for inhibiting bone resorption in mammals. Thus, the compounds of the present invention are useful for preventing or reducing the incidence of osteoporosis. Additionally, the $\alpha v \beta 3$ ligands of the present invention are also useful for treating and/or inhibiting restenosis, diabetic retinopathy, macular degeneration, 15 atherosclerosis and/or angiogenesis in mammals.

SUMMARY OF THE INVENTION

- The present invention provides a method of eliciting a vitronectin receptor antagonizing effect in a mammal in need thereof, 20 comprising administering to the mammal a therapeutically effective amount of a compound of the formula



wherein X is selected from



- 25 a 5- or 6-membered monocyclic aromatic or nonaromatic ring system containing 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S wherein the 5- or 6-membered ring system is either


- 5 -

unsubstituted or substituted on a carbon or nitrogen atom with one or more groups chosen from R¹, R², R¹⁵ or R¹⁶;

5 a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups chosen from R¹, R², R¹⁵ or R¹⁶;

10

Y is selected from



Z is

15 a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; provided that Z is not a 6-membered monocyclic aromatic ring system, an isoxazoline ring or an isoxazole ring;

20

- 6 -

R¹, R², R⁴, R⁵, R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from

5 hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl,
C₃-8 cycloheteroalkyl, C₃-8 cycloalkyl C₁-6 alkyl,
C₃-8 cycloheteroalkyl C₁-6 alkyl, aryl, aryl C₁-8 alkyl, amino,
amino C₁-8 alkyl, C₁-3 acylamino, C₁-3 acylamino C₁-8 alkyl,
(C₁-6 alkyl)_qamino, (C₁-6 alkyl)_qamino C₁-8 alkyl,
C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl,
10 hydroxycarbonyl C₁-6 alkyl, C₁-3 alkoxycarbonyl,
C₁-3 alkoxycarbonyl C₁-6 alkyl, hydroxycarbonyl-
C₁-6 alkyloxy, hydroxy, hydroxy C₁-6 alkyl, C₁-6 alkyloxy-
C₁-6 alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,
trifluoroethoxy, C₁-8 alkyl-S(O)_q, (C₁-8 alkyl)_qaminocarbonyl,
C₁-8 alkyloxycarbonylamino, (C₁-8 alkyl)_qaminocarbonyloxy,
15 oxo, (aryl C₁-8 alkyl)_qamino, (aryl)_qamino,
aryl C₁-8 alkylsulfonylamino or C₁-8 alkylsulfonylamino;

R³ is selected from

20 hydrogen,
aryl,
aryl-(CH₂)_p-,
hydroxyl,
C₁-5 alkoxy,
aminocarbonyl,
25 C₃-8 cycloalkyl,
amino C₁-6 alkyl,
(aryl)_qaminocarbonyl,
(aryl C₁-5 alkyl)_qaminocarbonyl,
hydroxycarbonyl C₁-6 alkyl,
30 C₁-8 alkyl,
aryl C₁-6 alkyl,
(C₁-6 alkyl)_qamino C₁-6 alkyl,
(aryl C₁-6 alkyl)_qamino C₁-6 alkyl,
C₁-8 alkylsulfonyl,

- 7 -

C₁-8 alkoxy carbonyl,
aryloxy carbonyl,
aryl C₁-8 alkoxy carbonyl,
C₁-8 alkyl carbonyl,
5 aryl carbonyl,
aryl C₁-6 alkyl carbonyl,
(C₁-8 alkyl)_q aminocarbonyl,
aminosulfonyl,
C₁-8 alkyl aminosulfonyl,
10 (aryl)_q aminosulfonylamino,
(aryl C₁-8 alkyl)_q aminosulfonyl,
C₁-6 alkyl sulfonyl,
aryl sulfonyl,
aryl C₁-6 alkyl sulfonyl,
15 aryl C₁-6 alkyl carbonyl,
C₁-6 alkyl thiocarbonyl,
aryl thiocarbonyl, or
aryl C₁-6 alkyl thiocarbonyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
20 R¹³ and R¹⁴;

R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from
hydrogen,
aryl,
25 aryl-(CH₂)_p-,
aryl-(CH₂)_n-O-(CH₂)_m-,
aryl-(CH₂)_n-S(O)_q-(CH₂)_m-,
aryl-(CH₂)_n-C(O)-(CH₂)_m-,
aryl-(CH₂)_n-C(O)-N(R³)-(CH₂)_m-,
30 aryl-(CH₂)_n-N(R³)-C(O)-(CH₂)_m-,
aryl-(CH₂)_n-N(R³)-(CH₂)_m-,
halogen,
hydroxyl,
C₁-8 alkyl carbonylamino,

- 8 -

- aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_qaminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 5 C₃₋₈ cycloalkyl,
 oxo,
 (C₁₋₆ alkyl)_qamino,
 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 10 aryl C₁₋₅ alkylaminocarbonyl,
 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 15 C₁₋₈ alkyl, either unsubstituted or substituted, with one or more
 groups selected from: halogen, hydroxyl,
 C₁₋₅ alkylcarbonylamino, aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl, aminocarbonyl, (C₁₋₅ alkyl)_qamino-
 carbonyl, C₁₋₅ alkylcarbonyloxy, C₃₋₈ cycloalkyl, oxo,
 20 (C₁₋₃ alkyl)_qamino, amino C₁₋₃ alkyl,
 (aryl)_qaminocarbonyl, (aryl C₁₋₅ alkyl)_qaminocarbonyl,
 aminocarbonyl, aminocarbonyl C₁₋₄ alkyl,
 hydroxycarbonyl or hydroxycarbonyl C₁₋₅ alkyl,
 CH≡C-(CH₂)_s-,
 25 C₁₋₆ alkyl-C≡C-(CH₂)_s-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,
 aryl-C≡C-(CH₂)_s-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,
 CH₂=CH-(CH₂)_s-,
 30 C₁₋₆ alkyl-CH=CH-(CH₂)_s-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,
 aryl-CH=CH-(CH₂)_s-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,
 C₁₋₆ alkyl-SO₂-(CH₂)_s-,

- 9 -

- C₁₋₆ alkylaryl-SO₂-(CH₂)_s-,
C₁₋₆ alkoxy,
aryl C₁₋₆ alkoxy,
aryl C₁₋₆ alkyl,
5 (C₁₋₆ alkyl)qamino C₁₋₆ alkyl,
(aryl)qamino,
(aryl)qamino C₁₋₆ alkyl,
(aryl C₁₋₆ alkyl)qamino,
(aryl C₁₋₆ alkyl)qamino C₁₋₆ alkyl,
10 arylcarbonyloxy,
aryl C₁₋₆ alkylcarbonyloxy,
(C₁₋₆ alkyl)qaminocarbonyloxy,
C₁₋₈ alkylsulfonylamino,
arylsulfonylamino,
15 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxycarbonylamino,
20 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
25 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
-arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
30 (C₁₋₈ alkyl)qaminocarbonylamino,
(C₁₋₈ alkyl)qaminocarbonylamino C₁₋₆ alkyl,
(aryl)qaminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)qaminocarbonylamino,
(aryl C₁₋₈ alkyl)qaminocarbonylamino C₁₋₆ alkyl,

- 10 -

- 5 aminosulfonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)_qaminosulfonylamino,
 (C₁₋₈ alkyl)_qaminosulfonylamino C₁₋₆ alkyl,
 (aryl)_qaminosulfonylamino C₁₋₆ alkyl,
 10 (aryl C₁₋₈ alkyl)_qaminosulfonylamino,
 (aryl C₁₋₈ alkyl)_qaminosulfonylamino C₁₋₆ alkyl,
 C₁₋₆ alkylsulfonyl,
 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 arylsulfonyl C₁₋₆ alkyl,
 15 aryl C₁₋₆ alkylsulfonyl,
 aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 C₁₋₆ alkylcarbonyl,
 C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 arylcabonyl C₁₋₆ alkyl,
 20 aryl C₁₋₆ alkylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 C₁₋₆ alkylthiocarbonylamino,
 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 arylthiocarbonylamino C₁₋₆ alkyl,
 25 aryl C₁₋₆ alkylthiocarbonylamino,
 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,
 (aryl)_qaminocarbonyl C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_qaminocarbonyl, or
 30 (aryl C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,
 wherein any of the alkyl groups may be unsubstituted or substituted with
 R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
 are attached is itself attached to no more than one heteroatom; and
 provided further that the carbon atom to which R¹⁰ and R¹¹ are
 30 attached is itself attached to no more than one heteroatom;

R¹² is selected from
 hydrogen,
 C₁₋₈ alkyl,

- 11 -

5 aryl,
aryl C₁₋₈ alkyl,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
C₁₋₈ alkylaminocarbonylmethylene, or
C₁₋₈ dialkylaminocarbonylmethylene;


m, s and t are each independently an integer from 0 to 3;
n is an integer from 1 to 3;
10 p is an integer from 1 to 4;
q is an integer from 0 to 2;
r is an integer from 0 to 6;
and the pharmaceutically acceptable salts thereof.

15 In one embodiment of the present invention is the method
of eliciting a vitronectin antagonizing effect wherein
X is


20 a 9- to 10-membered polycyclic ring system, wherein one or
more of the rings is aromatic, and wherein the polycyclic ring
system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or
S, and wherein the polycyclic ring system is either unsubstituted
or substituted on a carbon atom with R¹ and R²; and

Z is selected from

- 12 -




- 13 -




and all other variables are as defined above;

and the pharmaceutically acceptable salts thereof. Preferably, Z is


5 selected from



- 14 -




In a class of the invention is the method of eliciting a vitronectin antagonizing effect wherein the compound has the formula



wherein X is selected from

- 15 -



Y is selected from $-(CH_2)_r-$ or $-(CH_2)_m-NR^3-(CH_2)_t-$;

R^3 is selected from

- hydrogen,
- 5 aryl- $(CH_2)_p-$,
- C₁₋₅ alkoxycarbonyl,
- C₃₋₈ cycloalkyl,
- (aryl)_qaminocarbonyl,
- (aryl C₁₋₅ alkyl)_qaminocarbonyl,
- 10 C₁₋₈ alkyl,
- aryl C₁₋₆ alkyl,
- C₁₋₈ alkylsulfonyl,
- arylsulfonyl,
- aryl C₁₋₆ alkylsulfonyl,
- 15 C₁₋₈ alkoxycarbonyl,
- aryloxycarbonyl,
- aryl C₁₋₈ alkoxycarbonyl,
- C₁₋₈ alkylcarbonyl,
- arylcarbonyl,
- 20 aryl C₁₋₆ alkylcarbonyl,
- (C₁₋₈ alkyl)_qaminocarbonyl,
- C₁₋₆ alkylsulfonyl, or
- aryl C₁₋₆ alkylcarbonyl,
- wherein any of the alkyl groups may be unsubstituted or
- 25 substituted with R^{13} and R^{14} ;

R^4 is selected from

- hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloheteroalkyl,
- C₃₋₈ cycloalkyl C₁₋₆ alkyl, C₃₋₈ cycloheteroalkyl C₁₋₆ alkyl,
- 30 aryl or aryl C₁₋₈ alkyl,


- 16 -

- R^8 is selected from
- hydrogen,
 - aryl,
 - 5 aryl-(CH₂)_p-,
 - CH≡C-(CH₂)_s-,
 - C₁-6 alkyl-C≡C-(CH₂)_s-,
 - C₃-7 cycloalkyl-C≡C-(CH₂)_s-,
 - aryl-C≡C-(CH₂)_s-,
 - 10 C₁-6 alkylaryl-C≡C-(CH₂)_s-,
 - CH₂=CH-(CH₂)_s-,
 - C₁-6 alkyl-CH=CH-(CH₂)_s-,
 - C₃-7 cycloalkyl-CH=CH-(CH₂)_s-,
 - aryl-CH=CH-(CH₂)_s-,
 - 15 C₁-6 alkylaryl-CH=CH-(CH₂)_s-,
 - C₁-6 alkyl-SO₂-(CH₂)_s-,
 - C₁-6 alkylaryl-SO₂-(CH₂)_s-; and

r is an integer from 0 to 3;


- 20 and all other variables are as defined above;
and the pharmaceutically acceptable salts thereof.

In a subclass of the invention is the method wherein the compound has the formula




- 25 wherein Z is selected from

- 17 -



R⁸ is selected from
hydrogen,



- 5 indolyl-(CH₂)_p-,
CH≡C-(CH₂)_s-,
C₁₋₆ alkyl-C≡C-(CH₂)_s-,
C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,
aryl-C≡C-(CH₂)_s-,
10 C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,
CH₂=CH-(CH₂)_s-,
C₁₋₆ alkyl-CH=CH-(CH₂)_s-,
C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,
aryl-CH=CH-(CH₂)_s-,
15 C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,

- 18 -

C₁₋₆ alkyl-SO₂-(CH₂)_s-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_s-; and

R¹² is selected from hydrogen or C₁₋₈ alkyl;

5 s is an integer from 0 to 3;

and all other variables are as defined above;

and the pharmaceutically acceptable salts thereof.

Illustrative of the invention is the method of eliciting a
 vitronectin antagonizing effect wherein the compound is selected from

10

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperidin-
 1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

15 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperin-
 1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine trifluoroacetate;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
 yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;

20 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
 yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-
 pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

25

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
 yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;

30 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
 yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
 yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine ethyl ester;

5 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

10 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

15 Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

20 Ethyl 2-oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;

25 Ethyl 3-(2-{2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}-pyrrolidin-1-yl)-acetyl-amino)-3-(S)-pyridin-3-yl-propionic acid;

30 3-(2-{2-Oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}pyrrolidin-1-yl)-acetyl-amino)-3-(S)-pyridin-3-yl-propionic acid;

3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetyl-amino}-3-(S)-pyridin-3-yl-propionic acid;

- 20 -

- 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3(S)-
pyridin-3-yl-propionic acid;
- 5
- 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-
ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;
- 10
- 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;
- 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-
ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;
- 15
- 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;
- 20
- 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl
ester;
- 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;
- 25
- 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-
tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-
acetylamino)-propionic acid ethyl ester; or
- 30
- 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-
tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-
acetylamino)-propionic acid;
- 3-{2-(2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-
pyrrolidin-1-yl)-acetylamino}-3(S)-quinolin-3-yl-propionic acid;

3-(2-(5(S)-Ethyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-acetylamino)-3-(S)-quinolin-3-yl-propionic acid trifluoroacetate;

5

3-(2-(6-Methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl)-acetylamino)-3(S)-pyridin-3-yl-propionic acid bis trifluoroacetate; or

10 3-(2-(6-Methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl)-acetylamino)-3(S)-pyridin-3-yl-propionic acid ethyl ester;

and the pharmaceutically acceptable salts thereof.

15

Preferably, the compound is selected from

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine trifluoroacetate;

20 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

25

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

30

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3-(S)-pyridin-3-yl- β -alanine;

- 22 -

- 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-
tetrahydropyrimidin-1-yl-acetyl-3-(S)-pyridin-3-yl-β-alanine;
- 5 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)-β-alanine;
- 3-(2-{2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
amino]pyrrolidin-1-yl)-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;
- 10 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-
pyridin-3-yl-propionic acid;
- 15 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-
pyridin-3-yl-propionic acid;
- 20 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;
- 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;
- 25 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine; or
- 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-
tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-
acetylamino)-propionic acid;
- 30

and the pharmaceutically acceptable salts thereof.

Exemplifying the invention is the method wherein the vitronectin receptor antagonizing effect is an αvβ3 antagonizing effect. An illustration of the invention is the method wherein the αvβ3


- 23 -

antagonizing effect is selected from inhibition of: bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth. Preferably, the $\alpha v \beta 3$ antagonizing effect is the inhibition of bone resorption.

5 An example of the invention is the method wherein the vitronectin receptor antagonizing effect is an $\alpha v \beta 5$ antagonizing effect. More specifically, the $\alpha v \beta 5$ antagonizing effect is selected from inhibition of: restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth.


10 Illustrating the invention is the method wherein the vitronectin receptor antagonizing effect is a dual $\alpha v \beta 3 / \alpha v \beta 5$ antagonizing effect. More particularly, the dual $\alpha v \beta 3 / \alpha v \beta 5$ antagonizing effect is selected from inhibition of: bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth.

15 In a second embodiment of the present invention is a method of eliciting an $\alpha v \beta 3$ antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound of the formula



20

wherein X is selected from




25

a 5- or 6-membered monocyclic aromatic or nonaromatic ring system containing 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S wherein the 5- or 6-membered ring system is either unsubstituted or substituted on a carbon atom with R^1 and R^2 , or

- 24 -


- 5 a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted on a carbon atom with R^1 and R^2 ;

Y is selected from




Z is

- 10 a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted on a carbon or
- 15 nitrogen atom with one or more groups independently selected from R^4 , R^5 , R^6 and R^7 ; provided that Z is not a 6-membered monocyclic aromatic ring system; preferably, Z is selected from



- 25 -



R¹, R², R³, R⁴, R⁵, R¹³ and R¹⁴ are each independently selected from

- 26 -

- hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, aryl,
aryl C₁-8 alkyl, amino, amino C₁-8 alkyl, C₁-3 acylamino,
C₁-3 acylamino C₁-8 alkyl, C₁-6 alkylamino, C₁-6 alkylamino-
C₁-8 alkyl, C₁-6 dialkylamino, C₁-6 dialkylamino C₁-8 alkyl,
5 C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl,
hydroxycarbonyl C₁-6 alkyl, C₁-3 alkoxycarbonyl,
C₁-3 alkoxycarbonyl C₁-6 alkyl, hydroxycarbonyl-
C₁-6 alkyloxy, hydroxy or hydroxy C₁-6 alkyl;
- 10 R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from
hydrogen,
aryl,
-(CH₂)_p-aryl,
halogen,
15 hydroxyl,
C₁-8 alkylcarbonylamino,
aryl C₁-5 alkoxy,
C₁-5 alkoxycarbonyl,
aminocarbonyl,
20 C₁-8 alkylaminocarbonyl,
C₁-6 alkylcarbonyloxy,
C₃-8 cycloalkyl,
oxo,
amino,
25 C₁-6 alkylamino,
amino C₁-6 alkyl,
arylaminocarbonyl,
aryl C₁-5 alkylaminocarbonyl,
aminocarbonyl,
30 aminocarbonyl C₁-6 alkyl,
hydroxycarbonyl,
hydroxycarbonyl C₁-6 alkyl,
C₁-8 alkyl, either unsubstituted or substituted, with one or more
groups selected from: halogen, hydroxyl,

- 27 -

- C₁₋₅ alkylcarbonylamino, aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl, aminocarbonyl, C₁₋₅ alkylamino-
 carbonyl, C₁₋₅ alkylcarbonyloxy, C₃₋₈ cycloalkyl, oxo,
 amino, C₁₋₃ alkylamino, amino C₁₋₃ alkyl, arylamino-
 carbonyl, aryl C₁₋₅ alkylaminocarbonyl, aminocarbonyl,
 aminocarbonyl C₁₋₄ alkyl, hydroxycarbonyl, or
 hydroxycarbonyl C₁₋₅ alkyl,
- 5 -(CH₂)_s C≡CH,
 -(CH₂)_s C≡C-C₁₋₆ alkyl,
 10 -(CH₂)_s C≡C-C₃₋₇ cycloalkyl,
 -(CH₂)_s C≡C-aryl,
 -(CH₂)_s C≡C-C₁₋₆ alkylaryl,
 -(CH₂)_s CH=CH₂,
 -(CH₂)_s CH=CH C₁₋₆ alkyl,
 15 -(CH₂)_s CH=CH-C₃₋₇ cycloalkyl,
 -(CH₂)_s CH=CH aryl,
 -(CH₂)_s CH=CH C₁₋₆ alkylaryl,
 -(CH₂)_s SO₂C₁₋₆ alkyl,
 -(CH₂)_s SO₂C₁₋₆ alkylaryl,
 20 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 C₁₋₆ alkylamino C₁₋₆ alkyl,
 arylamino,
 25 arylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylamino,
 aryl C₁₋₆ alkylamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 30 C₁₋₆ dialkylamino,
 C₁₋₆ dialkylamino C₁₋₆ alkyl,
 C₁₋₆ alkylaminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,
 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,

- 28 -

- arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxycarbonylamino,
5 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
10 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
15 C₁₋₈ alkylaminocarbonylamino,
C₁₋₈ alkylaminocarbonylamino C₁₋₆ alkyl,
arylaminocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₈ alkylaminocarbonylamino,
aryl C₁₋₈ alkylaminocarbonylamino C₁₋₆ alkyl,
20 aminosulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkylaminosulfonylamino,
C₁₋₈ alkylaminosulfonylamino C₁₋₆ alkyl,
arylaminosulfonylamino C₁₋₆ alkyl,
aryl C₁₋₈ alkylaminosulfonylamino,
25 aryl C₁₋₈ alkylaminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
30 aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,

- 29 -

aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
5 aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
C₁₋₈ alkylaminocarbonyl C₁₋₆ alkyl,
arylaminocarbonyl C₁₋₆ alkyl,
aryl C₁₋₈ alkylaminocarbonyl, or
10 aryl C₁₋₈ alkylaminocarbonyl C₁₋₆ alkyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
are attached is itself attached to no more than one heteroatom; and
provided further that the carbon atom to which R¹⁰ and R¹¹ are
15 attached is itself attached to no more than one heteroatom;

R¹² is selected from
hydrogen,
C₁₋₈ alkyl,
20 aryl,
aryl C₁₋₈ alkyl,
hydroxy,
C₁₋₈ alkoxy,
aryloxy,
25 aryl C₁₋₆ alkoxy,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
C₁₋₈ alkylaminocarbonylmethyleneoxy, or
C₁₋₈ dialkylaminocarbonylmethyleneoxy;


30
m is an integer from 0 to 3;
n is an integer from 1 to 3;
p is an integer from 1 to 4;
q is an integer from 0 to 2;

- 30 -


r is an integer from 0 to 6;
 s is an integer from 0 to 3; and
 t is an integer from 0 to 3;

5 and the pharmaceutically acceptable salts thereof.

In a third embodiment of the invention is a method of eliciting an $\alpha\text{v}\beta 3$ antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound of the formula



wherein X is selected from




15 a 5- or 6-membered monocyclic aromatic or nonaromatic ring system containing 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S wherein the 5- or 6-membered ring system is either unsubstituted or substituted on a carbon atom with R^1 and R^2 , or

20 a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted on a carbon atom with R^1 and R^2 ;


Y is selected from

- 31 -




Z is

- 5 a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; provided that Z is not a 6-membered monocyclic aromatic ring system; preferably Z is selected from



10

- 32 -



5

R^1 , R^2 , R^4 , R^5 , R^{13} and R^{14} are each independently selected from hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, aryl, aryl C₁-8 alkyl, amino, amino C₁-8 alkyl, C₁-3 acylamino, C₁-3 acylamino C₁-8 alkyl, C₁-6 alkylamino, C₁-6 alkylamino-C₁-8 alkyl, C₁-6 dialkylamino, C₁-6 dialkylamino C₁-8 alkyl,

10

- 33 -

- 5 C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl,
hydroxycarbonyl C₁-6 alkyl, C₁-3 alkoxycarbonyl,
C₁-3 alkoxycarbonyl C₁-6 alkyl, hydroxycarbonyl-
C₁-6 alkyloxy, hydroxy, hydroxy C₁-6 alkyl, C₁-6 alkyloxy-
C₁-6 alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,
trifluoroethoxy, C₁-8 alkyl-S(O)_q, C₁-8 aminocarbonyl,
C₁-8 dialkylaminocarbonyl, C₁-8 alkyloxycarbonylamino,
C₁-8 alkylaminocarbonyloxy or C₁-8alkylsulfonylamino;
- 10 R³ is selected from
hydrogen,
aryl,
-(CH₂)_p-aryl,
hydroxyl,
15 C₁-5 alkoxycarbonyl,
aminocarbonyl,
C₃-8 cycloalkyl,
amino C₁-6 alkyl,
arylamino carbonyl,
20 aryl C₁-5 alkylaminocarbonyl,
hydroxycarbonyl C₁-6 alkyl,
C₁-8 alkyl,
aryl C₁-6 alkyl,
C₁-6 alkylamino C₁-6 alkyl,
25 aryl C₁-6 alkylamino C₁-6 alkyl,
C₁-6 dialkylamino C₁-6 alkyl,
C₁-8 alkylsulfonyl,
C₁-8 alkoxycarbonyl,
aryloxycarbonyl,
30 aryl C₁-8 alkoxycarbonyl,
C₁-8 alkylcarbonyl,
arylcarbonyl,
aryl C₁-6 alkylcarbonyl,
C₁-8 alkylaminocarbonyl,

- 34 -

- 5 aminosulfonyl,
 C1-8 alkylaminosulfonyl,
 arylaminosulfonylamino,
 aryl C1-8 alkylaminosulfonyl,
 C1-6 alkylsulfonyl,
 arylsulfonyl,
 aryl C1-6 alkylsulfonyl,
 aryl C1-6 alkylcarbonyl,
 C1-6 alkylthiocarbonyl,
10 arylthiocarbonyl, or
 aryl C1-6 alkylthiocarbonyl,
 wherein any of the alkyl groups may be unsubstituted or substituted with
 R13 and R14;
- 15 R6, R7, R8, R9, R10 and R11 are each independently selected from
 hydrogen,
 aryl,
 -(CH2)_p-aryl,
 halogen,
20 hydroxyl,
 C1-8 alkylcarbonylamino,
 aryl C1-5 alkoxy,
 C1-5 alkoxy carbonyl,
 aminocarbonyl,
25 C1-8 alkylaminocarbonyl,
 C1-6 alkylcarbonyloxy,
 C3-8 cycloalkyl,
 oxo,
 amino,
30 C1-6 alkylamino,
 amino C1-6 alkyl,
 arylaminocarbonyl,
 aryl C1-5 alkylaminocarbonyl,
 aminocarbonyl,

- 35 -

- aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 C₁₋₈ alkyl, either unsubstituted or substituted, with one or more
 5 groups selected from: halogen, hydroxyl,
 C₁₋₅ alkylcarbonylamino, aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxy carbonyl, aminocarbonyl, C₁₋₅ alkylamino-
 carbonyl, C₁₋₅ alkylcarbonyloxy, C₃₋₈ cycloalkyl, oxo,
 amino, C₁₋₃ alkylamino, amino C₁₋₃ alkyl, arylamino-
 10 carbonyl, aryl C₁₋₅ alkylaminocarbonyl, aminocarbonyl,
 aminocarbonyl C₁₋₄ alkyl, hydroxycarbonyl, or
 hydroxycarbonyl C₁₋₅ alkyl,
 -(CH₂)_s C≡CH,
 -(CH₂)_s C≡C-C₁₋₆ alkyl,
 15 -(CH₂)_s C≡C-C₃₋₇ cycloalkyl,
 -(CH₂)_s C≡C-aryl,
 -(CH₂)_s C≡C-C₁₋₆ alkylaryl,
 -(CH₂)_s CH=CH₂,
 -(CH₂)_s CH=CH C₁₋₆ alkyl,
 20 -(CH₂)_s CH=CH-C₃₋₇ cycloalkyl,
 -(CH₂)_s CH=CH aryl,
 -(CH₂)_s CH=CH C₁₋₆ alkylaryl,
 -(CH₂)_s SO₂C₁₋₆ alkyl,
 -(CH₂)_s SO₂C₁₋₆ alkylaryl,
 25 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 C₁₋₆ alkylamino C₁₋₆ alkyl,
 arylamino,
 30 arylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylamino,
 aryl C₁₋₆ alkylamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,

- 36 -

- C1-6 dialkylamino,
C1-6 dialkylamino C1-6 alkyl,
C1-6 alkylaminocarbonyloxy,
C1-8 alkylsulfonylamino,
5 C1-8 alkylsulfonylamino C1-6 alkyl,
arylsulfonylamino C1-6 alkyl,
aryl C1-6 alkylsulfonylamino,
aryl C1-6 alkylsulfonylamino C1-6 alkyl,
C1-8 alkoxycarbonylamino,
10 C1-8 alkoxycarbonylamino C1-8 alkyl,
aryloxycarbonylamino C1-8 alkyl,
aryl C1-8 alkoxycarbonylamino,
aryl C1-8 alkoxycarbonylamino C1-8 alkyl,
C1-8 alkylcarbonylamino,
15 C1-8 alkylcarbonylamino C1-6 alkyl,
arylcarbonylamino C1-6 alkyl,
aryl C1-6 alkylcarbonylamino,
aryl C1-6 alkylcarbonylamino C1-6 alkyl,
aminocarbonylamino C1-6 alkyl,
20 C1-8 alkylaminocarbonylamino,
C1-8 alkylaminocarbonylamino C1-6 alkyl,
arylaminocarbonylamino C1-6 alkyl,
aryl C1-8 alkylaminocarbonylamino,
aryl C1-8 alkylaminocarbonylamino C1-6 alkyl,
25 aminosulfonylamino C1-6 alkyl,
C1-8 alkylaminosulfonylamino,
C1-8 alkylaminosulfonylamino C1-6 alkyl,
arylaminosulfonylamino C1-6 alkyl,
aryl C1-8 alkylaminosulfonylamino,
30 aryl C1-8 alkylaminosulfonylamino C1-6 alkyl,
C1-6 alkylsulfonyl,
C1-6 alkylsulfonyl C1-6 alkyl,
arylsulfonyl C1-6 alkyl,
aryl C1-6 alkylsulfonyl,

- 37 -

aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcarbonyl C₁₋₆ alkyl,
5 aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
10 aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
C₁₋₈ alkylaminocarbonyl C₁₋₆ alkyl,
arylaminocarbonyl C₁₋₆ alkyl,
aryl C₁₋₈ alkylaminocarbonyl, or
15 aryl C₁₋₈ alkylaminocarbonyl C₁₋₆ alkyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
are attached is itself attached to no more than one heteroatom; and
provided further that the carbon atom to which R¹⁰ and R¹¹ are
20 attached is itself attached to no more than one heteroatom;

R¹² is selected from
hydrogen,
C₁₋₈ alkyl,
25 aryl,
aryl C₁₋₈ alkyl,
hydroxy,
C₁₋₈ alkoxy,
aryloxy,
30 aryl C₁₋₆ alkoxy,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
C₁₋₈ alkylaminocarbonylmethyleneoxy, or
C₁₋₈ dialkylaminocarbonylmethyleneoxy;

- 38 -

m is an integer from 0 to 3;

n is an integer from 1 to 3;

p is an integer from 1 to 4;

5 q is an integer from 0 to 2;

r is an integer from 0 to 6; and


s is an integer from 0 to 3;

and the pharmaceutically acceptable salts thereof.

10 Illustrating the invention is the method wherein the $\alpha v\beta 3$ antagonizing effect is selected from inhibition of bone resorption, inhibition of restenosis, inhibition of angiogenesis, inhibition of diabetic retinopathy, inhibition of macular degeneration, inhibition of

15 atherosclerosis, inflammation or inhibition of tumor growth. Preferably, the $\alpha v\beta 3$ antagonizing effect is the inhibition of bone resorption.

An illustration of the invention is a compound of the formula




20 wherein X is

a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups chosen from R¹, R², R¹⁵ or R¹⁶;

25

Y is selected from

- 39 -



Z is

5 a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; provided that Z is not a 6-membered monocyclic aromatic ring system, an isoxazoline ring or an isoxazole ring;

10 R¹, R², R⁴, R⁵, R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from

15 hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, C₃-8 cycloheteroalkyl, C₃-8 cycloalkyl C₁-6 alkyl, C₃-8 cycloheteroalkyl C₁-6 alkyl, aryl, aryl C₁-8 alkyl, amino, amino C₁-8 alkyl, C₁-3 acylamino, C₁-3 acylamino C₁-8 alkyl, (C₁-6 alkyl)_qamino, (C₁-6 alkyl)_qamino C₁-8 alkyl, C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl,

20 hydroxycarbonyl C₁-6 alkyl, C₁-3 alkoxy carbonyl, C₁-3 alkoxy carbonyl C₁-6 alkyl, hydroxycarbonyl-C₁-6 alkyloxy, hydroxy, hydroxy C₁-6 alkyl, C₁-6 alkyloxy-C₁-6 alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,

- 40 -

trifluoroethoxy, C₁₋₈ alkyl-S(O)_q, (C₁₋₈ alkyl)_qaminocarbonyl,
C₁₋₈ alkyloxycarbonylamino, (C₁₋₈ alkyl)_qaminocarbonyloxy,
oxo, (aryl C₁₋₈ alkyl)_qamino, (aryl)_qamino,
aryl C₁₋₈ alkylsulfonylamino or C₁₋₈ alkylsulfonylamino;

5

R³ is selected from

hydrogen,

aryl,

aryl-(CH₂)_p-,

10

hydroxyl,

C₁₋₅ alkoxy,

aminocarbonyl,

C₃₋₈ cycloalkyl,

amino C₁₋₆ alkyl,

15

(aryl)_qaminocarbonyl,

(aryl C₁₋₅ alkyl)_qaminocarbonyl,

hydroxycarbonyl C₁₋₆ alkyl,

C₁₋₈ alkyl,

aryl C₁₋₆ alkyl,

20

(C₁₋₆ alkyl)_qamino C₁₋₆ alkyl,

(aryl C₁₋₆ alkyl)_qamino C₁₋₆ alkyl,

C₁₋₈ alkylsulfonyl,

C₁₋₈ alkoxycarbonyl,

aryloxycarbonyl,

25

aryl C₁₋₈ alkoxycarbonyl,

C₁₋₈ alkylcarbonyl,

arylcarbonyl,

aryl C₁₋₆ alkylcarbonyl,

(C₁₋₈ alkyl)_qaminocarbonyl,

30

aminosulfonyl,

C₁₋₈ alkylaminosulfonyl,

(aryl)_qaminosulfonylamino,

(aryl C₁₋₈ alkyl)_qaminosulfonyl,

C₁₋₆ alkylsulfonyl,

- 41 -

- arylsulfonyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylthiocarbonyl,
5 arylthiocarbonyl, or
aryl C₁₋₆ alkylthiocarbonyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴;
- 10 R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from
hydrogen,
aryl,
aryl-(CH₂)_p-,
15 aryl-(CH₂)_n-O-(CH₂)_m-,
aryl-(CH₂)_n-S(O)_q-(CH₂)_m-,
aryl-(CH₂)_n-C(O)-(CH₂)_m-,
aryl-(CH₂)_n-C(O)-N(R³)-(CH₂)_m-,
aryl-(CH₂)_n-N(R³)-C(O)-(CH₂)_m-,
20 aryl-(CH₂)_n-N(R³)-(CH₂)_m-,
halogen,
hydroxyl,
C₁₋₈ alkylcarbonylamino,
aryl C₁₋₅ alkoxy,
C₁₋₅ alkoxycarbonyl,
25 (C₁₋₈ alkyl)_qaminocarbonyl,
- C₁₋₆ alkylcarbonyloxy,
C₃₋₈ cycloalkyl,
oxo,
(C₁₋₆ alkyl)_qamino,
30 amino C₁₋₆ alkyl,
arylaminocarbonyl,
aryl C₁₋₅ alkylaminocarbonyl,
aminocarbonyl,
aminocarbonyl C₁₋₆ alkyl,

- 42 -

- hydroxycarbonyl,
hydroxycarbonyl C1-6 alkyl,
C1-8 alkyl, either unsubstituted or substituted, with one or more
groups selected from: halogen, hydroxyl,
5 C1-5 alkylcarbonylamino, aryl C1-5 alkoxy,
C1-5 alkoxycarbonyl, aminocarbonyl, (C1-5 alkyl)qamino-
carbonyl, C1-5 alkylcarbonyloxy, C3-8 cycloalkyl, oxo,
(C1-3 alkyl)qamino, amino C1-3 alkyl,
(aryl)qaminocarbonyl, (aryl C1-5 alkyl)qaminocarbonyl,
10 aminocarbonyl, aminocarbonyl C1-4 alkyl,
hydroxycarbonyl or hydroxycarbonyl C1-5 alkyl,
 $\text{CH}\equiv\text{C}-(\text{CH}_2)_{\text{s-}}$,
C1-6 alkyl- $\text{C}\equiv\text{C}-(\text{CH}_2)_{\text{s-}}$,
C3-7 cycloalkyl- $\text{C}\equiv\text{C}-(\text{CH}_2)_{\text{s-}}$,
15 aryl- $\text{C}\equiv\text{C}-(\text{CH}_2)_{\text{s-}}$,
C1-6 alkylaryl- $\text{C}\equiv\text{C}-(\text{CH}_2)_{\text{s-}}$,
 $\text{CH}_2=\text{CH}-(\text{CH}_2)_{\text{s-}}$,
C1-6 alkyl- $\text{CH}=\text{CH}-(\text{CH}_2)_{\text{s-}}$,
C3-7 cycloalkyl- $\text{CH}=\text{CH}-(\text{CH}_2)_{\text{s-}}$,
20 aryl- $\text{CH}=\text{CH}-(\text{CH}_2)_{\text{s-}}$,
C1-6 alkylaryl- $\text{CH}=\text{CH}-(\text{CH}_2)_{\text{s-}}$,
C1-6 alkyl- $\text{SO}_2-(\text{CH}_2)_{\text{s-}}$,
C1-6 alkylaryl- $\text{SO}_2-(\text{CH}_2)_{\text{s-}}$,
C1-6 alkoxy,
25 aryl C1-6 alkoxy,
aryl C1-6 alkyl,
(C1-6 alkyl)qamino C1-6 alkyl,
(aryl)qamino,
(aryl)qamino C1-6 alkyl,
30 (aryl C1-6 alkyl)qamino,
(aryl C1-6 alkyl)qamino C1-6 alkyl,
arylcarbonyloxy,
aryl C1-6 alkylcarbonyloxy,
(C1-6 alkyl)qaminocarbonyloxy,

- 43 -

- C₁-8 alkylsulfonylamino,
arylsulfonylamino,
C₁-8 alkylsulfonylamino C₁-6 alkyl,
arylsulfonylamino C₁-6 alkyl,
5 aryl C₁-6 alkylsulfonylamino,
aryl C₁-6 alkylsulfonylamino C₁-6 alkyl,
C₁-8 alkoxycarbonylamino,
C₁-8 alkoxycarbonylamino C₁-8 alkyl,
aryloxycarbonylamino C₁-8 alkyl,
10 aryl C₁-8 alkoxycarbonylamino,
aryl C₁-8 alkoxycarbonylamino C₁-8 alkyl,
C₁-8 alkylcarbonylamino,
C₁-8 alkylcarbonylamino C₁-6 alkyl,
arylcarbonylamino C₁-6 alkyl,
15 aryl C₁-6 alkylcarbonylamino,
aryl C₁-6 alkylcarbonylamino C₁-6 alkyl,
aminocarbonylamino C₁-6 alkyl,
(C₁-8 alkyl)qaminocarbonylamino,
(C₁-8 alkyl)qaminocarbonylamino C₁-6 alkyl,
20 (aryl)qaminocarbonylamino C₁-6 alkyl,
(aryl C₁-8 alkyl)qaminocarbonylamino,
(aryl C₁-8 alkyl)qaminocarbonylamino C₁-6 alkyl,
aminosulfonylamino C₁-6 alkyl,
(C₁-8 alkyl)qaminosulfonylamino,
25 (C₁-8 alkyl)qaminosulfonylamino C₁-6 alkyl,
(aryl)qaminosulfonylamino C₁-6 alkyl,
(aryl C₁-8 alkyl)qaminosulfonylamino,
(aryl C₁-8 alkyl)qaminosulfonylamino C₁-6 alkyl,
C₁-6 alkylsulfonyl,
30 C₁-6 alkylsulfonyl C₁-6 alkyl,
arylsulfonyl C₁-6 alkyl,
aryl C₁-6 alkylsulfonyl,
aryl C₁-6 alkylsulfonyl C₁-6 alkyl,
C₁-6 alkylcarbonyl,

- 44 -


- C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 arylcarbonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 5 C₁₋₆ alkylthiocarbonylamino,
 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 arylthiocarbonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylthiocarbonylamino,
 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 10 (C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,
 (aryl)_qaminocarbonyl C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_qaminocarbonyl, or
 (aryl C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,
 wherein any of the alkyl groups may be unsubstituted or substituted with
 15 R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
 are attached is itself attached to no more than one heteroatom; and
 provided further that the carbon atom to which R¹⁰ and R¹¹ are
 attached is itself attached to no more than one heteroatom;
- 20 R¹² is selected from
 hydrogen,
 C₁₋₈ alkyl,
 aryl,
 aryl C₁₋₈ alkyl,
 25 C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
 aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
 C₁₋₈ alkylaminocarbonylmethylene, or
 C₁₋₈ dialkylaminocarbonylmethylene;
- 30 m, s and t are each independently an integer from 0 to 3;
 n is an integer from 1 to 3;
 p is an integer from 1 to 4;
 q is an integer from 0 to 2;
 r is an integer from 0 to 6;

- 45 -





and the pharmaceutically acceptable salts thereof.

- 5 Particularly illustrative of the invention is the compound wherein Z is a 5-11 membered nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; and all other variables are as defined above.


- 10 Exemplifying the invention is the compound wherein Z is selected from




- 46 -



5 and the pharmaceutically acceptable salts thereof. Preferably Z is selected from




- 47 -




- 48 -

An example of the invention is the compound of the formula



wherein X is selected from



Y is selected from $-(\text{CH}_2)_r-$ or $-(\text{CH}_2)_m-\text{NR}^3-(\text{CH}_2)_t-$;

R³ is selected from

- hydrogen,
 aryl- $(\text{CH}_2)_p-$,
 10 C₁-5 alkoxy carbonyl,
 C₃-8 cycloalkyl,
 (aryl)_qaminocarbonyl,
 (aryl C₁-5 alkyl)_qaminocarbonyl,
 C₁-8 alkyl,
 15 aryl C₁-6 alkyl,
 C₁-8 alkylsulfonyl,
 arylsulfonyl,
 aryl C₁-6 alkylsulfonyl,
 C₁-8 alkoxy carbonyl,
 20 aryloxy carbonyl,
 aryl C₁-8 alkoxy carbonyl,
 C₁-8 alkylcarbonyl,
 arylcarbonyl,
 aryl C₁-6 alkylcarbonyl,
 25 (C₁-8 alkyl)_qaminocarbonyl,
 C₁-6 alkylsulfonyl, or
 aryl C₁-6 alkylcarbonyl,

- 49 -

wherein any of the alkyl groups may be unsubstituted or substituted with R^{13} and R^{14} ;

R^4 is selected from

5 hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloheteroalkyl, C₃₋₈ cycloalkyl C₁₋₆ alkyl, C₃₋₈ cycloheteroalkyl C₁₋₆ alkyl, aryl or aryl C₁₋₈ alkyl,

R^8 is selected from

10 hydrogen,
aryl,
aryl-(CH₂)_p-,
CH≡C-(CH₂)_s-,
C₁₋₆ alkyl-C≡C-(CH₂)_s-,
15 C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,
aryl-C≡C-(CH₂)_s-,
C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,
CH₂=CH-(CH₂)_s-,
C₁₋₆ alkyl-CH=CH-(CH₂)_s-,
20 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,
aryl-CH=CH-(CH₂)_s-,
C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,
C₁₋₆ alkyl-SO₂-(CH₂)_s-,
C₁₋₆ alkylaryl-SO₂-(CH₂)_s-; and

25


r is an integer from 0 to 3;

wherein all other variables are as defined above;

and the pharmaceutically acceptable salts thereof.


Further illustrating the invention is the compound of the

30 formula




- 50 -

wherein Z is selected from



R⁸ is selected from
hydrogen,



5

-indolyl-(CH₂)_p-,CH≡C-(CH₂)_s-,C₁₋₆ alkyl-C≡C-(CH₂)_s-,C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,

10

aryl-C≡C-(CH₂)_s-,C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,CH₂=CH-(CH₂)_s-,C₁₋₆ alkyl-CH=CH-(CH₂)_s-,C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,

15

aryl-CH=CH-(CH₂)_s-,

- 51 -

C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,
C₁₋₆ alkyl-SO₂-(CH₂)_s-,
C₁₋₆ alkylaryl-SO₂-(CH₂)_s-; and

- 5 R¹² is selected from hydrogen or C₁₋₈ alkyl; and
s is an integer from 0 to 3;
and all other variables are as defined above;
and the pharmaceutically acceptable salts thereof.

Further exemplifying the invention is the compound
10 selected from

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperidin-
1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

- 15 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperin-
1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine trifluoroacetate;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;

20

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

- 25 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-
pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;

- 30 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

- 52 -

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine ethyl ester;

- 5 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

10 Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

- 15 Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;

- 20 Ethyl 2-oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;

25 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;

Ethyl 3-(2-{2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}-pyrrolidin-1-yl)-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;

- 30 3-(2-{2-Oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}pyrrolidin-1-yl)-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;

- 53 -

3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-pyridin-3-yl-propionic acid;

- 5 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-pyridin-3-yl-propionic acid;

- 10 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine ethyl ester;

2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;

- 15 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine ethyl ester;

2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

- 20 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine ethyl ester;

- 25 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;

- 30 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-acetylamino)-propionic acid ethyl ester; or

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-acetylamino)-propionic acid;

- 54 -

and the pharmaceutically acceptable salts thereof.

Preferably, the compound is selected from

- 5 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine trifluoroacetate;
- 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;
- 10 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;
- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;
- 15 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;
- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;
- 20 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3-(S)-pyridin-3-yl- β -alanine;
- 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3-(S)-pyridin-3-yl- β -alanine;
- 25 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;
- 3-(2-{ 2-Oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}pyrrolidin-1-yl)-acetyl-amino)-3-(S)-pyridin-3-yl-propionic acid;
- 30 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetyl-amino}-3-(S)-pyridin-3-yl-propionic acid; or

- 55 -

3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-
pyridin-3-yl-propionic acid;

5

2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;


10 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;

5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine; or

15 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-
tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-
acetylamino)-propionic acid;

and the pharmaceutically acceptable salts thereof.

20 An additional example of the invention is a compound of
the formula




wherein X is

25 a 9- to 10-membered polycyclic ring system, wherein one or
more of the rings is aromatic, and wherein the polycyclic ring
system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or
S, and wherein the polycyclic ring system is either unsubstituted
or substituted on a carbon atom with R¹ and R²;

30 Y is selected from


- 56 -




Z is

5 a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; provided that Z is not a 6-membered monocyclic aromatic ring system; preferably, Z is selected from

10



- 57 -



5 R¹, R², R³, R⁴, R⁵, R¹³ and R¹⁴ are each independently selected from hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, aryl, aryl C₁-8 alkyl, amino, amino C₁-8 alkyl, C₁-3 acylamino, C₁-3 acylamino C₁-8 alkyl, C₁-6 alkylamino, C₁-6 alkylamino-C₁-8 alkyl, C₁-6 dialkylamino, C₁-6 dialkylamino C₁-8 alkyl, C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl,

- 58 -

hydroxycarbonyl C₁₋₆ alkyl, C₁₋₃ alkoxy carbonyl,
C₁₋₃ alkoxy carbonyl C₁₋₆ alkyl, hydroxycarbonyl-
C₁₋₆ alkyloxy, hydroxy or hydroxy C₁₋₆ alkyl;

- 5 R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from
hydrogen,
aryl,
-(CH₂)_p-aryl,
halogen,
10 hydroxyl,
C₁₋₈ alkylcarbonylamino,
aryl C₁₋₅ alkoxy,
C₁₋₅ alkoxy carbonyl,
aminocarbonyl,
15 C₁₋₈ alkylaminocarbonyl,
C₁₋₆ alkylcarbonyloxy,
C₃₋₈ cycloalkyl,
oxo,
amino,
20 C₁₋₆ alkylamino,
amino C₁₋₆ alkyl,
arylaminocarbonyl,
aryl C₁₋₅ alkylaminocarbonyl,
aminocarbonyl,
25 aminocarbonyl C₁₋₆ alkyl,
hydroxycarbonyl,
hydroxycarbonyl C₁₋₆ alkyl,
C₁₋₈ alkyl, either unsubstituted or substituted, with one or more
groups selected from: halogen, hydroxyl,
30 C₁₋₅ alkylcarbonylamino, aryl C₁₋₅ alkoxy,
C₁₋₅ alkoxy carbonyl, aminocarbonyl, C₁₋₅ alkylamino-
carbonyl, C₁₋₅ alkylcarbonyloxy, C₃₋₈ cycloalkyl, oxo,
amino, C₁₋₃ alkylamino, amino C₁₋₃ alkyl, arylamino-
carbonyl, aryl C₁₋₅ alkylaminocarbonyl, aminocarbonyl,

- 59 -

- aminocarbonyl C₁₋₄ alkyl, hydroxycarbonyl, or
hydroxycarbonyl C₁₋₅ alkyl,
- 5 -(CH₂)_s C≡CH,
 -(CH₂)_s C≡C-C₁₋₆ alkyl,
 -(CH₂)_s C≡C-C₃₋₇ cycloalkyl,
 -(CH₂)_s C≡C-aryl,
 -(CH₂)_s C≡C-C₁₋₆ alkylaryl,
 -(CH₂)_s CH=CH₂,
10 -(CH₂)_s CH=CH C₁₋₆ alkyl,
 -(CH₂)_s CH=CH-C₃₋₇ cycloalkyl,
 -(CH₂)_s CH=CH aryl,
 -(CH₂)_s CH=CH C₁₋₆ alkylaryl,
 -(CH₂)_s SO₂C₁₋₆ alkyl, or
 -(CH₂)_s SO₂C₁₋₆ alkylaryl;
15 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 C₁₋₆ alkylamino C₁₋₆ alkyl,
 arylamino,
20 arylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylamino,
 aryl C₁₋₆ alkylamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
25 C₁₋₆ dialkylamino,
 -C₁₋₆ dialkylamino C₁₋₆ alkyl,
 C₁₋₆ alkylaminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,
 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
30 arylsulfonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylsulfonylamino,
 aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
 C₁₋₈ alkoxycarbonylamino,
 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,

- 60 -

- aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
5 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
10 C₁₋₈ alkylaminocarbonylamino,
C₁₋₈ alkylaminocarbonylamino C₁₋₆ alkyl,
arylaminocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₈ alkylaminocarbonylamino,
aryl C₁₋₈ alkylaminocarbonylamino C₁₋₆ alkyl,
15 aminosulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkylaminosulfonylamino,
C₁₋₈ alkylaminosulfonylamino C₁₋₆ alkyl,
arylaminosulfonylamino C₁₋₆ alkyl,
aryl C₁₋₈ alkylaminosulfonylamino,
20 aryl C₁₋₈ alkylaminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
25 aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
30 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,

- 61 -


aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
C₁₋₈ alkylaminocarbonyl C₁₋₆ alkyl,
arylamino carbonyl C₁₋₆ alkyl,
aryl C₁₋₈ alkylaminocarbonyl, or
5 aryl C₁₋₈ alkylaminocarbonyl C₁₋₆ alkyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
are attached is itself attached to no more than one heteroatom; and
provided further that the carbon atom to which R¹⁰ and R¹¹ are
10 attached is itself attached to no more than one heteroatom;

R¹² is selected from
hydrogen,
C₁₋₈ alkyl,
15 aryl,
aryl C₁₋₈ alkyl,
hydroxy,
C₁₋₈ alkoxy,
aryloxy,
20 aryl C₁₋₆ alkoxy,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
C₁₋₈ alkylaminocarbonylmethyleneoxy, or
C₁₋₈ dialkylaminocarbonylmethyleneoxy;

25 m is an integer from 0 to 3;
n is an integer from 1 to 3;
p is an integer from 1 to 4;
q is an integer from 0 to 2;
30 r is an integer from 0 to 6; and
s is an integer from 0 to 3;
and the pharmaceutically acceptable salts thereof.

An additional illustration of the invention is a compound of
the formula


- 62 -



wherein X is

5 a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted on a carbon atom with R¹ and R²;


Y is selected from




Z is

15 a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted on a carbon or nitrogen atom with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; provided that Z is not a 6-membered monocyclic aromatic ring system; preferably, Z is selected from

- 63 -



- 64 -



5 R¹, R², R⁴, R⁵, R¹³ and R¹⁴ are each independently selected from
 hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, aryl,
 aryl C₁-8 alkyl, amino, amino C₁-8 alkyl, C₁-3 acylamino,
 C₁-3 acylamino C₁-8 alkyl, C₁-6 alkylamino, C₁-6 alkylamino-
 C₁-8 alkyl, C₁-6 dialkylamino, C₁-6 dialkylamino C₁-8 alkyl,
 C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl,
 hydroxycarbonyl C₁-6 alkyl, C₁-3 alkoxy carbonyl,
 10 C₁-3 alkoxy carbonyl C₁-6 alkyl, hydroxycarbonyl-
 C₁-6 alkyloxy, hydroxy, hydroxy C₁-6 alkyl, C₁-6 alkyloxy-
 C₁-6 alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,
 trifluoroethoxy, C₁-8 alkyl-S(O)_q, C₁-8 aminocarbonyl,
 C₁-8 dialkylaminocarbonyl, C₁-8 alkyloxycarbonylamino,
 15 C₁-8 alkylaminocarbonyloxy or C₁-8 alkylsulfonylamino;

R³ is selected from
 hydrogen,
 aryl,
 20 -(CH₂)_p-aryl,
 hydroxyl,

- 65 -

- 5 C₁-5 alkoxycarbonyl,
aminocarbonyl,
C₃-8 cycloalkyl,
amino C₁-6 alkyl,
arylamino carbonyl,
aryl C₁-5 alkylaminocarbonyl,
hydroxycarbonyl C₁-6 alkyl,
C₁-8 alkyl,
aryl C₁-6 alkyl,
10 C₁-6 alkylamino C₁-6 alkyl,
aryl C₁-6 alkylamino C₁-6 alkyl,
C₁-6 dialkylamino C₁-6 alkyl,
C₁-8 alkylsulfonyl,
C₁-8 alkoxycarbonyl,
15 aryloxycarbonyl,
aryl C₁-8 alkoxycarbonyl,
C₁-8 alkylcarbonyl,
arylcarbonyl,
aryl C₁-6 alkylcarbonyl,
20 C₁-8 alkylaminocarbonyl,
aminosulfonyl,
C₁-8 alkylaminosulfonyl,
arylaminosulfonylamino,
aryl C₁-8 alkylaminosulfonyl,
25 C₁-6 alkylsulfonyl,
arylsulfonyl,
aryl C₁-6 alkylsulfonyl,
aryl C₁-6 alkylcarbonyl,
C₁-6 alkylthiocarbonyl,
30 arylthiocarbonyl, or
aryl C₁-6 alkylthiocarbonyl,

wherein any of the alkyl groups may be unsubstituted or substituted with R¹³ and R¹⁴;

- 66 -

R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from

hydrogen,

aryl,

-(CH₂)_p-aryl,

5 halogen,

hydroxyl,

C₁₋₈ alkylcarbonylamino,

aryl C₁₋₅ alkoxy,

C₁₋₅ alkoxy carbonyl,

10 aminocarbonyl,

C₁₋₈ alkylaminocarbonyl,

C₁₋₆ alkylcarbonyloxy,

C₃₋₈ cycloalkyl,

oxo,

15 amino,

C₁₋₆ alkylamino,

amino C₁₋₆ alkyl,

arylaminocarbonyl,

aryl C₁₋₅ alkylaminocarbonyl,

20 aminocarbonyl,

aminocarbonyl C₁₋₆ alkyl,

hydroxycarbonyl,

hydroxycarbonyl C₁₋₆ alkyl,

C₁₋₈ alkyl, either unsubstituted or substituted, with one or more

25 groups selected from: halogen, hydroxyl,

C₁₋₅ alkylcarbonylamino, aryl C₁₋₅ alkoxy,

C₁₋₅ alkoxy carbonyl, aminocarbonyl, C₁₋₅ alkylamino-

carbonyl, C₁₋₅ alkylcarbonyloxy, C₃₋₈ cycloalkyl, oxo,

amino, C₁₋₃ alkylamino, amino C₁₋₃ alkyl, arylamino-

30 carbonyl, aryl C₁₋₅ alkylaminocarbonyl, aminocarbonyl,

aminocarbonyl C₁₋₄ alkyl, hydroxycarbonyl, or

hydroxycarbonyl C₁₋₅ alkyl,

-(CH₂)_s C≡CH,

-(CH₂)_s C≡C-C₁₋₆ alkyl,

- 67 -

- 5 -(CH₂)_s C≡C-C₃₋₇ cycloalkyl,
 -(CH₂)_s C≡C-aryl,
 -(CH₂)_s C≡C-C₁₋₆ alkylaryl,
 -(CH₂)_s CH=CH₂,
 -(CH₂)_s CH=CH C₁₋₆ alkyl,
 -(CH₂)_s CH=CH-C₃₋₇ cycloalkyl,
 -(CH₂)_s CH=CH aryl,
 -(CH₂)_s CH=CH C₁₋₆ alkylaryl,
10 -(CH₂)_s SO₂C₁₋₆ alkyl, or
 -(CH₂)_s SO₂C₁₋₆ alkylaryl;
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 C₁₋₆ alkylamino C₁₋₆ alkyl,
15 arylamino,
 arylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylamino,
 aryl C₁₋₆ alkylamino C₁₋₆ alkyl,
 arylcarbonyloxy,
20 aryl C₁₋₆ alkylcarbonyloxy,
 C₁₋₆ dialkylamino,
 C₁₋₆ dialkylamino C₁₋₆ alkyl,
 C₁₋₆ alkylaminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,
25 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
 arylsulfonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylsulfonylamino,
 aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
 C₁₋₈ alkoxycarbonylamino,
30 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
 aryloxycarbonylamino C₁₋₈ alkyl,
 aryl C₁₋₈ alkoxycarbonylamino,
 aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
 C₁₋₈ alkylcarbonylamino,

- 68 -

- 5 C1-8 alkylcarbonylamino C1-6 alkyl,
arylcarbonylamino C1-6 alkyl,
aryl C1-6 alkylcarbonylamino,
aryl C1-6 alkylcarbonylamino C1-6 alkyl,
aminocarbonylamino C1-6 alkyl,
C1-8 alkylaminocarbonylamino,
C1-8 alkylaminocarbonylamino C1-6 alkyl,
arylaminocarbonylamino C1-6 alkyl,
10 aryl C1-8 alkylaminocarbonylamino,
aryl C1-8 alkylaminocarbonylamino C1-6 alkyl,
aminosulfonylamino C1-6 alkyl,
C1-8 alkylaminosulfonylamino,
C1-8 alkylaminosulfonylamino C1-6 alkyl,
15 arylaminosulfonylamino C1-6 alkyl,
aryl C1-8 alkylaminosulfonylamino,
aryl C1-8 alkylaminosulfonylamino C1-6 alkyl,
C1-6 alkylsulfonyl,
C1-6 alkylsulfonyl C1-6 alkyl,
arylsulfonyl C1-6 alkyl,
20 aryl C1-6 alkylsulfonyl,
aryl C1-6 alkylsulfonyl C1-6 alkyl,
C1-6 alkylcarbonyl,
C1-6 alkylcarbonyl C1-6 alkyl,
arylcarbonyl C1-6 alkyl,
25 aryl C1-6 alkylcarbonyl,
aryl C1-6 alkylcarbonyl C1-6 alkyl,
C1-6 alkylthiocarbonylamino,
C1-6 alkylthiocarbonylamino C1-6 alkyl,
arylthiocarbonylamino C1-6 alkyl,
30 aryl C1-6 alkylthiocarbonylamino,
aryl C1-6 alkylthiocarbonylamino C1-6 alkyl,
C1-8 alkylaminocarbonyl C1-6 alkyl,
arylaminocarbonyl C1-6 alkyl,
aryl C1-8 alkylaminocarbonyl, or

- 69 -

aryl C₁₋₈ alkylaminocarbonyl C₁₋₆ alkyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
are attached is itself attached to no more than one heteroatom; and
5 provided further that the carbon atom to which R¹⁰ and R¹¹ are
attached is itself attached to no more than one heteroatom;

R¹² is selected from

- hydrogen,
- 10 C₁₋₈ alkyl,
- aryl,
- aryl C₁₋₈ alkyl,
- hydroxy,
- C₁₋₈ alkoxy,
- 15 aryloxy,
- aryl C₁₋₆ alkoxy,
- C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
- aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkoxy,
- C₁₋₈ alkylaminocarbonylmethyleneoxy, or
- 20 C₁₋₈ dialkylaminocarbonylmethyleneoxy;

m is an integer from 0 to 3;

n is an integer from 1 to 3;

p is an integer from 1 to 4;

25 q is an integer from 0 to 2;

r is an integer from 0 to 6; and

s is an integer from 0 to 3;

and the pharmaceutically acceptable salts thereof.

More particularly illustrating the invention is a
30 pharmaceutical composition comprising any of the compounds described
above and a pharmaceutically acceptable carrier. Another example of
the invention is a pharmaceutical composition made by combining any
of the compounds described above and a pharmaceutically acceptable
carrier. Another illustration of the invention is a process for making a

- 70 -

pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

Further illustrating the invention is a method of treating and/or preventing a condition mediated by antagonism of a vitronectin receptor in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds described above. Preferably, the condition is selected from bone resorption, osteoporosis, restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, cancer and tumor growth. More preferably, the condition is selected from osteoporosis and cancer. Most preferably, the condition is osteoporosis.

More specifically exemplifying the invention is a method of eliciting a vitronectin antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds or any of the pharmaceutical compositions described above. Preferably, the vitronectin antagonizing effect is an $\alpha v\beta 3$ antagonizing effect; more specifically the $\alpha v\beta 3$ antagonizing effect is selected from inhibition of bone resorption, inhibition of restenosis, inhibition of atherosclerosis, inhibition of angiogenesis, inhibition of diabetic retinopathy, inhibition of macular degeneration, inhibition of inflammation or inhibition of tumor growth. Most preferably, the $\alpha v\beta 3$ antagonizing effect is inhibition of bone resorption. Alternatively, the vitronectin antagonizing effect is an $\alpha v\beta 5$ antagonizing effect or a dual $\alpha v\beta 3/\alpha v\beta 5$ antagonizing effect. Examples of $\alpha v\beta 5$ antagonizing effects are inhibition of: restenosis, atherosclerosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth. Examples of dual $\alpha v\beta 3/\alpha v\beta 5$ antagonizing effects are inhibition of: bone resorption, restenosis, atherosclerosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth.

Additional examples of the invention are methods of inhibiting bone resorption and of treating and/or preventing osteoporosis in a mammal in need thereof, comprising administering to

- 71 -

the mammal a therapeutically effective amount of any of the compounds or any of the pharmaceutical compositions described above.

Further exemplifying the invention is any of the compositions described above, further comprising a therapeutically effective amount of a second bone resorption inhibitor; preferably, the
5 second bone resorption inhibitor is alendronate.

More particularly illustrating the invention is any of the methods of treating and/or preventing osteoporosis and/or of inhibiting bone resorption described above, wherein the compound is administered
10 in combination with a second bone resorption inhibitor; preferably, the second bone resorption inhibitor is alendronate.

Additional illustrations of the invention are methods of treating hypercalcemia of malignancy, osteopenia due to bone metastases, periodontal disease, hyperparathyroidism, periarticular
15 erosions in rheumatoid arthritis, Paget's disease, immobilization-induced osteopenia, and glucocorticoid treatment in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds or any of the pharmaceutical compositions described above.

More particularly exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment and/or prevention of osteoporosis in a mammal in need thereof. Still further exemplifying the invention is the use of any of the compounds described above in the preparation of a
20 medicament for the treatment and/or prevention of: bone resorption, tumor growth, cancer, restenosis, atherosclerosis, diabetic retinopathy, macular degeneration, inflammation and/or angiogenesis.

Additional illustrations of the invention are methods of treating tumor growth in a mammal in need thereof, comprising
30 administering to the mammal a therapeutically effective amount of a compound described above and one or more agents known to be cytotoxic or antiproliferative, e.g., taxol and doxorubicin.

- 72 -

DETAILED DESCRIPTION OF THE INVENTION

Representative compounds of the present invention are integrin antagonists which display submicromolar affinity for the human $\alpha v \beta 3$ receptor. Compounds of this invention are therefore useful for
5 treating mammals suffering from a bone condition caused or mediated by increased bone resorption, who are in need of such therapy. Pharmacologically effective amounts of the compounds, including pharmaceutically acceptable salts thereof, are administered to the mammal, to inhibit the activity of mammalian osteoclasts.

10 The compounds of the present invention are administered in dosages effective to antagonize the $\alpha v \beta 3$ receptor where such treatment is needed, as, for example, in the prevention or treatment of osteoporosis. For use in medicine, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other
15 salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or
20 inorganic acid. Representative salts include the following:

Acetate, Benzenesulfonate, Benzoate, Bicarbonate, Bisulfate, Bitartrate, Borate, Bromide, Calcium, Camsylate, Carbonate, Chloride, Clavulanate, Citrate, Dihydrochloride, Edetate, Edisylate, Estolate, Esylate, Fumarate, Gluceptate, Gluconate, Glutamate,
25 Glycolylarsanilate, Hexylresorcinatate, Hydrabamine, Hydrobromide, Hydrochloride, Hydroxynaphthoate, Iodide, Isothionate, Lactate, Lactobionate, Laurate, Malate, Maleate, Mandelate, Mesylate, Methylbromide, Methylnitrate, Methylsulfate, Mucate, Napsylate, Nitrate, N-methylglucamine ammonium salt, Oleate, Oxalate, Pamoate
30 (Embonate), Palmitate, Pantothenate, Phosphate/diphosphate, Polygalacturonate, Salicylate, Stearate, Sulfate, Subacetate, Succinate, Tannate, Tartrate, Teoclate, Tosylate, Triethiodide and Valerate. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include

- 73 -

alkali metal salts, e.g., sodium or potassium salts; alkaline earth metal salts, e.g., calcium or magnesium salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts.

The compounds of the present invention, may have chiral
5 centers and occur as racemates, racemic mixtures, diastereomeric mixtures, and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention. Therefore, where a compound is chiral, the separate enantiomers, substantially free of the other, are included within the scope of the invention; further
10 included are all mixtures of the two enantiomers. Also included within the scope of the invention are polymorphs and hydrates of the compounds of the instant invention.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be
15 functional derivatives of the compounds of this invention which are readily convertible in vivo into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not
20 be specifically disclosed, but which converts to the specified compound in vivo after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs," ed. H. Bundgaard, Elsevier, 1985. Metabolites of these compounds include active species
25 produced upon introduction of compounds of this invention into the biological milieu.

The term "therapeutically effective amount" shall mean that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being
30 sought by a researcher or clinician.

The term "vitronectin receptor antagonist," as used herein, refers to a compound which binds to and antagonizes either the $\alpha v \beta 3$ receptor or the $\alpha v \beta 5$ receptor, or a compound which binds to and

- 74 -

antagonizes both the $\alpha v\beta 3$ and $\alpha v\beta 5$ receptors (i.e., a dual $\alpha v\beta 3/\alpha v\beta 5$ receptor antagonist).

The term "bone resorption," as used herein, refers to the process by which osteoclasts degrade bone.

5 The term "alkyl" shall mean straight or branched chain alkanes of one to ten total carbon atoms, or any number within this range (i.e., methyl, ethyl, 1-propyl, 2-propyl, n-butyl, s-butyl, t-butyl, etc.).

10 The term "alkenyl" shall mean straight or branched chain alkenes of two to ten total carbon atoms, or any number within this range.

The term "alkynyl" shall mean straight or branched chain alkynes of two to ten total carbon atoms, or any number within this range.

15 The term "cycloalkyl" shall mean cyclic rings of alkanes of three to eight total carbon atoms, or any number within this range (i.e., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl).

20 The term "cycloheteroalkyl," as used herein, shall mean a 3- to 8-membered fully saturated heterocyclic ring containing one or two heteroatoms chosen from N, O or S. Examples of cycloheteroalkyl groups include, but are not limited to piperidinyl, pyrrolidinyl, azetidiny, morpholinyl, piperazinyl.

25 The term "alkoxy," as used herein, refers to straight or branched chain alkoxides of the number of carbon atoms specified (e.g., C1-5 alkoxy), or any number within this range (i.e., methoxy, ethoxy, etc.).

30 The term "aryl," as used herein, refers to a monocyclic or polycyclic system composed of 5- and 6-membered fully unsaturated or partially unsaturated rings, such that the system comprises at least one fully unsaturated ring, wherein the rings contain 0, 1, 2, 3 or 4 heteroatoms chosen from N, O or S, and either unsubstituted or substituted with one or more groups independently selected from hydrogen, halogen, C1-10 alkyl, C3-8 cycloalkyl, aryl, aryl C1-8 alkyl,

- 75 -

- amino, amino C₁₋₈ alkyl, C₁₋₃ acylamino, C₁₋₃ acylamino C₁₋₈ alkyl, C₁₋₆ alkylamino, C₁₋₆ alkylamino C₁₋₈ alkyl, C₁₋₆ dialkylamino, C₁₋₆ dialkylamino-C₁₋₈ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxy C₁₋₆ alkyl, hydroxycarbonyl, hydroxycarbonyl C₁₋₆ alkyl, C₁₋₅ alkoxycarbonyl, C₁₋₃ alkoxycarbonyl C₁₋₆ alkyl, hydroxycarbonyl C₁₋₆ alkyloxy, hydroxy, hydroxy C₁₋₆ alkyl, cyano, trifluoromethyl, oxo or C₁₋₅ alkylcarbonyloxy. Examples of aryl include, but are not limited to, phenyl, naphthyl, pyridyl, pyrazinyl, pyrimidinyl, imidazolyl, benzimidazolyl, indolyl, thienyl, furyl, dihydrobenzofuryl, benzo(1,3)dioxolane, oxazolyl, isoxazolyl and thiazolyl, which are either unsubstituted or substituted with one or more groups independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₈ cycloalkyl, aryl, aryl C₁₋₈ alkyl, amino, amino C₁₋₈ alkyl, C₁₋₃ acylamino, C₁₋₃ acylamino C₁₋₈ alkyl, C₁₋₆ alkylamino, C₁₋₆ alkylamino-C₁₋₈ alkyl, C₁₋₆ dialkylamino, C₁₋₆ dialkylamino C₁₋₈ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxy C₁₋₆ alkyl, hydroxycarbonyl, hydroxycarbonyl C₁₋₆ alkyl, C₁₋₅ alkoxycarbonyl, C₁₋₃ alkoxycarbonyl C₁₋₆ alkyl, hydroxycarbonyl C₁₋₆ alkyloxy, hydroxy, hydroxy C₁₋₆ alkyl, cyano, trifluoromethyl, oxo or C₁₋₅ alkylcarbonyloxy. Preferably, the aryl group is unsubstituted, mono-, di-, tri- or tetra-substituted with one to four of the above-named substituents; more preferably, the aryl group is unsubstituted, mono-, di- or tri-substituted with one to three of the above-named substituents; most preferably, the aryl group is unsubstituted, mono- or di-substituted with one to two of the above-named substituents.

- Whenever the term "alkyl" or "aryl" or either of their prefix roots appear in a name of a substituent (e.g., aryl C₀₋₈ alkyl) it shall be interpreted as including those limitations given above for "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C₁₋₁₀) shall refer independently to the number of carbon atoms in an alkyl or cyclic alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

The terms "arylalkyl" and "alkylaryl" include an alkyl portion where alkyl is as defined above and to include an aryl portion where aryl is as defined above. The C_{0-m} or C_{1-m} designation where

- 76 -

m may be an integer from 1-10 or 2-10 respectively refers to the alkyl component of the arylalkyl or alkylaryl unit. Examples of arylalkyl include, but are not limited to, benzyl, fluorobenzyl, chlorobenzyl, phenylethyl, phenylpropyl, fluorophenylethyl, chlorophenylethyl, thienylmethyl, thienylethyl, and thienylpropyl. Examples of alkylaryl include, but are not limited to, toluene, ethylbenzene, propylbenzene, methylpyridine, ethylpyridine, propylpyridine and butylpyridine.

- 5 When substituent R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} or R^{14} includes the definition C0 (e.g., aryl C0-8 alkyl), the group modified by C0 is not present in the substituent. Similarly, when any of the variables m, q, r or s is zero, then the group modified by the variable is not present; for example, when s is zero, the group $-(CH_2)_s C \equiv CH$ is $-C \equiv CH$. In addition, the substituent $-(C_{1-6} \text{ alkyl})_q \text{amino}$ where q is zero, one or two, refers to an amino, 15 C_{1-6} alkylamino and C_{1-6} dialkylamino group, respectively. When a C_{1-6} dialkylamino substituent is intended, the C_{1-6} alkyl groups can be the same (e.g., dimethylamino) or different (e.g., $N(CH_3)(CH_2CH_3)$). Similarly, the substituent $-(aryl)_q \text{amino}$ or $-(aryl \ C_{1-6} \text{ alkyl})_q \text{amino}$, where q is zero, one or two, refers to an amino, 20 arylamino and diarylamino group, [or an amino, aryl C_{1-6} alkylamino or di-(aryl C_{1-6} alkyl)amino] respectively, where the aryl [or aryl C_{1-6} alkyl] groups in a diarylamino [or di-(aryl C_{1-6} alkyl)amino] substituent can be the same or different.


- 25 The term "halogen" shall include iodine, bromine, chlorine and fluorine.

The term "oxy" means an oxygen (O) atom. The term "thio" means a sulfur (S) atom. The term "oxo" shall mean $=O$.

- The term "substituted" shall be deemed to include multiple degrees of substitution by a named substituent. Where multiple 30 substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally. By independently substituted, it is meant that the (two or more) substituents can be the same or different.

- 77 -

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. For example, a C₁₋₅ alkylcarbonylamino C₁₋₆ alkyl substituent is equivalent to



The present invention is also directed to combinations of the compounds of the present invention with one or more agents useful in the prevention or treatment of osteoporosis. For example, the compounds of the instant invention may be effectively administered in combination with effective amounts of other agents used in the treatment of osteoporosis such as bisphosphonate bone resorption inhibitors; preferably, the bone resorption inhibitor is the bisphosphonate alendronate, now sold as FOSAMAX®. Preferred combinations are simultaneous or alternating treatments of an $\alpha_v\beta_3$ receptor antagonist of the present invention and FOSAMAX®.

In addition, the integrin ($\alpha_v\beta_3$) antagonist compounds of the present invention may be effectively administered in combination with a growth hormone secretagogue in the therapeutic or prophylactic treatment of disorders in calcium or phosphate metabolism and associated diseases. These diseases include conditions which can benefit from a reduction in bone resorption. A reduction in bone resorption should improve the balance between resorption and formation, reduce bone loss or result in bone augmentation. A reduction in bone resorption can alleviate the pain associated with osteolytic lesions and reduce the incidence and/or growth of those lesions. These diseases include: osteoporosis (including estrogen deficiency, immobilization, glucocorticoid induced and senile), osteodystrophy, Paget's disease, myositis ossificans, Bechterew's disease, malignant hypercalcemia, metastatic bone disease, periodontal disease, cholelithiasis, nephrolithiasis, urolithiasis, urinary calculus, hardening of the arteries (sclerosis), arthritis, bursitis, neuritis and tetany. Increased bone

- 78 -

resorption can be accompanied by pathologically high calcium and phosphate concentrations in the plasma, which would be alleviated by this treatment. Similarly, the present invention would be useful in increasing bone mass in patients with growth hormone deficiency.

- 5 Thus, preferred combinations are simultaneous or alternating treatments of an $\alpha v \beta 3$ receptor antagonist of the present invention and a growth hormone secretagogue, optionally including a third component comprising FOSAMAX®.

- 10 In addition, the vitronectin receptor antagonist compounds of the present invention may be effectively administered in combination with one or more agents known to be cytotoxic or antiproliferative, e.g., taxol and doxorubicin.

- 15 In accordance with the method of the present invention, the individual components of the combination can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly. It will be understood that the scope of combinations of the compounds of this invention with other agents useful for treating $\alpha v \beta 3$ related conditions includes in principle any combination with any pharmaceutical composition useful for treating osteoporosis.

- 20 As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

- 25 The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixers, tinctures, suspensions, syrups and emulsions. Likewise, they may also be administered in intravenous (bolus or infusion), intraperitoneal, topical (e.g., ocular eyedrop), subcutaneous,

- 79 -

intramuscular or transdermal (e.g., patch) form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed as an $\alpha v\beta 3$ inhibitor.

5 The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound
10 or salt thereof employed. An ordinarily skilled physician, veterinarian or clinician can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

 Oral dosages of the present invention, when used for the
15 indicated effects, will range between about 0.01 mg per kg of body weight per day (mg/kg/day) to about 100 mg/kg/day, preferably 0.01 to 10 mg/kg/day, and most preferably 0.1 to 5.0 mg/kg/day. For oral administration, the compositions are preferably provided in the form of tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0,
20 50.0, 100 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A medicament typically contains from about 0.01 mg to about 500 mg of the active ingredient, preferably, from about 1 mg to about 100 mg of active ingredient. Intravenously, the most preferred doses will range
25 from about 0.1 to about 10 mg/kg/minute during a constant rate infusion. Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, preferred compounds for the present invention can be
30 administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage

- 80 -

administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as 'carrier' materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the

- 81 -

compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxy-ethylaspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polyactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and crosslinked or amphipathic block copolymers of hydrogels.

In the schemes and examples below, various reagent symbols and abbreviations have the following meanings:

15	AcOH:	Acetic acid.
	BH ₃ •DMS:	Borane•dimethylsulfide.
	BOC(Boc):	t-Butyloxycarbonyl.
	BOP:	Benzotriazol-1-yloxytris(dimethylamino)-
20		phosphonium hexafluorophosphate.
	CBZ(Cbz):	Carbobenzyloxy or benzyloxycarbonyl.
	CDI:	Carbonyldiimidazole.
	CH ₂ Cl ₂ :	Methylene chloride.
	CHCl ₃ :	Chloroform.
25	DEAD:	Diethyl azodicarboxylate.
	DIAD:	Diisopropyl azodicarboxylate.
	DIBAH or	
	DIBAL-H:	Diisobutylaluminum hydride.
	DIPEA:	Diisopropylethylamine.
30	DMAP:	4-Dimethylaminopyridine.
	DME:	1,2-Dimethoxyethane.
	DMF:	Dimethylformamide.
	DMSO:	Dimethylsulfoxide.
	DPFN:	3,5-Dimethyl-1-pyrazolylformamidinium nitrate.

- 82 -

	EDC:	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide.
	EtOAc:	Ethyl acetate.
	EtOH:	Ethanol.
	HOAc:	Acetic acid.
5	HOAT:	1-Hydroxy-7-azabenzotriazole
	HOBT:	1-Hydroxybenzotriazole.
	LDA:	Lithium diisopropylamide.
	MeOH:	Methanol.
	NEt ₃ :	Triethylamine.
10	NMM:	N-methylmorpholine.
	PCA•HCl:	Pyrazole carboxamidine hydrochloride.
	Pd/C:	Palladium on activated carbon catalyst.
	Ph:	Phenyl.
	pTSA	p-Toluene sulfonic acid.
15	TEA:	Triethylamine.
	TFA:	Trifluoroacetic acid.
	THF:	Tetrahydrofuran.
	TLC:	Thin Layer Chromatography.
	TMEDA:	N,N,N',N'-Tetramethylethylenediamine.
20	TMS:	Trimethylsilyl.

The novel compounds of the present invention were prepared according to the procedure of the following schemes and examples, using appropriate materials and are further exemplified by the following specific examples. The most preferred compounds of the invention are any or all of those specifically set forth in these examples. These compounds are not, however, to be construed as forming the only genus that is considered as the invention, and any combination of the compounds or their moieties may itself form a genus. The following examples further illustrate details for the preparation of the compounds of the present invention. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures are degrees Celsius unless otherwise noted.

- 83 -


The following Schemes and Examples describe procedures for making representative compounds of the present invention.

Moreover, by utilizing the procedures described in detail in PCT International Application Publication Nos. WO95/32710, published 7


- 5 December 1995, and WO95/17397, published 29 June 1995, in conjunction with the disclosure contained herein, one of ordinary skill in the art can readily prepare additional compounds of the present invention claimed herein.

- More specifically, procedures for preparing the N-terminus
10 of the compounds of the present invention are described in WO 95/32710. Additionally, for a general review describing the synthesis of β -alanines which can be utilized as the C-terminus of the compounds of the present invention, see Cole, D.C., *Recent Stereoselective Synthetic Approaches to β -Amino Acids*, *Tetrahedron*, 1994, 50, 9517-9582;
15 Juaristi, E, et al., *Enantioselective Synthesis of β -Amino Acids*, *Aldrichemica Acta*, 1994, 27, 3. In particular, synthesis of the 3-methyl β -alanine is taught in Duggan, M.F. et al., *J. Med. Chem.*, 1995, 38, 3332-3341; the 3-ethynyl β -alanine is taught in Zablocki, J.A., et al., *J. Med. Chem.*, 1995, 38, 2378-2394; the 3-pyrid-3-yl β -alanine is taught
20 in Rico, J.G. et al., *J. Org. Chem.*, 1993, 58, 7948-7951; and the 2-amino and 2-tosylamino β -alanines are taught in Xue, C-B, et al., *Biorg. Med. Chem. Letts.*, 1996, 6, 339-344.

- 84 -

SCHEME 1

- 85 -

SCHEME 1 (CONTD)

2-Oxo-3-(3-oxobutyl)piperidine (1-3)

A stirred solution of TMEDA (3.0 g, 20 mmol), 0.5 M LDA (6 mL, in THF), and THF (10 mL) at -78°C was treated with 1-1 (1.7 g, 10 mmol) (for preparation, see: JOC, 1990, 55, 3682) to effect
5 an orange solution.

After 1 h, the iodide 1-2 (2.4 g, 10 mmol) (*J. Org. Chem.*, 1983, 48, 5381) was added to the orange solution and the resulting solution stirred for 2 h at -78°C, 3 h at -15°C and then 16 h at ambient temperature. The reaction mixture was concentrated and then treated
10 with 1N HCl (30 mL). The mixture was then basified with 1N NaOH/brine followed by extraction with EtOAc (3x). The combined extracts were dried (MgSO₄) and concentrated to give a yellow oil. Flash chromatography (silica, EtOAc → 10% CH₃OH/EtOAc) gave 1-3 as a colorless solid.
15 TLC R_f 0.42 (silica, 10% CH₃OH/EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 5.75 (bs, 1H), 3.28 (m, 2H), 2.64 (t, 7Hz, 2H), 2.30-1.50 (m, 7H), 2.16 (s, 3H).

2-Oxo-3-[2-([1,8]-naphthyridin-2-yl)ethyl]piperidine (1-5)

A solution of 1-3 (0.25 g, 1.5 mmol), L-proline (85 mg, 0.75 mmol), 1-4 (0.18 g, 1.5 mmol) (for preparation see: *Synth. Commun.* 1987, 17, 1695), and ethanol (10 mL) was refluxed for 24 hr. The cooled solution was concentrated and the residue purified by flash chromatography (silica, EtOAc → 20% CH₃OH/EtOAc) to give 1-5 as a
25 solid.
TLC R_f = 0.32 (silica, 20% CH₃OH/EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 9.08 (m, 1H), 8.16 (m, 1H), 8.10 (d, J=8Hz, 1H), 7.50 (d, J=8Hz, 1H), 7.45 (m, 1H), 5.64 (bs, 1H), 3.31 (m, 2H), 3.18 (m, 2H), 2.50-1.60 (m, 7H).
30

Ethyl 2-Oxo-3-[2-([1,8]-naphthyridin-2-yl)ethyl]piperidin-1-yl-acetate (1-6)

A solution of 1-5 (0.28 g, 1.1 mmol) and DMF (10 mL) at -15°C was treated with NaN(TMS)₂ (1.2 mL, 1.2 mmol, 1M in hexanes)

- 87 -

- to give a red solution. After 30 min, the red solution was treated with ethyl bromoacetate (128 μ L, 1.2 mmol), followed by continued stirring for 1 h. The reaction mixture was then quenched with sat. NH_4Cl and then extracted with EtOAc (3x). The combined extracts were washed with brine, dried (MgSO_4), and concentrated. Flash chromatography (silica, 10% $\text{CH}_3\text{OH}/\text{EtOAc}$) gave 1-6 as a yellow gum.
- 5 TLC R_f = 0.50 (silica, 10% $\text{CH}_3\text{OH}/\text{EtOAc}$)
 ^1H NMR (300 MHz, CDCl_3) δ 9.07 (m, 1H), 8.16 (m, 1H), 8.10 (d, $J=8\text{Hz}$, 1H), 7.50 (d, $J=8\text{Hz}$, 1H), 7.44 (m, 1H), 4.30-3.90 (m, 4H),
10 3.50-3.30 (m, 2H), 3.17 (m, 2H), 2.46 (m, 2H), 2.20-1.70 (m, 5H), 1.28 (t, $J=7\text{Hz}$, 3H).

Ethyl 2-Oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]-naphthyridin-2-yl)ethyl]piperidin-1-yl-acetate (1-7)

- 15 A mixture of 1-6 (102 mg, 0.3 mmol), 10% Pd/C (50 mg), and EtOAc (25 mL) was stirred under a hydrogen atmosphere (1 atm) for 24 h. The catalyst was then removed by filtration through celite and the filtrate concentrated. Flash chromatography (silica, 20% $\text{CH}_3\text{OH}/\text{EtOAc}$) gave 1-7 as a yellow gum.
- 20 TLC R_f = 0.45 (silica, 30% $\text{CH}_3\text{OH}/\text{EtOAc}$)
 ^1H NMR (300 MHz, CDCl_3) δ 7.05 (d, $J=6\text{Hz}$, 1H), 6.41 (d, $J=6\text{Hz}$, 1H), 4.80 (bs, 1H), 4.18 (q, $J=7\text{Hz}$, 2H), 4.08 (m, 2H), 3.37 (m, 4H), 2.80-1.60 (m, 13H), 1.26 (t, 7Hz, 3H).

- 25 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperidin-1-yl-acetic acid (1-8)

- A solution of 1-7 (71 mg, 0.21 mmol) and 6N HCl (15 mL) was stirred at 55°C for 2h, followed by concentration to give 1-8 as a pale yellow gum.
- 30 TLC R_f = 0.09 (silica, 20% $\text{CH}_3\text{OH}/\text{EtOAc}$)

- 88 -

2-Oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]-naphthyridin-2-yl)ethyl]piperidin-1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester (1-10)

A stirred mixture of 1-8 (71 mg, 0.20 mmol), 1-9 (59 mg, 0.22 mmol) (Rico *et al.*, *J. Org. Chem.*, 1993, 58, 7948), NMM (88 μL, 0.8 mmol), and CH₃CN (25 mL) was treated with BOP (97 mg, 0.22 mmol). After 24h, the reaction mixture was concentrated to dryness, dissolved in EtOAc, and then washed with H₂O, dried (MgSO₄), and concentrated. Flash chromatography (silica, 10% (NH₃/EtOH/EtOAc) gave 1-10 as a colorless gum.

TLC R_f = 0.9 (silica, 10% (NH₃/EtOH)/EtOAc)

¹H NMR (300 MHz, CD₃OD) δ 8.55 (m, 1H), 8.43 (m, 1H), 7.83 (m, 1H), 7.40 (m, 1H), 7.11 (m, 1H), 6.37 (m, 1H), 5.38 (m, 1H), 4.08 (q, J=7Hz, 2H), 4.00 (m, 2H), 3.37 (m, 4H), 2.90 (m, 1H), 2.70-1.60 (m, 14H), 1.14 (t, J=7Hz, 3H).


2-Oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]-naphthyridin-2-yl)ethyl]piperidin-1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine trifluoroacetate (1-11)

A stirred solution of 1-11 (52 mg, 0.10 mmol) and 6N HCl (10 mL) was heated at 55°C for 2 h, followed by concentration.


Preparative HPLC (VYDAC C₁₈ semiprep column, gradient elution: [95:5 (0.1% TFA/H₂O/0.1% TFA/CH₃CN) to 50:50 (0.1% TFA/H₂O/0.1% TFA/CH₃CN) 80 min] gave 1-11 as a colorless solid.


¹H NMR (300 MHz, CD₃OD) δ 8.90 (s, 1H), 8.74 (d, J=5Hz, 1H), 8.61 (d, J=8Hz, 1H), 8.03 (m, 1H), 7.56 (d, J=7Hz, 1H), 6.59 (d, J=7Hz, 1H), 5.43 (m, 1H), 4.03 (m, 2H), 3.40 (m, 5H), 3.00 (m, 2H), 2.78 (m, 4H), 2.40-1.60 (m, 12H).

- 89 -

SCHEME 2

- 90 -

SCHEME 2 (CONTD)



(2-Oxo-3-(3-(ethylenedioxy)butyl)pyrrolidin-1-yl)benzyl (2-2)

- To a stirred solution of 2-1 (5.3 g, 30 mmol) and THF (100 mL) at -78°C was added LDA (17.5 mL, 35 mmol, 2.0 M in hexanes) dropwise over a 10 minute period. After 30 min, 1-2 (5.0 g, 21 mmol) was added followed by removal of the cooling bath. After 1 h, the reaction was quenched with AcOH (10 mL) and then diluted with EtOAc, washed with sat. NaHCO₃ and brine, dried (MgSO₄) and concentrated. Flash chromatography (silica, 25% → 75% EtOAc/hexanes) gave 2-2 as an oil.
- 10 TLC R_f = 0.38 (silica, EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 7.25 (m, 5H), 4.48 (d, J=15Hz, 1H), 4.40 (d, J=15Hz, 1H), 3.94 (s, 4H), 3.18 (m, 2H), 2.44 (m, 1H), 2.30-1.30 (m, 9H).

15 2-Oxo-3-(3-(ethylenedioxy)butyl)pyrrolidine (2-3)

- To a stirred solution of 2-2 (4.1 g, 14.2 mmol) in THF (100 mL) at -78°C was added a solution of Li 4,4'-di-tert-butylbiphenyl (188 mL, 0.5 M in THF) in 4 portions. After 1 h, the reaction was quenched with AcOH (25 mL). The resulting mixture was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, and brine, dried (MgSO₄) and concentrated. Flash chromatography (silica, EtOAc → 10% CH₃OH/EtOAc) gave 2-3 as a yellow oil.
- 20 TLC R_f = 0.1 (silica, EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 6.23 (bs, 1H), 3.94 (s, 4H), 3.30 (m, 2H), 2.70 (m, 2H), 2.10-1.30 (m, 9H).
- 25

Ethyl (2-Oxo-3-(3-(ethylenedioxy)butyl)pyrrolidin-1-yl)acetate (2-4)

- To a rapidly stirred solution of 2-3 (0.86 g, 4.3 mmol) and THF (25 mL) at -78°C was added NaN(TMS)₂ (5.2 mL, 5.2 mmol, 1.0 M in THF). After 20 min, ethyl bromoacetate (0.58 mL, 5.2 mmol) was added followed by removal of the cooling bath. After 1 h, the reaction mixture was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃ and brine, dried (MgSO₄), and concentrated to give 2-4 as a yellow oil.
- 30

- 93 -

TLC R_f = 0.53 (silica, EtOAc)

¹H NMR (300 MHz, CDCl₃) δ 4.18 (q, J=7Hz, 2H), 4.04 (m, 2H), 3.93 (s, 4H), 3.39 (m, 2H), 2.44 (m, 1H), 2.23 (m, 1H), 2.00-1.30 (m, 9H), 1.25 (t, J=7H, 3H).

5

Ethyl (2-Oxo-3-(3-oxobutyl)pyrrolidin-1-yl)acetate (2-5)

A solution of 2-4 (1.1 g, 3.9 mmol), p-TSA (5 mg) and acetone (50 mL) was heated at reflux for 1 hr. The cooled reaction mixture was diluted with EtOAc and then washed with sat. NaHCO₃ and

10

brine, dried (MgSO₄), and concentration to afford 2-5 as a yellow oil.

TLC R_f = 0.48 (silica, EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 4.18 (q, J=7Hz, 2H), 4.01 (s, 2H), 3.40 (m, 2H), 2.67 (t, J=7Hz, 2H), 2.48 (m, 1H), 2.30-1.60 (m, 4H), 2.15 (s, 3H), 1.25 (t, J=7Hz, 3H).

15

Ethyl (2-Oxo-3-(2-([1,8]naphthyridin-2-yl)ethyl)pyrrolidin-1-yl)-acetate (2-6)

A mixture of 2-5 (0.77 g, 3.0 mmol), 1-4 (0.55 g, 4.5 mmol, for preparation see *Het*, 1993, 36, 2513), L-proline (0.17 g, 1.5 mmol) and ethanol (25 mL) was heated at reflux for 20 hr. The cooled reaction mixture was concentrated and the residue purified by flash chromatography (silica, EtOAc → 5% CH₃OH/EtOAc) to give 2-6 as a yellow oil.

20

TLC R_f = 0.13 (silica, 10% CH₃OH/EtOAc)

25

¹H NMR (300 MHz, CDCl₃) δ 9.08 (m, 1H), 8.17 (m, 1H), 8.12 (d, J=8Hz, 1H), 7.49 (d, J=8Hz, 1H), 7.46 (m, 1H), 4.15 (q, J=7Hz, 2H), 4.04 (m, 2H), 3.42 (m, 2H), 3.21 (t, J=8Hz, 2H), 2.60-1.80 (m, 5H), 1.25 (t, J=7Hz, 3H).

30

Ethyl (2-Oxo-3-(2-(5,6,7,8-tetrahydro-[1,8]-naphthyridin-2-yl)ethyl)pyrrolidin-1-yl)acetate (2-7)

A mixture of 2-6 (0.87 g, 2.6 mmol), 10% Pd/C (0.5 g), and CH₃OH (25 mL) was stirred under a hydrogen atmosphere (1 atm) for 2 hr. The catalyst was then removed by filtration through a celite

- 94 -

pad followed by concentration of the filtrate. Flash chromatography (silica, EtOAc → 5% CH₃OH/EtOAc) gave 2-7 as a yellow oil.

TLC R_f = 0.18 (silica, 5% CH₃OH/EtOAc)

¹H NMR (300 MHz, CDCl₃) δ 7.05 (d, J=7Hz, 1H), 6.40 (d, J=7Hz, 1H), 4.83 (bs, 1H), 4.17 (q, J=7Hz, 2H), 4.03 (m, 2H), 3.40 (m, 4H), 2.80-1.60 (m, 11H), 1.27 (t, J=7Hz, 3H).

(2-Oxo-3-(2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl)pyrrolidin-1-yl)acetic acid hydrochloride (2-8)

10 A stirred mixture of 2-7 (0.45 g, 1.4 mmol) and 6N HCl (10 mL) was heated at 50°C for 1 h, followed by concentration to give 2-8 as a yellow oil.

¹H NMR (300 MHz, CD₃OD) δ 7.60 (d, J=7Hz, 1H), 6.66 (d, J=7Hz, 1H), 4.05 (s, 2H), 3.50 (m, 4H), 2.83 (m, 4H), 2.54 (m, 1H), 2.32 (m, 15 1H), 2.10 (m, 1H), 2.00-1.75 (m, 4H).

(2-Oxo-3-(2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl)pyrrolidin-1-yl)acetyl-3(S)-ethynyl-β-alanine ethyl ester (2-11)

To a stirred solution of 2-8 (50 mg, 0.15 mmol), 2-9 (29 mg, 0.17 mmol) (Zablocki *et al.*, *J. Med. Chem.*, 1995, 38, 2378), NMM (83 μL, 0.75 mmol), and CH₃CN (1 mL) was added BOP (74 mg, 0.17 mmol). After 20 h, the reaction mixture was diluted with EtOAc and then washed with sat. NaHCO₃, H₂O and brine, dried (MgSO₄), and concentrated to give 2-11 as a yellow oil.

25 TLC R_f = 0.24 (silica, 10% CH₃OH/EtOAc).

(2-Oxo-3-(2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl)pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester (2-12)

To a stirred solution of 2-8 (50 mg, 0.15 mmol), 2-10 (44 mg, 0.17 mmol) (Rico *et al.*, *J. Org. Chem.*, 1993, 58, 7948), NMM (83 μL, 0.75 mmol), and CH₃CN (1 mL) was added BOP (74 mg, 0.17 mmol). After 20 h, the reaction mixture was diluted with EtOAc and then washed with sat. NaHCO₃, H₂O and brine, dried (MgSO₄), and concentrated to give 2-12 as a brown oil.

- 95 -

TLC R_f = 0.24 (silica, 20% CH₃OH/EtOAc).


(2-Oxo-3-(2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl)pyrrolidin-1-yl)acetyl-3(S)-ethynyl-β-alanine (2-13)

- 5 A mixture of 2-11 (0.1 g, 0.15 mmol), 1N NaOH (300 μL, and ethanol (1 mL) was stirred at ambient temperature for 1 hr. Concentration and then flash chromatography (silica, 25:10:1:1 → 15:10:1:1 EtOAc/EtOH/NH₄OH/H₂O) gave 2-13 as a white solid. TLC R_f = 0.18 (silica, 10:10:1:1 EtOAc/EtOH/NH₄OH/H₂O)
- 10 ¹H NMR (300 MHz, CD₃OD) δ 7.45 (m, 1H), 6.50 (m, 1H), 4.53 (m, 1H), 3.80-3.30 (m, 5H), 3.05 (m, 1H), 2.80-2.15 (m, 9H), 2.00-1.75 (m, 4H).

(2-Oxo-3-(2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl)pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine (2-14)


- 15 A mixture of 2-12 (0.1 g, 0.15 mmol), 1N NaOH (300 μL) and ethanol (1 mL) was stirred at ambient temperature for 1 hr. Concentration and the flash chromatography (silica, 25:10:1:1 → 15:10:1:1 EtOAc/EtOH/NH₄OH/H₂O) gave 2-14 as a white solid.
- 20 TLC R_f = 0.10 (silica, 10:10:1:1 EtOAc/EtOH/NH₄OH/H₂O)
- ¹H NMR (300 MHz, CD₃OD) δ 8.57 (m, 1H), 8.40 (m, 1H), 7.86 (m, 1H), 7.40 (m, 2H), 6.50 (m, 1H), 5.28 (m, 1H), 4.65-4.40 (m, 1H), 3.90-1.80 (M, 19H).

- 96 -


SCHEME 3

- 97 -

SCHEME 3 (cont'd)



- 98 -

SCHEME 3 (cont'd)4-(Propyl-2-ene)butyric acid (3-2)

To a stirred suspension of methyltriphenylphosphonium bromide (67.7 g, 190 mmol) in 1 L THF at 0° C was added a solution of sodium bis(trimethylsilyl)amide (190 mL, 190 mmol, 1M THF). After an additional 30 minutes, 3-1 ethyl 4-acetylbutyrate (Aldrich Chemical Co.) (25.0 g, 158 mmol) was added, and the mixture stirred for 18 h. The mixture was filtered, and the filtrate concentrated. The residue was triturated with hexanes, and then filtered. Following evaporative

- 99 -

removal of the solvent, the residue was chromatographed on silica gel, eluting with 10% ethyl acetate/hexanes to give the olefin as a colorless oil. TLC R_f = 0.52 (10% ethyl acetate/hexanes).

¹H NMR (300 MHz, CHCl₃) δ 4.71 (d, 2H, J=13 Hz), 4.13 (q, 2H, J=7 Hz), 2.29 (t, 2H, J=7 Hz), 2.05 (t, 2H, J= 8 Hz), 1.77 (m, 2H), 1.72 (s, 3H), 1.26 (t, 3H, J=7Hz).

A solution of the above olefin (15.4 g, 98.6 mmol), 1 N NaOH (150 mL), and EtOH (300 mL) was stirred at ambient temperature for 2 h. Following acidification with 1 N HCl, the mixture was extracted with ether. The ether layer was washed with brine, dried over magnesium sulfate, and concentrated to give 3-2 as a colorless oil. ¹H NMR (300 MHz, CHCl₃) δ 4.70 (d, 2H, J=13 Hz), 2.27 (t, 2H, J=7 Hz), 2.06 (t, 2H, J= 7 Hz), 1.72 (m, 5H).

15 (4-(Propyl-2-ene)butanoyl)-4(R)-benzyl-2-oxazolidinone (3-3)

To a solution of 3-2 (6.0 g , 46.8 mmol) in THF (200 ml) at -78° C was added triethylamine (7.19 mL , 51.5 mmol) followed by pivaloyl chloride (6.35 mL , 51.5 mmol). The mixture was warmed to 0° C for 1 h, then recooled to -78° C. In a separate flask, of (R)-(+)-4-benzyl-2-oxazolidinone (9.15 g, 51.5 mmol) was dissolved in THF (100 mL), cooled to -78° C, and n-BuLi (32.3 mL, 51.5 mmol; 1.6 M hexanes) was added dropwise. After 10 minutes, the lithium oxazolidinone was added to the pivalic anhydride. After 10 minutes, the mixture was warmed to 0° C for 1.5 h. The mixture was then poured into ethyl acetate, washed with aqueous sodium bicarbonate, and dried over magnesium sulfate. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, dichloromethane) to give 3-3 as a slightly yellow oil.

TLC R_f = 0.8 (CH₂Cl₂).
30 ¹H NMR (300 MHz, CHCl₃) δ 7.40-7.18 (m, 5H), 4.80-4.60 (m, 3H), 4.18 (m, 2H), 3.30 (dd, 1H, J=3.2, 13.2 Hz), 2.95 (m, 2H), 2.76 (dd, 1H, J=9.5, 13.1 Hz), 2.11 (t, 2H, J=7.5 Hz), 1.87 (m, 2H), 1.74 (s, 3H).

- 100 -

2-Chloroethyltriflate (3-4)

To a solution of 1.67 mL (24.8 mmol) of 2-chloroethanol and 3.47 mL (29.8 mmol) of 2,6-lutidine in 20 mL of dichloromethane at 0° C was added 4.59 mL (27.3 mmol) of triflic anhydride. After 1 h, the mixture was diluted with hexanes, washed with ice-cold 1N HCl, and dried over sodium sulfate. The solvents were evaporated to give 3-4 as a pink oil.

¹H NMR (300 MHz, CHCl₃) δ 4.69 (t, 2H, J=5.3 Hz), 3.78 (t, 2H, J=5.6 Hz).

2(S)-Chloroethyl-4-(propyl-2-ene)butanoyl-(4(R)-benzyl-2-oxazolidinone) (3-5)

To a solution of 3-3 (11.0 g, 38.3 mmol) in THF (60 mL) at -78° C was added a solution of sodium bis(trimethylsilyl)amide (42.1 mL, 42.1 mmol; 1M/THF). After 20 min, 3-4 (16.2 mL, 115 mmol) was added over 5 min, and the resulting mixture stirred for 1.5 h at -78° C, then 2 h at -15° C. The mixture was diluted with hexanes, washed with sat. ammonium chloride, and dried over sodium sulfate. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 14% ethyl acetate/hexanes) to give 3-5 as a colorless oil. TLC R_f = 0.5 (20% ethyl acetate/hexanes).

¹H NMR (300 MHz, CHCl₃) δ 7.30-7.18 (m, 5H), 4.67 (m, 3H), 4.19 (m, 2H), 3.99 (m, 1H), 3.58 (m, 2H), 3.33 (dd, 1H, J=3.2, 12.0 Hz), 2.75 (dd, 1H, J=9.7, 13.5 Hz), 2.23 (m, 1H), 2.18-1.82 (m, 4H), 1.77-1.60 (m, 1H), 1.71 (s, 3H).

Ethyl 2-oxo-3(S)-(3-methylenebutyl)pyrrolidine (3-6)

A mixture of 3-5 (8.15 g, 23.3 mmol) and NaN₃ (4.54 g, 69.8 mmol) in DMSO (120 mL) was heated at 75° C for 2 h. After cooling, the mixture was diluted with ether and hexanes, washed with water, and dried over sodium sulfate. Evaporative removal of the solvent gave the azide as a colorless oil. TLC R_f = 0.5 (20% ethyl acetate/hexanes).

- 101 -

¹H NMR (300 MHz, CHCl₃) δ 7.30-7.22 (m, 5H), 4.69 (m, 3H), 4.17 (d, 2H, J=5.1 Hz), 3.89 (m, 1H), 3.38 (m, 3H), 2.74 (m, 1H), 2.13-1.63 (m, 6H), 1.71 (s, 3H).

- To a solution of of this azide (8.0 g , 22.4 mmol) in THF (250 mL) and water (40 mL) was added triphenylphosphine (8.24 g , 31.4 mmol) in 4 portions over 5 minutes. This mixture was heated at reflux for 2 h, cooled, and evaporated. The residue was chromatographed (silica gel, 10% chloroform/ethyl acetate) to give 3-6 as a colorless oil.
- 5
- 10 TLC R_f = 0.40 (20% chloroform/ethyl acetate).
¹H NMR (300 MHz, CHCl₃) δ 6.47 (br s, 1H), 4.73 (m, 2H), 3.31 (m, 2H), 2.33 (m, 2H), 2.08 (m, 3H), 1.81 (m, 1H), 1.74 (s, 3H), 1.44 (s, 1H).

15 Ethyl 2-oxo-3(S)-(3-methylenebutyl)pyrrolidin-1-yl)acetate (3-7)

- To a solution of 3-6 (2.50 g, 16.3 mmol) in THF (40 mL) at -78° C was added sodium bis(trimethylsilyl)amide (17.1 mL, 17.1 mmol; 1M/ THF) dropwise. After an additional 20 min, ethyl bromoacetate (2.17 mL, 19.6 mmol) was added dropwise over 3 min.
- 20 After an additional 20 min, 20 mL sat. aqueous NH₄Cl was added, and the cooling bath removed. The layers were separated, the aqueous layer washed with ether, and the combined organic extracts were dried over sodium sulfate. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 40% ethyl acetate/hexanes) to
- 25 give 3-7 as a colorless oil.
TLC R_f = 0.85 (50% chloroform/ethyl acetate).
¹H NMR (300 MHz, CHCl₃) δ 4.73 (m, 2H), 4.18 (q, 2H, J=7.1Hz), 4.06 (dd, 2H, J=17.6, 20.8 Hz), 3.42 (m, 2H), 2.44 (m, 1H), 2.27 (m, 1H), 2.12 (m, 3H), 1.75 (m, 1H), 1.74 (s, 3H), 1.50 (m, 1H), 1.28 (t, 3H, J=7.3 Hz).
- 30

Ethyl 2-oxo-3(S)-(3-oxo-butyl)pyrrolidin-1-yl)acetate (3-8)

To a solution of 3-7 (3.35 g, 14.0 mmol) and N-methylmorpholine-N-oxide (3.27 g, 28.0 mmol) in THF (10 mL) and

- 102 -

water (1 mL) was added OsO₄ (5.7 mL, 0.56 mmol; 2.5 % t-butanol). After 1 h, NaIO₄ (5.99 g, 28 mmol) in warm water (30 mL) was added over 2 min, and the resulting mixture stirred for 1 h. Water was then added, and the aqueous layer washed with ether and ethyl acetate, and
5 the combined organic extracts were dried over sodium sulfate. Evaporative removal of the solvent gave 3-8 as a dark oil containing residual OsO₄.

TLC R_f = 0.78 (70:20:10 chloroform/ethyl acetate/MeOH).

¹H NMR (300 MHz, CHCl₃) δ 4.19 (m, 2H, J=7.2 Hz), 4.03 (s, 2H),
10 3.41 (m, 2H), 2.68 (t, 2H, J=9.4 Hz) 2.45 (m, 1H), 2.27 (m, 1H), 2.17 (s, 3H), 1.97 (m, 1H), 1.78 (m, 2H), 1.28 (t, 3H, J=7.2 Hz).

Ethyl 2-oxo-3(S)-[2-([1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetate (3-9)

15 A mixture of 3-8 (3.25 g, 13.5 mmol), 1-4, 2-amino-3-formylpyridine (2.2 g, 18.2 mmol; for preparation see *Synth. Commun.* 1987, 17, 1695) and proline (0.62 g, 5.39 mmol) in absolute ethanol (45 mL) was heated at reflux for 15 h. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:25:5
20 chloroform/ethyl acetate/MeOH to give 3-9 as a colorless oil. TLC R_f = 0.24 (70:25:5 chloroform/ethyl acetate/MeOH).
¹H NMR (300 MHz, CHCl₃) δ 9.08 (m, 1H), 8.16 (m, 2H), 7.47 (m, 2H), 4.17 (m, 4H), 3.42 (m, 2H), 3.21 (t, 2H, J=6.0 Hz), 2.56 (m, 1H), 2.39 (m, 2H), 2.08 (m, 1H), 1.87 (m, 1H), 1.27 (t, 3H, J=7.1 Hz).

25 Ethyl 2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetate (3-10)

A mixture of 3-9 (3.33 g, 10.2 mmol) and 10% Pd/carbon (1.5 g) in EtOH (50 mL) was stirred under a balloon of hydrogen for
30 13 h. Following filtration and evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:20:10 chloroform/ethyl acetate/MeOH to give 3-10 as a colorless oil. TLC R_f = 0.20 (70:20:10 chloroform/ethyl acetate/MeOH).

- 103 -

¹H NMR (300 MHz, CHCl₃) δ 7.05 (d, 1H, J=7.3 Hz), 6.38 (d, 1H, J=7.3 Hz), 4.88 (br s, 1H), 4.17 (dd, 2H, J=7.0, 14.4 Hz), 4.04 (dd, 2H, J=17.6, 27.3 Hz), 3.40 (m, 4H), 2.69 (m, 4H), 2.51 (m, 1H), 2.28 (m, 2H), 1.90 (m, 2H), 1.78 (m, 2H), 1.27 (t, 3H, J=6.9 Hz).

5

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetic acid (3-11)

A mixture of 3-10 (0.60 g, 1.81 mmol) and 6N HCl (25 mL) was heated at 60° C for 1 h. Evaporative removal of the solvent gave 3-11 as a yellow oil.

10

¹H NMR (300 MHz, DMSO-d₆) δ 8.4 (br s, 1H), 7.60 (d, 1H, J=7.3 Hz), 6.63 (d, 1H, J=7.3 Hz), 3.92 (dd, 2H, J=17.6, 25.9 Hz), 3.43 (m, 2H), 3.35 (m, 2H), 2.74 (m, 4H), 2.28 (m, 2H), 2.03 (m, 1H), 1.82 (m, 2H), 1.67 (m, 2H).

15

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester (3-12)

A mixture of 3-11 (0.20 g, 0.588 mmol), 2-9 (0.157 g, 0.882 mmol), EDC (0.147 g, 0.765 mmol), HOBT (0.095 g, 0.706 mmol) and NMM (0.453 mL, 4.12 mmol) in CH₃CN (3 mL) and DMF (2 mL) was stirred for 20 h. The mixture was diluted with ethyl acetate, washed with water, brine, and dried over sodium sulfate. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:20:10 chloroform/ethyl acetate/MeOH) to give 3-12 as a colorless foam.

25

TLC R_f = 0.44 (70:20:10 chloroform/ethyl acetate/MeOH).

¹H NMR (300 MHz, CHCl₃) δ 7.06 (d, 1H, J=7.3 Hz), 6.39 (d, 1H, J=7.3 Hz), 5.07 (m, 1H), 4.94 (br s, 1H), 4.18 (q, 2H, J=6.1 Hz), 3.95 (q, 2H, J=16.1 Hz), 3.39 (m, 4H), 2.90 (s, 1H), 2.68 (m, 6H), 2.50 (m, 1H), 2.27 (m, 3H), 1.82 (m, 4H), 1.27 (t, 3H, J=7.1 Hz).

30

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine (3-13)

- 104 -

To a solution of 3-12 (0.050 g, 0.117 mmol) in EtOH (1 mL) was added 1N NaOH (0.164 ml, 0.164 mmol). After stirring for 2 h, the solvents were evaporated and the residue was chromatographed (silica gel, 25:10:1:1 ethyl acetate/EtOH/water/NH₄OH to give 3-13 as a colorless foam.

TLC R_f = 0.26 (25:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

¹H NMR (300 MHz, DMSO-d₆) δ 7.75 (br s, 1H), 7.14 (d, 1H, J=7.3 Hz), 6.31 (d, 1H, J=7.3 Hz), 4.74 (m, 1H), 3.90 (d, 1H, J=16.6 Hz), 3.67 (d, 1H, J=16.6 Hz), 3.23 (m, 4H), 2.57 (m, 7H), 2.30 (m, 1H), 2.11 (m, 2H), 1.73 (m, 2H), 1.59 (m, 2H).

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester (3-14)

A mixture of 3-11 (0.30 g, 0.882 mmol), 2-10 (0.354 g, 1.32 mmol), EDC (0.220 g (1.15 mmol), HOBT (0.143 g, 1.05 mmol) and NMM (0.680 mL (6.18 mmol) in CH₃CN (5 mL) and DMF (3 mL) at 0° C was stirred for 10 min, then allowed to warm and stir for 20 h. The mixture was diluted with ethyl acetate, washed with water, brine, and dried over sodium sulfate. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:20:10 chloroform/ethyl acetate/MeOH) to give 3-14 as a colorless foam.

TLC R_f = 0.31 (70:20:10 chloroform/ethyl acetate/MeOH).

¹H NMR (300 MHz, CHCl₃) δ 8.55 (d, 1H, J=2.2 Hz), 8.50 (dd, 1H, J=1.5, 4.6 Hz), 7.64 (m, 2H), 7.23 (m, 1H), 7.05 (d, 1H, J=7.3 Hz), 6.38 (d, 1H, J=7.3 Hz), 5.40 (m, 1H), 4.98 (br s, 1H), 4.01 (m, 4H), 3.39 (m, 4H), 2.85 (m, 2H), 2.68 (m, 4H), 2.49 (m, 1H), 2.25 (m, 2H), 1.83 (m, 4H), 1.16 (t, 3H, J=7.2 Hz).

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine (3-15)

To a solution of 3-14 (0.049 g, 0.102 mmol) in THF (1 mL) and water (0.3 mL) at 0° C was added 1M LiOH (0.112 ml, 0.112 mmol). After warming to ambient temperature and stirring for 2 h, the solvents were evaporated and the residue was chromatographed (silica


- 105 -

gel, 25:10:1:1 ethyl acetate/EtOH/water/NH₄OH to give 3-15 as a colorless foam.


TLC R_f = 0.15 (25:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

¹H NMR (300 MHz, DMSO-d₆) δ 8.74 (d, 1H, J=8.3 Hz), 8.51 (m, 1H), 8.42 (m, 2H), 7.70 (d, 1H, J=8.1 Hz), 7.33 (m, 1H), 7.21 (d, 1H, J=7.3 Hz), 6.36 (d, 1H, J=7.3 Hz), 5.14 (m, 1H), 4.00 (d, 1H, J=16.8 Hz), 3.70 (d, 1H, J=16.6 Hz), 3.30 (m, 4H), 2.68 (m, 7H), 2.20 (m, 3H), 1.71 (m, 4H).

SCHEME 4



- 106 -

SCHEME 4 (cont'd)

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-ylacetyl-3(S)-alkynyl-β-alanine ethyl ester (4-2)

- Prepared from 4-1 (prepared by the method used to
 5 prepare 3-11, utilizing (S)-(-)-4-benzyl-2-oxazolidinone) and 2-9, by the
 method used to prepare 3-12.

- 107 -

¹H NMR (300 MHz, CHCl₃) δ 7.06 (d, 1H, J=7 Hz), 6.39 (d, 1H, J=7 Hz), 5.06 (m, 1H), 4.84 (br s, 1H), 4.16 (q, 2H, J=6 Hz), 3.93 (m, 2H), 3.38 (m, 4H), 2.68 (m, 6H), 2.52 (m, 1H), 2.25 (m, 2H), 1.90 (m, 2H), 1.78 (m, 2H), 1.26 (t, 3H, J=7Hz).

5

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethylpyrrolidin-1-yl]acetyl-3(S)-alkynyl-β-alanine (4-3)

Prepared from 4-2 (0.05 g, 0.11 mmol) by the method used to prepare 3-13.

10 ¹H NMR (300 MHz, CD₃OD, 1 drop 1N NaOD) δ 7.11 (d, 1H, J=7 Hz), 6.40 (d, 1H, J=7 Hz), 4.90 (m, 1H), 3.94 (q, 2H, J= 17 Hz), 3.39 (m, 4H), 2.69 (d, 2H, J= 6 Hz), 2.60 (m, 2H), 2.52 (d, J=7 Hz), 2.49 (m, 1H), 2.27 (m, 1H), 2.13 (m, 1H), 1.85 (m, 4H), 1.68 (m, 1H).

15 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl]acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester (4-4)

Prepared from 4-1 (0.35 g, 1.0 mmol) and 2-10 (0.33 g, 1.2 mmol) by the method used to prepare 3-14.


20 ¹H NMR (300 MHz, CHCl₃) δ 8.55 (d, 1H, J= 2 Hz), 8.55 (dd, 1H, J= 2 , 5 Hz), 7.61 (m, 1H), 7.54 (m, 1H), 7.06 (d, 1H), 6.38 (d, 1H, J=7 Hz), 5.40 (m, 1H), 4.90 (br s, 1H), 4.05 (q, 2H, J=7 Hz), 3.95 (m, 2H), 3.42 (m, 4H), 2.85 (dd, 2H, J=2, 6 Hz), 2.67 (m, 4H), 2.53 (m, 1H), 2.27 (m, 2H), 1.90 (m, 2H), 1.78 (m, 2H), 1.16 (m, 3H, J=7 Hz).

25 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro-[1,8]-naphthyridin-2-yl)ethylpyrrolidin-1-yl]acetyl-3(S)-pyridin-3-yl-β-alanine (4-5)


Prepared from 4-4 (0.16 g, 0.33 mmol) by the method used to prepare 3-15.

30 ¹H NMR (300 MHz, CD₃OD) δ 8.57 (s, 1H), 8.42 (m, 1H), 7.86 (d, 1H, J=6 Hz), 7.43 (m, 2H), 6.51 (d, 1H, J=7 Hz), 5.28 (m, 1H), 4.63 (d, 1H, J=17 Hz), 3.60 (m, 2H), 3.47 (d, 1H, J= 17 Hz), 3.35 (m, 3H), 3.14 (td, 1H, J=5, 13 Hz), 2.75 (m, 5H), 2.42 (m, 1H), 2.23 (m, 1H), 1.90 (m, 4H).

- 108 -

SCHEME 5

- 109 -

SCHEME 5 (cont'd)

- 110 -

1,3-Di-tert-butyloxycarbonyl-tetrahydropyrimidine (5-2)

- A heterogeneous mixture of 5-1 (10.0 g, 100 mmol), BOC₂O (48 g, 220 mmol), DMAP (20 mg), and CH₃CN (500 mL) was heated for 40 hr at 65°C followed by addition of DMF (100 mL) and then continued heating for 24 hr. The cooled reaction mixture was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, 1N HCl, and brine, dried (MgSO₄), and concentrated. The residue was triturated with hexanes to give 5-2 as a yellow solid.
- TLC RF = 0.93 (EtOAc);
- ¹HNMR (300 MHz, CDCl₃) δ 3.68 (t, J = 7 Hz, 4H), 2.00 (m, 2H), 1.48 (s, 18 H).

Tert-Butyloxycarbonyl-tetrahydropyrimidine (5-3)

- A solution of 5-2 (19.0 g, 63 mmol), Mg(ClO₄)₂ (2.8 g, 12.7 mmol), and CH₃CN was heated at 50 °C for 2 hr. The cooled solution was diluted with CHCl₃ and then washed with 1N HCl, sat. NaHCO₃, and brine, dried (MgSO₄), and concentrated. Flash chromatography (silica, 75% EtOAc/ hexanes → EtOAc) gave 5-3 as a brown solid.
- TLC RF = 0.26 (silica, EtOAc);
- ¹HNMR (300 MHz, CDCl₃) δ 5.50 (bs, 1H), 3.70 (m, 2H), 3.29 (m, 2H), 1.97 (m, 2H), 1.48 (s, 9H).

Tert-Butyloxycarbonyl-2-oxo-3-(3-ethylene glycolbutyl)-tetrahydropyrimidine (5-4)

- To a stirred solution of 5-3 (3.2g, 16.1 mmol) and DMF (50 mL) was added LiN(TMS)₂ (21 mL, 1M/hexanes). After 20 minutes, the iodide 1-2 (8.6 g, 35.2 mmol) in DMF (10 mL) was added and the reaction mixture heated at 50 °C for 2 hours. The cooled solution was diluted with CHCl₃ and then washed with H₂O and brine, dried (MgSO₄), and concentrated. Flash chromatography (silica, 60% to 75% EtOAc/hexanes) gave 5-4 as an orange oil.
- TLC RF = 0.74 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);

- 111 -

¹H NMR (300 MHz, CDCl₃) δ 3.93 (s, 4H), 3.66 (t, J= 6Hz, 2H), 3.44 (m, 2H), 3.30 (m, 2H), 1.96 (m, 2H), 1.48 (s, 9H), 1.32 (s, 3H).

1-Oxo-2-(3-ethylene glycol-butyl) tetrahydro-pyrimidine (5-5)

- 5 A mixture of 5-4 (3.0 g, 9.5 mmol), TFA (1.5 mL, and toluene (30 mL) was stirred at ambient temperature for 20 minutes, concentrated and the residue azeotroped with toluene to remove excess TFA. The residue was then dissolved in toluene (30 mL) and treated with NaHCO₃ (3g), filtered, and the filtrate concentrated to give a
- 10 yellow oil. Flash chromatography (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH) gave 5-5 as a yellow oil.
- TLC RF = 0.63 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);
- ¹H NMR (300 MHz, CDCl₃) δ 5.16 (bs, 1H), 3.94 (s, 4H), 3.40 (m, 2H), 3.24 (m, 4H), 1.90 (m, 2H), 1.34 (s, 3H).

15

Ethyl 2-oxo-3-[3-ethylene glycol-butyl]tetrahydropyrimidin-1-yl-acetate (5-6)

- To a stirred solution of 5-5 (2.0 g, 9.3 mmol) and DMF (50 mL) was added LiN(TMS)₂ (12.1 mL, 1.0 M/THF). After 20 min,
- 20 ethyl iodoacetate (1.66 mL, 14.0 mmol) was added followed by heating at 60°C for 1 hr. The cooled solution was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, and brine, dried (MgSO₄), and concentrated. Flash chromatography (silica, 50% to 75% EtOAc/hexanes) gave 5-6 as a colorless oil.
- 25 TLC RF = 0.72 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);
- ¹H NMR (300 MHz, CDCl₃) δ 4.18 (q, J=7Hz, 2H), 3.93 (s, 4H), 3.42 (m, 2H), 3.34 (m, 4H), 1.98 (m, 2H), 1.92 (m, 2H), 1.34 (s, 3H), 1.25 (t, J=7Hz, 3H).

30 Ethyl 2-oxo-3-[3-oxo-butyl]tetrahydro-pyrimidin-1-yl-acetate (5-7)

A solution of 5-6 (750 mg, 2.5 mmol), p-TSA (10 mg), and acetone (30 mL) was refluxed for 1 hr. The cooled solution was diluted with CHCl₃ and then washed with sat. NaHCO₃ and brine, dried (MgSO₄), and concentrated to give 5-7 as a yellow oil.

- 112 -

TLC RF = 0.36 (silica, 10% CH₃OH/EtOAc);

¹H NMR (300 MHz, CDCl₃) δ 4.17 (q, J=7Hz, 2H), 3.56 (m, 2H), 3.34 (m, 4H), 2.76 (t, J=7Hz, 2H), 2.17 (s, 3H), 2.00 (m, 2H), 1.27 (t, J=7Hz, 3H).

5

Ethyl 2-oxo-3-[2-naphthyridin-2-yl]ethyl]-tetrahydropyrimidin-1-yl-acetate (5-8)

A mixture of 5-7 (600 mg, 2.3 mmol), 1-4 (343 mg, 2.8 mmol), L-proline (175 mg), and ethanol (25 mL) was heated at reflux for 18 hr. The cooled reaction mixture was concentrated and the residue purified by flash chromatography (silica, 10% CH₃OH/EtOAc) gave 5-8 as a yellow solid.

10

TLC RF = 0.21 (silica, 10% CH₃OH/EtOAc);

¹H NMR (300 MHz, CDCl₃) δ 9.10 (m, 1H), 8.19 (m, 1H), 8.14 (d, J=8Hz, 1H), 7.52 (d, J=8Hz, 1H), 7.44 (m, 1H), 4.18 (q, J=7Hz, 2H), 3.83 (m, 2H), 3.32 (m, 6H), 1.93 (m, 2H), 1.24 (t, J=7Hz, 3H).

15

Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]-naphthyridine-2-yl)ethyl]tetrahydropyrimidine-1-yl-acetate (5-9)

A mixture of 5-8 (600 mg, 1.75 mmol), 10% Pd/C (300 mg), and ethanol (10 mL) was stirred at ambient temperature under a hydrogen atmosphere (1 atm) for 20 hr. The catalyst was removed by filtration through a celite pad and the filtrate concentrated to give 5-9 as a yellow oil.

20

¹H NMR (300 MHz, CDCl₃) δ 7.04 (d, J=8Hz, 1H), 6.42 (d, J=8Hz, 1H), 4.80 (bs, 1H), 4.22-4.03 (m, 4H), 3.60 (m, 2H), 2.78 (m, 2H), 2.66 (m, 2H), 1.96 (m, 4H), 1.24 (t, J=7Hz, 3H).

25

2-Oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]naphthridin-2-yl)tetrahydropyrimidin-1-yl]-acetic acid (5-10)

30

A solution of 5-9 (600 mg, 1.73 mmol) and 6N HCl (20 mL) was heated at 50°C for 2 hr. The solution was concentrated followed by azeotropic removal of H₂O with CH₃CN to give 5-10 as a yellow solid.

- 113 -

¹H NMR (300 MHz, CD₃OD) δ 7.58 (d, J=8Hz, 1H), 6.63 (d, J=8Hz, 1H), 3.98 (s, 2H), 3.62 (t, J=7Hz, 2H), 3.50 (m, 2H), 3.36 (m, 4H), 2.93 (m, 2H), 2.80 (m, 2H), 2.00 (m, 4H).

5 Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine (5-11)

To a stirred solution of 5-10 (250 mg, 0.70 mmol), 1-9 (210 mg, 0.77 mmol), EDC (148 mg, 0.77 mmol), HOBT (95 mg, 0.70 mmol), CH₃CN (2 mL), and DMF (2 mL) was added NMM (542 μL, 4.9 mmol). After stirring at ambient temperature for 20 hr, the reaction mixture was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, brine, dried (MgSO₄), and concentrated. Flash chromatography (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH) gave 5-11 as a colorless oil.

15 TLC RF = 0.31 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);

¹H NMR (300 MHz, CDCl₃) δ 8.58 (m, 1H), 8.50 (m, 1H), 7.94 (m, 1H), 7.66 (m, 1H), 7.22 (m, 1H), 7.05 (d, J=8Hz, 1H), 6.40 (d, J=8Hz, 1H), 5.43 (m, 1H), 4.06 (q, J=7Hz, 2H), 4.02 (m, 1H), 3.90 (m, 1H), 3.60 (m, 2H), 3.39 (m, 2H), 3.29 (m, 2H), 3.19 (m, 2H), 2.88 (m, 2H), 2.77 (m, 2H), 2.70 (m, 2H), 1.90 (m, 4H), 1.16 (t, J=7Hz, 3H).

2-Oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3-(S)-pyridin-3-yl-β-alanine (5-12)


A mixture of 5-11 (100 mg, 0.22 mmol), 1N NaOH (300 μL), and ethanol (1 mL) was stirred at ambient temperature for 1 hr, followed by concentration. Flash chromatography (silica, 25:10:1:1 to 15:10:1:1 EtOAc/EtOH/NH₄OH/H₂O) gave 5-12 as a white solid.

TLC RF = 0.22 (silica, 10:10:1:1 EtOAc/ethanol/NH₄OH, H₂O);

¹H NMR (300 MHz, CD₃OD) δ 8.66 (m, 1H), 8.39 (m, 1H), 7.95 (m, 1H), 7.53 (d, J=8Hz, 1H), 7.40 (m, 1H), 6.66 (d, J=8Hz, 1H), 5.18 (m, 1H), 4.27 (d, J=7Hz, 1H), 4.16 (m, 1H), 3.64 (d, J=7Hz, 1H), 3.50-3.10 (m, 8H), 3.00-2.65 (m, 6H), 1.95 (m, 4H).


- 114 -

SCHEME 6



- 115 -

SCHEME 6 (cont'd)



- 116 -

1,3-Di-tert-butyloxycarbonyl-imidazolidin-2-one (6-2)

A heterogeneous mixture of 6-1 (10.0 g, 116 mmol), BOC₂O (56 g, 255 mmol), DMAP (20 mg), and CH₃CN (400 mL) was heated for 18 hr at 60°C. The cooled reaction mixture was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, 1N HCl, and brine, dried (MgSO₄), and concentrated. The residue was triturated with hexanes to give 6-2 as a white solid.

TLC RF = 0.91 (EtOAc);

¹H NMR (300 MHz, CDCl₃) δ 3.73 (s, 4H), 1.53 (s, 18 H).

Tert-Butyloxycarbonyl-imidazolidin-2-one (6-3)

A solution of 6-2 (28.0 g, 98 mmol), Mg(ClO₄)₂ (4.3 g, 20 mmol), and CH₃CN (400 mL) was heated at 50 °C for 3 hr. The cooled solution was diluted with CHCl₃ and then washed with 1N HCl, sat. NaHCO₃, and brine, dried (Mg SO₄), and concentrated. Flash chromatography (silica, 50% EtOAc/ hexanes →EtOAc) gave 6-3 as a yellow solid.

TLC RF = 0.31 (silica, EtOAc);

¹H NMR (300 MHz, CDCl₃) δ 6.27 (bs, 1H), 3.86 (m, 2H), 3.47 (m, 2H), 1.50 (s, 9H).

1-Tert-Butyloxycarbonyl-3-(3-ethylene glycol-butyl)imidazolidin-2-one (6-4)

To a stirred solution of 6-3 (4.5 g, 24 mmol) and DMF (50 mL) was added LiN(TMS)₂ (26.6 mL, 1M/hexanes). After 20 minutes, the iodide 1-2 (8.6 g, 35.2 mmol) in DMF (10 mL) was added and the reaction mixture heated at 60 °C for 4 hours. The cooled solution was diluted with CHCl₃ and then washed with H₂O and brine, dried (MgSO₄), and concentrated. Flash chromatography (silica, 75%

EtOAc/hexanes) gave 6-4 as an yellow solid.

TLC RF = 0.71 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);

¹H NMR (300 MHz, CDCl₃) δ 3.93 (s, 4H), 3.75 (m, 2H), 3.36 (m, 4H), 1.90 (m, 2H), 1.53 (s, 9H), 1.34 (s, 3H).

- 117 -

1-(3-Ethylene glycol-butyl)imidazolidin-2-one (6-5)

A mixture of 6-4 (4.0 g, 13.3 mmol), TFA (3 mL, and toluene (60 mL) was stirred at 50°C for 60 minutes, concentrated and the residue azeotroped with toluene to remove excess TFA. The residue
5 was then dissolved in toluene (30 mL) and treated with NaHCO₃ (3g), filtered, and the filtrate concentrated to give a yellow oil. Flash chromatography (silica, 70:25:5 CHCl₃/EtOAc/CH₃OH) gave 6-5 as a white solid.

TLC RF = 0.58 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);

10 ¹H NMR (300 MHz, CDCl₃) δ 4.25 (bs, 1H), 3.94 (s, 4H), 3.44 (m, 4H), 3.32 (m, 2H), 1.90 (m, 2H), 1.35 (s, 3H).

Ethyl 2-oxo-3-[3-ethylene glycol-butyl]imidazolidin-1-yl-acetate (6-6)

To a stirred solution of 6-5 (2.0 g, 10 mmol) and DMF (50
15 mL) was added LiN(TMS)₂ (11 mL, 1.0 M/THF). After 20 min, ethyl iodoacetate (3.5 mL, 30 mmol) was added at ambient temperature. After 3 hr the solution was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, and brine, dried (MgSO₄), and concentrated. Flash
20 chromatography (silica, 50% to 75% EtOAc/hexanes) gave 6-6 as a colorless oil.

TLC RF = 0.71 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);

25 ¹H NMR (300 MHz, CDCl₃) δ 4.18 (q, J=7Hz, 2H), 3.93 (s, 4H), 3.91 (m, 2H), 3.50-3.30 (m, 6H), 1.90 (m, 2H), 1.92 (m, 2H), 1.35 (s, 3H), 1.25 (t, J=7Hz, 3H).

Ethyl 2-oxo-3-[3-oxo-butyl]imidazolidin-1-yl-acetate (6-7)

A solution of 6-6 (1.4 g, 4.9 mmol), p-TSA (10 mg), and acetone (30 mL) was refluxed for 1 hr. The cooled solution was diluted with CHCl₃ and then washed with sat. NaHCO₃ and brine, dried
30 (MgSO₄), and concentrated to give 6-7 as a yellow oil.

TLC RF = 0.34 (silica, EtOAc);

¹H NMR (300 MHz, CDCl₃) δ 4.17 (q, J=7Hz, 2H), 3.94 (s, 2H), 3.48 (m, 2H), 3.42 (m, 4H), 2.72 (t, J=7Hz, 2H), 2.17 (s, 3H), 1.27 (t, J=7Hz, 3H).

- 118 -

Ethyl 2-oxo-3-[2-naphthyridin-2-yl]ethylimidazolidin-1-yl-acetate (6-8)

- A mixture of 6-7 (1.0 g, 4.1 mmol), 1-4 (604 mg, 4.9 mmol), L-proline (238 mg), and ethanol (50 mL) was heated at reflux for 20 hr. The cooled reaction mixture was concentrated and the residue purified by flash chromatography (silica, 70:25:5 CHCl₃/EtOAc/CH₃OH) gave 6-8 as a yellow oil.
- TLC RF = 0.42 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);
- ¹H NMR (300 MHz, CDCl₃) δ 9.10 (m, 1H), 8.19 (m, 1H), 8.14 (d, J=8Hz, 1H), 7.52 (d, J=8Hz, 1H), 7.44 (m, 1H), 4.17 (q, J=7Hz, 2H), 3.81 (m, 2H), 3.42 (m, 4H), 3.32 (m, 4H), 1.24 (t, J=7Hz, 3H).

Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]-naphthyridine-2-yl)ethylimidazolidin-1-yl-acetate (6-9)

- A mixture of 6-8 (1.1 g, 3.35 mmol), 10% Pd/C (500 mg), and ethanol (30 mL) was stirred at ambient temperature under a hydrogen atmosphere (1 atm) for 20 hr. The catalyst was removed by filtration through a celite pad and the filtrate concentrated to give 6-9 as a colorless oil.
- TLC RF = 0.11 (silica, 70:25:5 CHCl₃/EtOAc/CH₃OH);
- ¹H NMR (300 MHz, CDCl₃) δ 7.04 (d, J=8Hz, 1H), 6.42 (d, J=8Hz, 1H), 4.80 (bs, 1H), 4.22-4.03 (m, 4H), 3.96 (s, 2H), 3.55 (m, 2H), 3.40 (m, 2H), 2.78 (m, 2H), 2.68 (m, 2H), 1.90 (m, 2H), 1.24 (t, J=7Hz, 3H).

- 2-Oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]naphthridin-2-yl)imidazolidin-1-yl-acetic acid (6-10)

- A solution of 6-9 (1.0 g, 3.0 mmol) and 6N HCl (40 mL) was heated at 60°C for 1 hr. The solution was concentrated followed by azeotropic removal of H₂O with CH₃CN to give 6-10 as a yellow solid.
- ¹H NMR (300 MHz, CD₃OD) δ 7.58 (d, J=8Hz, 1H), 6.63 (d, J=8Hz, 1H), 3.98 (s, 2H), 3.50 (m, 4H), 3.36 (m, 4H), 2.93 (m, 2H), 2.82 (m, 2H), 1.97 (m, 4H).

- 119 -


Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)ethyl]imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine (6-11)

- To a stirred solution of 6-10 (240 mg, 0.70 mmol), 1-9 (207 mg, 0.77 mmol), EDC (269 mg, 1.4 mmol), HOBT (95 mg, 0.70 mmol), and CH₃CN (3 mL) was added NMM (619 μL, 5.6 mmol). After stirring at ambient temperature for 20 hr, the reaction mixture was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, brine, dried (MgSO₄), and concentrated. Flash chromatography (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH) gave 6-11 as a colorless oil.
- 10 TLC RF = 0.41 (silica, 70:15:15 CHCl₃/EtOAc/CH₃OH);
¹H NMR (300 MHz, CDCl₃) δ 8.58 (m, 1H), 8.50 (m, 1H), 7.94 (m, 1H), 7.66 (m, 1H), 7.22 (m, 1H), 7.05 (d, J=8Hz, 1H), 6.40 (d, J=8Hz, 1H), 5.43 (m, 1H), 4.06 (q, J=7Hz, 2H), 3.85 (m, 1H), 3.55 (m, 2H), 3.40 (m, 2H), 3.33 (m, 4H), 2.90 (m, 2H), 2.77 (m, 2H), 2.70 (m, 2H),
15 1.90 (m, 2H), 1.77 (m, 2H), 1.18 (t, J=7Hz, 3H).

2-Oxo-3-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3-(S)-pyridin-3-yl-β-alanine (6-12)

- A mixture of 6-11 (160 mg, 0.33 mmol), 1N NaOH (500 μL), and ethanol (1 mL) was stirred at ambient temperature for 1 hr, followed by concentration. Flash chromatography (silica, 25:10:1:1 to 15:10:1:1 EtOAc/EtOH/NH₄OH/H₂O) gave 6-12 as a white solid. TLC RF = 0.21 (silica, 10:10:1:1 EtOAc/ethanol/NH₄OH, H₂O);
¹H NMR (300 MHz, CD₃OD) δ 8.66 (m, 1H), 8.39 (m, 1H), 7.95 (m, 1H), 7.53 (d, J=8Hz, 1H), 7.40 (m, 1H), 6.66 (d, J=8Hz, 1H), 5.22 (m, 1H), 3.93 (d, J=17 Hz, 1H), 3.74 (d, J=17Hz, 1H), 4.00-3.20 (m, 9H), 3.00-2.65 (m, 6H), 1.89 (m, 4H).
- 20
25

- 120 -

SCHEME 7

- 121 -

Ethyl 2-oxo-3(R)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)ethylpyrrolidin-1-yl]acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine (7-2)

To a stirred solution of 4-1 (175 mg, 0.52 mmol), 7-1 (214 mg, 0.72 mmol; for preparation see US 5,321,034), EDC (197 mg, 1.0 mmol), HOBT (70 mg, 0.52 mmol), and CH₃CN (3 mL) was added
5 NMM (498 μ L, 4.1 mmol). After stirring at ambient temperature for 20 hr, the reaction mixture was diluted with EtOAc and then washed with H₂O, sat. NaHCO₃, brine, dried (MgSO₄), and concentrated. Flash chromatography (silica, 70:25:5 CHCl₃/EtOAc/CH₃OH) gave 7-2
10 as a white solid.

TLC RF = 0.11 (silica, 70:25:5 CHCl₃/ EtOAc/CH₃OH);

¹H NMR (300 MHz, CDCl₃) δ 8.29 (bs, 1H), 7.55 (d, J = 7 Hz, 1H), 7.36 (d, J = 7 Hz, 1H), 7.20-7.00 (m, 3H), 6.63 (d, J = 7 Hz, 1H), 6.39 (d, J = 7 Hz, 1H), 4.30 (m, 1H), 4.10 (q, J=7Hz, 2H), 3.94 (d, J = 17 Hz, 1H), 3.83 (d, J = 17 Hz, 1H), 3.36 (m, 4H), 2.80 (m, 2H), 2.69 (m, 3H),
15 2.53 (d, J= 6 Hz, 2H), 2.50 (m, 1H), 2.24 (m, 2H), 1.93 (m, 4H), 1.75 (m, 2H), 1.18 (t, J=7Hz, 3H).

20 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)ethylpyrrolidin-1-yl]acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine (7-3)

A mixture of 7-2 (60 mg, 0.11 mmol), 1N NaOH (132 μ L), and ethanol (1 mL) was stirred at ambient temperature for 1 hr, followed by concentration. Flash chromatography (silica, 25:10:1:1 15:10:1:1 EtOAc/EtOH/NH₄OH/H₂O) gave 7-3 as a white solid.


25 TLC RF = 0.12 (silica, 10:10:1:1 EtOAc/ethanol/NH₄OH/H₂O);

¹H NMR (300 MHz, CD₃OD) δ 7.52 (d, J = 7 Hz, 1H), 7.43 (d, J = 7 Hz, 1H), 7.30 (d, J = 8 Hz, 1H), 7.05 (m, 2H), 6.92 (m, 1H), 6.48 (d, J = 7 Hz, 1H), 4.54 (d, J = 17 Hz, 1H), 4.27 (m, 1H), 3.50-1.70 (m, 22H).


- 122 -

SCHEME 8


Synthesis of Radioligand for SPA Assay



- 123 -

SCHEME 8 (Cont'd)

- 124 -

SCHEME 8 (Cont'd)

- 125 -

N-(4-Iodo-phenylsulfonylamino)-L-asparagine (8-2)

- To a stirred solution of acid 8-1 (4.39 g, 33.2 mmol), NaOH (1.49 g, 37.2 mmol), dioxane (30 ml) and H₂O (30 ml) at 0°C was added pipsyl chloride (10.34 g, 34.2 mmol). After ~5 minutes, NaOH (1.49, 37.2 mmol) dissolved in 15 ml H₂O, was added followed by the removal of the cooling bath. After 2.0 h, the reaction mixture was concentrated. The residue was dissolved in H₂O (300 ml) and then washed with EtOAc. The aqueous portion was cooled to 0°C and then acidified with concentrated HCl. The solid was collected and then washed with Et₂O to provide acid 8-2 as a white solid.
- ¹H NMR (300 MHz, D₂O) δ 7.86 (d, 2H, J=8Hz), 7.48 (d, 2H, J=8Hz), 3.70 (m, 1H), 2.39 (m, 2H).

2(S)-(4-Iodo-phenylsulfonylamino)-β-alanine (8-3)

- To a stirred solution of NaOH (7.14 g, 181.8 mmol) and H₂O (40 ml) at 0°C was added Br₂ (1.30 ml, 24.9 mmol) dropwise over a ten minute period. After ~5 minutes, acid 8-2 (9.9 g, 24.9 mmol), NaOH (2.00 g, 49.8 mmol) and H₂O (35 ml) were combined, cooled to 0°C and then added in a single portion to the reaction. After stirring for 20 minutes at 0°C, the reaction was heated to 90°C for 30 minutes and then recooled to 0°C. The pH was adjusted to ~7 by dropwise addition of concentrated HCl. The solid was collected, washed with EtOAc, and then dried *in vacuo* to provide acid 8-3 as a white solid.
- ¹H NMR (300 MHz, D₂O) δ 8.02 (d, 2H, J=8Hz), 7.63 (d, 2H, J=8Hz), 4.36 (m, 1H), 3.51 (dd, 1H, J=5Hz, 13Hz) 3.21 (m, 1H).

Ethyl 2(S)-(4-iodo-phenylsulfonylamino)-β-alanine-hydrochloride (8-4)

- HCl gas was rapidly bubbled through a suspension of acid 8-3 (4.0 g, 10.81 mmol) in EtOH (50 ml) at 0°C for 10 minutes. The cooling bath was removed and the reaction was heated to 60°C. After 18 h, the reaction was concentrated to provide ester 8-4 as a white solid.
- ¹H NMR (300 MHz, CD₃OD) δ 7.98 (d, 2H, J=8Hz), 7.63 (d, 2H, J=8Hz), 4.25 (q, 1H, J=5Hz), 3.92 (m, 2H), 3.33 (m, 1H), 3.06 (m, 1H), 1.01 (t, 3H, J=7Hz).

Ethyl 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoate (8-5)

A mixture of ester 8-5a (700 mg, 2.63 mmol), (for preparation, see: Scheme 29 of PCT International Application Publication No. WO 95/32710, published December 7, 1995) 10% Pd/C (350 mg) and EtOH were stirred under 1 atm H₂. After 20 h, the reaction was filtered through a celite pad and then concentrated to provide ester 8-5 as a brown oil.

TLC R_f = 0.23 (silica, 40% EtOAc/hexanes)

- 10 ¹H NMR (300 MHz, CDCl₃) δ 7.95 (d, 2H, J=8Hz), 7.26 (m, 3H), 6.43 (d, 1H, J=7Hz), 6.35 (d, 1H, J=8Hz), 4.37 (m, 4H), 3.05 (m, 2H), 2.91 (m, 2H), 1.39 (t, 3H, J=7Hz).

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoic acid hydrochloride (8-6)

- 15 A suspension of ester 8-5 (625 mg, 2.31 mmol) in 6N HCl (12 ml) was heated to 60°C. After ~20 h, the reaction was concentrated to give acid 8-6 as a tan solid.

¹H NMR (300 MHz, CD₃OD) δ 7.96 (d, 2H, J=8Hz), 7.80 (m, 1H), 7.33 (d, 2H, J=8Hz), 6.84 (d, 1H, J=9Hz), 6.69 (d, 1H, J=7Hz), 3.09 (m, 4H).

20

Ethyl 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-iodo-phenyl)sulfonylamino)-β-alanine (8-7)

- 25 A solution of acid 8-6 (400 mg, 1.43 mmol), amine 8-4 (686 mg, 1.57 mmol), EDC (358 mg, 1.86 mmol), HOBT (252 mg, 1.86 mmol), NMM (632 μl, 5.72 mmol) and DMF (10 ml) was stirred for ~20 h. The reaction was diluted with EtOAc and then washed with sat NaHCO₃, brine, dried (MgSO₄) and concentrated. Flash chromatography (silica, EtOAc → 5% isopropanol/EtOAc) provided amide 8-7 as a white solid.

- 30 TLC R_f = 0.4 (silica, 10% isopropanol/EtOAc)

¹H NMR (300 MHz, CD₃OD) δ 7.79 (d, 2H, J=9Hz) 7.61 (d, 2H, J=8Hz), 7.52 (d, 2H, J=9Hz), 7.29 (m, 1H), 7.27 (d, 2H, J=8Hz), 4.20 (m, 1H), 3.95 (q, 2H, J=7Hz), 3.66 (dd, 1H, J=6Hz, 14Hz), 3.49 (dd, 1H, J=8Hz, 13Hz), 3.01 (m, 2H), 2.86 (m, 2H), 1.08 (t, 3H, J=7Hz).

- 127 -

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-iodophenyl-sulfonylamino)- β -alanine (8-8)

5 A solution of ester 8-7 (200 mg, 0.3213 mmol) and 6N HCl (30 ml) was heated to 60°C. After ~20 h, the reaction mixture was concentrated. Flash chromatography (silica, 20:20:1:1 EtOAc/EtOH/NH₄OH/H₂O) provided acid 8-8 as a white solid.

TLC R_f = 0.45 (silica, 20:20:1:1 EtOAc/EtOH/NH₄OH/H₂O)

10 ¹H NMR (400 MHz, DMSO) δ 8.40 (m, 1H), 8.14 (Bs, 1H), 7.81 (d, 2H, J=8Hz), 7.62 (d, 2H, J=8Hz), 7.48 (d, 2H, J=8Hz), 7.27 (m, 3H), 6.34 (d, 1H, J=7Hz), 6.25 (d, 1H, J=8Hz), 5.85 (bs, 2H), 3.89 (bs, 1H), 3.35 (m, 2H), 2.97 (m, 2H), 2.79 (m, 2H).

15 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-trimethylstannyl-phenylsulfonylamino)- β -alanine (8-9)

A solution of iodide 8-8 (70 mg, 0.1178 mmol), (CH₃Sn)₂ (49 μ l, 0.2356 mmol), Pd(PPh₃)₄ (5 mg) and dioxane (7 ml) was heated to 90°C. After 2 h, the reaction was concentrated and then purified by prep HPLC (Delta-Pak C₁₈ 15 μ M 100A°, 40 x 100 mm; 20 95:5 \rightarrow 5:95 H₂O/CH₃CN) provided the trifluoroacetate salt. The salt was suspended in H₂O (10 ml), treated with NH₄OH (5 drops) and then lyophilized to provide amide 8-9 as a white solid.

25 ¹H NMR (400 MHz, DMSO) δ 8.40 (m, 1H), 8.18 (d, 1H, J=8Hz), 7.67 (m, 5H), 7.56 (d, 2H, J=8Hz), 7.29 (d, 2H, J=8Hz), 6.95-7.52 (m, 2H), 6.45 (bs, 2H), 4.00 (m, 1H), 3.50 (m, 1H), 3.33 (m, 1H), 2.97 (m, 2H), 2.86 (m, 2H).

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-4-¹²⁵Iodo-phenylsulfonylamino- β -alanine (8-10)


30 An iodobead (Pierce) was added to a shipping vial of 5 mCi of Na¹²⁵I (Amersham, IMS30) and stirred for five minutes at room temperature. A solution of 0.1 mg of 8-9 in 0.05 mL of 10% H₂SO₄/MeOH was made and immediately added to the Na¹²⁵I/iodobead vial. After stirring for three minutes at room temperature,

- 128 -


approximately 0.04-0.05 mL of NH₄OH was added so the reaction mixture was at pH 6-7. The entire reaction mixture was injected onto the HPLC for purification [Vydac peptide-protein C-18 column, 4.6 x 250 mm, linear gradient of 10% acetonitrile (0.1% (TFA):H₂O (0.1% TFA) to 90% acetonitrile (0.1% TFA):H₂O (0.1% TFA) over 30 minutes, 1 mL/min]. The retention time of 8-10 is 17 minutes under these conditions. Fractions containing the majority of the radioactivity were pooled, lyophilized and diluted with ethanol to give approximately 1 mCi of 8-10, which coeluted on HPLC analysis with an authentic sample of 8-8.

Instrumentation: Analytical and preparative HPLC was carried out using a Waters 600E Powerline Multi Solvent Delivery System with 0.1 mL heads with a Rheodyne 7125 injector and a Waters 990 Photodiode Array Detector with a Gilson FC203 Microfraction collector. For analytical and preparative HPLC a Vydac peptide-protein C-18 column, 4.6 x 250 mm was used with a C-18 Brownlee modular guard column. The acetonitrile used for the HPLC analyses was Fisher Optima grade. The HPLC radiodetector used was a Beckman 170 Radioisotope detector. A Vydac C-18 protein and peptide column, 3.9 x 250 mm was used for analytical and preparative HPLC. Solutions of radioactivity were concentrated using a Speedvac vacuum centrifuge. Calibration curves and chemical concentrations were determined using a Hewlett Packard Model 8452A UV/Vis Diode Array Spectrophotometer. Sample radioactivities were determined in a Packard A5530 gamma counter.

- 129 -

SCHEME 9

- 130 -

SCHEME 9 (Cont'd)

- 131 -

Methyl (S)-(3-amino-2-oxo-pyrrolidin-1-yl)-acetic acid hydrochloride(9-2)

- A solution of 9-1 (0.50 g, 1.84 mmol) (prepared as described by Freidinger, R. M.; Perlow, D. S.; Veber, D. F.; *J. Org. Chem.*, 1982, 26, 104) in anhydrous ethyl acetate (50 mL) was cooled to 0°C and saturated with HCl gas, then stirred at 0°C for 2 h. The resulting colorless solution was concentrated at reduced pressure and the residue triturated with anhydrous diethyl ether giving 9-2 as a hygroscopic white solid.
- 10 ¹H NMR (300 MHz, CD₃OD) δ 4.16 (d, 2H); 4.2 (m, 1H); 3.68 (s, 3H); 3.53 (m, 2H); 2.58 (m, 1H); 2.09 (m, 1H).

Methyl 2-oxo-3(S)-[1,8]naphthyridin-2-ylmethyl)-amino]-pyrrolidin-1-yl]-acetic acid (9-4)

- 15 A solution of 9-2 (232 mg, 1.11 mmol) and 9-3 (176 mg, 1.11 mmol) (prepared as reported by Weissenfels, M.; Ulrici, B.; *Z. Chem.* 1978, 18, 20.) in anhydrous methanol (10 mL) was treated with NaOAc (91 mg, 1.11 mmol), NaBH₃CN (70 mg, 1.11 mmol) and powdered 4 Å molecular sieves (450 mg). The resulting mixture was
- 20 stirred at 0° for 3.5 h, then concentrated and the residue subjected to flash chromatography on silica gel (95:4.5:0.5 CH₂Cl₂/MeOH/NH₄OH) to afford 9-4 as a colorless glass.
- FAB MS (315, M⁺);
- 25 ¹H NMR (300 MHz, CD₃OD) δ 9.04 (d, 1H); 8.41 (dd, 1H); 8.38(d, 1H); 7.72 (d, 1H); 7.62 (dd, 1H); 4.31 (d, 2H); 4.21 (m, 2H); 3.68 (s, 3H); 3.63 (m, 1H); 3.53 (m, 2H); 2.52 (m, 1H); 1.95 (m, 1H).

Methyl [3(S)-[tert-butoxycarbonyl-[1,8]naphthyridin-2-ylmethyl)-amino]-2-oxo-pyrrolidin-1-yl]-acetic acid (9-5)

- 30 A solution of amine 9-4 (69 mg, 0.22 mmol) in THF (5 mL) was treated with Boc₂O (83 mg, 0.24 mmol) and stirred at room temperature for 18 h. The solvent was removed in vacuo and the resulting residue isolated by chromatography on silica gel (5% MeOH/CH₂Cl₂) to afford 9-5 as a yellow glass.

- 132 -

FAB MS (415, M+1);

¹H NMR (300 MHz, CD₃OD) δ 9.04 (d, 1H); 8.20 (m, 2H); 7.88 (d, 0.5H (rotamer a)); 7.82 (d, 0.5H (rotamer b)); 7.46 (m, 1H); 5.1-4.3 (m, 5H); 3.81 (m, 2H); 3.72 (s, 3H); 3.41 (m, 2H); 2.36 (m, 2H); 1.47 (s, 4.5 H (rotamer a)); 1.30 (s, 4.5 H, (rotamer b)).

Methyl 3(S)-[*tert*-butoxycarbonyl-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-aminol-2-oxo-pyrrolidin-1-yl]-acetic acid (9-6)

10 A solution of 9-5 (40 mg, 0.097 mmol) in EtOH (5 mL) was treated with 10% Pd on C (8 mg) and then stirred under a H₂ filled balloon for 16 h. The catalyst was removed by filtration through celite and the filtrate concentrated to afford 9-6 as a colorless glass.

¹H NMR (300 MHz, CD₃OD) δ 7.10 (d, 1H) 6.78 (d, 0.5H (rotamer a)); 6.62 (d, 0.5H (rotamer b)); 4.8-3.9 (m, 5H); 3.81 (m, 2H); 3.72 (s, 3H); 3.38 (m, 2H); 2.36 (m, 2H); 1.21 (s, 4.5 H (rotamer a)); 1.15 (s, 4.5 H, (rotamer b)).

3(S)-[*tert*-butoxycarbonyl-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-aminol-2-oxo-pyrrolidin-1-yl]-acetic acid (9-7)

20 A solution of 9-6 (38 mg, 0.091 mmol) in 50 % aqueous THF (2 mL) was treated with 1.0 N NaOH (95 mL, 0.095 mmol) and stirred at room temperature for 2 h. The reaction was neutralized with 1N HCl, evaporated, and the residue dissolved in MeOH (2.5 mL),
25 filtered and evaporated to afford 9-7 as a colorless glass.

¹H NMR (300 MHz, CD₃OD) δ 7.31 (d, 1H) 6.78 (br, d, 1H); 4.8-3.9 (m, 5H); 3.81 (m, 2H); 3.38 (m, 2H); 2.36 (m, 2H); 1.21 (s, 4.5 H (rotamer a)); 1.15 (s, 4.5 H, (rotamer b)).

30 Ethyl 3-(2-{2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-pyrrolidin-1-yl}-acetyl-amino)-3-(S)-pyridin-3-yl-propionic acid (9-8)

9-7 (43 mg, 0.093 mmol), 1-9 (25 mg, 0.093 mmol), EDC (18 mg, 0.093 mmol), HOBT (13 mg, 0.093 mmol), and N-methyl

- 133 -

morpholine (31 mL, 0.28 mmol) in anhydrous DMF (5 mL) was stirred at room temperature for 18 h, then concentrated in vacuo and the residue chromatographed on silica gel using 5% MeOH/CH₂Cl₂ as eluent affording 9-8 as a colorless glass.

- 5 ¹H NMR (300 MHz, CDCl₃) δ 8.61 (s, 1H); 8.45 (d, 1H); 8.00 (m, 1H); 7.68, (d, 1H); 7.21 (m, 1H); 7.17 (d, 1H); 5.56 (m, 1H); 4.75 (s, 2H); 4.45 (m, 2H); 4.05 (q, 2H); 3.95 (m, 1H); 3.5-3.3 (m, 4H); 2.92 (m, 1H); 2.87 (m, 1H); 2.74 (m, 2H); 2.35 (m, 2H); 1.92 (m, 2H); 1.36 (s, 9H); 1.21 (t, 3H).

10

3-(2-{2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-pyrrolidin-1-yl}-acetylamino)-3-(S)-pyridin-3-yl-propionic acid (9-9)

- 15 9-8 (25 mg, 0.043 mmol) was dissolved in 6 N HCl (2 mL) and stirred at room temperature for 16 h, then evaporated to afford 9-9 as a pale yellow solid.


FAB MS (453, M⁺);

- 20 ¹H NMR (300 MHz, CD₃OD) δ 9.00 (s, 1H); 8.81 (d, 1H); 8.79(m, 1H); 8.10 (m, 1H); 7.71 (d, 1H); 7.01 (m, 1H); 5.56 (m, 1H); 4.75 (s, 2H); 4.61 (m, 1H); 4.50 (m, 1H); 4.35 (m, 1H); 4.10 (s, 2H); 3.62 (m, 4H); 3.4 -3.0 (m, 2H); 2.8 (m, 2H); 2.70 (m, 1H); 2.45 (m 1H); 1.98 (m, 2H).

- 25 Following the procedure described in Scheme 10, bicyclic compounds such as 10-6 are readily prepared by one of ordinary skill in the art.


- 134 -

SCHEME 10




- 135 -


SCHEME 11



- 136 -

SCHEME 11 (CONT'D)

- 137 -

SCHEME 11 (CONT'D)

- 138 -

3(R)-phenyl-tetrahydro-pyrrolo[1,2(S)-cloxazol-5-one (11-2)

A mixture of alcohol (S)-5-(hydroxymethyl)-2-pyrrolidinone (11-1, Fluka) (5.0 g, 43.4 mmol), benzaldehyde (5.7 mL, 56.4 mmol), p-TSA (80 mg, 0.4340 mmol) and toluene (125 mL) was heated to reflux with azeotropic removal of water for 18 hours. The solution was concentrated. Flash chromatography (silica, 50% EtOAc/hexanes) gave 11-2 as a yellow oil.

TLC R_f = 0.21 (silica, 50% EtOAc/hexanes)

¹H NMR (300 MHz, CDCl₃) δ 7.29 → 7.46 (m, 5H), 6.34 (s, 1H), 4.24 (m, 1H), 4.16 (t, J=5.8 Hz, 1H), 3.49 (t, J=7.8 Hz, 1H), 2.82 (m, 1H), 2.55 (m, 1H), 2.39 (m, 1H), 1.97 (m, 1H).

6(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-3(R)-phenyl-tetrahydro-pyrrolo[1,2(S)-cloxazol-5-one (11-3)

To a stirred solution of 11-2 (7.0 g, 34.4 mmol), HMPA (30.0 mL, 172 mmol) and THF (150 mL) at -78°C was added LDA (18.9 mL, 37.8 mmol, 2.0 M in heptane/THF). After 10 minutes, the reaction was warmed to -15°C. After 20 min, 1-2 (8.3 g, 34.4 mmol), dissolved in 10 mL of THF, was added. After 2 h, the reaction was warmed to ambient temperature for 3.0 hours and then recooled to -15°C for 18 hours. The reaction was warmed to ambient temperature for 2 hours and then diluted with Et₂O, washed with H₂O, dried (MgSO₄) and concentrated. Flash chromatography (silica, 40% → 60% EtOAc/hexanes) gave 11-3 as an oil.

TLC R_f = 0.28 (silica, 50% EtOAc/hexanes)

¹H NMR (300 MHz, CDCl₃) δ 7.25 → 7.46 (m, 5H), 6.33 (s, 1H), 4.24 (m, 1H), 4.11 (m, 1H), 3.95 (s, 4H), 3.52 (t, J=7.3 Hz, 1H), 2.87 (m, 1H), 2.57 (m, 1H), 2.10 (m, 1H) 1.40 → 1.86 (m, 4H), 1.34 (s, 3H).

1-benzyl-5(S)-hydroxymethyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-pyrrolidin-2-one (11-4)

A mixture of 11-3 (2.0 g, 6.30 mmol) and 10% Pd/carbon (2.0 g) in EtOH (30 mL) was stirred under a balloon of hydrogen for 1.0 h. Following filtration and evaporative removal of the solvent, the

- 139 -

residue dissolved in benzene (30 mL), treated with TsOH (10 mg) and ethylene glycol (1.05 mL, 18.9 mmol) and then heated to reflux with azeotropic removal of water for 1 hour. The reaction was concentrated. Flash chromatography (silica, 70:23:7 CHCl₃/EtOAc/MeOH) gave **11-4** as an oil.

¹H NMR (300 MHz, CDCl₃) δ 7.30 (m, 5H), 4.64 (d, J=15 Hz, 1H), 4.25 (d, J=15 Hz, 1H), 3.95 (s, 4H), 3.72 (m, 1H), 3.49 (m, 2H), 2.46 (m, 1H), 2.15 (m, 2H), 1.74 (m, 2H), 1.53 (m, 2H), 1.35 (s, 3H).

10 **1-benzyl-5(S)-iodomethyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-pyrrolidin-2-one (11-5)**

To a stirred solution of **11-4** (2.0 g, 6.26 mmol), PPh₃ (2.63 g, 10.0 mmol), imidazole (725 mg, 10.6 mmol) and CH₃CN (30 mL) at 0°C was added I₂ (2.39 g, 9.39 mmol) in five portions over 15 minutes. After 20 minutes, the reaction was warmed to 50°C for 30 minutes and then poured into 200 mL 1:1 EtOAc/hexanes. The solution was washed with 10% sodium bisulfite, sat NaHCO₃, brine, dried (MgSO₄) and concentrated. Flash chromatography (silica, 40% EtOAc/hexanes) gave **11-5** as an oil.

20 TLC R_f = 0.27 (silica, 50% EtOAc/hexanes)

¹H NMR (300 MHz, CDCl₃) δ 7.19 → 7.35 (m, 5H), 5.04 (d, J=15.1 Hz, 1H), 3.96 (m, 5H), 3.30 (m, 1H), 3.19 (m, 2H), 2.50 (m, 1H), 2.32 (m, 1H), 2.12 (m, 1H), 1.79 (m, 2H), 1.58 (m, 1H), 1.36 (m, 4H).

25 **1-benzyl-5(R)-methyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-pyrrolidin-2-one (11-6)**

To a stirred solution of **11-5** (900 mg, 6.26 mmol) and HMPA (30 mL) was added NaBH₄ (156 mg, 4.20 mmol). After 45 minutes, the reaction was poured into 50 mL 1:1 Et₂O/hexanes and then washed with H₂O, brine, dried (MgSO₄) and concentrated to provide **11-6** as an oil.

30 TLC R_f = 0.34 (silica, 50% EtOAc/hexanes)

- 140 -

¹H NMR (300 MHz, CDCl₃) δ 7.20 → 7.33 (m, 5H), 4.95 (d, J=15.1 Hz, 1H), 4.03 (d, J=14.9 Hz, 1H), 3.95 (s, 4H), 3.41 (m, 1H), 2.38 (m, 2H), 2.10 (m, 1H), 1.75 (m, 2H), 1.48 (m, 1H), 1.35 (s, 3H), 1.16 (m, 4H).

5 5(R)-methyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-pyrrolidin-2-one (11-7)

Into a 3-necked 500mL flask at -78°C was condensed 200 mL of ammonia. Lithium (64 mg, 9.25 mmol) was washed with MeOH, then THF and then added to the ammonia. After 20 minutes, 11-6 (560 mg, 1.85 mmol), dissolved in 25mL of THF, was added. After 30 minutes, the reaction was quenched with NH₄Cl; 200 mL of THF was added, the cooling bath was removed and the solution purged with argon for 30 minutes to remove the ammonia. The solution was dried (MgSO₄) and concentrated. Flash chromatography (silica, EtOAc → 5% MeOH/ EtOAc) gave 11-7 as an oil.

10 TLC R_f = 0.33 (silica, 10% MeOH/ EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 5.98 (br s, 1H), 3.94 (s, 4H), 3.67 (m, 1H), 2.40 (m, 2H), 2.02 (m, 1H), 1.70 (m, 2H), 1.40 (m, 1H), 1.33 (s, 3H), 1.22 (m, 4H).

20

{5(R)-methyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-2-oxo-pyrrolidin-1-yl}-acetic acid ethyl ester (11-8)

To a stirred solution of 11-7 (355 mg, 1.67 mmol) and THF (10 mL) at -78°C was added NaN(TMS)₂ (1.83 mL, 1.83 mmol, 1.0 M in THF). After 20 min, ethyl bromoacetate (0.203 mL, 1.84 mmol) was added and the reaction was warmed to 0°C. After 30 minutes, the reaction mixture was diluted with EtOAc and then washed with H₂O, brine, dried (MgSO₄), and concentrated to give 11-8 as a yellow oil.

25 TLC R_f = 0.90 (silica, 10% MeOH/ EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 4.35 (d, J=17.6 Hz, 1H), 4.18 (q, J=7.1 Hz, 2H), 3.94 (s, 4H), 3.74 (m, 3H), 2.44 (m, 2H), 2.05 (m, 1H), 1.73 (m, 2H), 1.43 (m, 1H), 1.33 (s, 3H), 1.27 (t, J=7.1 Hz, 3H), 1.20 (d, J=6.4 Hz, 3H).

30

- 141 -

[5(R)-methyl-2-oxo-3(S)-(3-oxo-butyl)-pyrrolidin-1-yl]-acetic acid ethyl ester (11-9)

5 A solution of 11-10 (360 mg, 1.20 mmol), p-TSA (10 mg) and acetone (20 mL) was heated at reflux for 1 hr. The cooled reaction mixture was diluted with EtOAc and then washed with sat. NaHCO₃ and brine, dried (MgSO₄), and concentrated to afford 11-9 as an oil.

TLC R_f = 0.54 (silica, 75%EtOAc/hexanes)

10 ¹H NMR (300 MHz, CDCl₃) δ 4.32 (d, J=17.6 Hz, 1H), 4.18 (q, J=7.1Hz, 2H), 3.73 (m, 3H), 2.72 (m, 2H), 2.42 (m, 1H), 2.16 (s, 3H), 1.99 (m, 1H), 1.78 (m, 1H), 1.27 (t, J=7.1Hz, 3H), 1.20 (d, J=6.1 Hz, 3H).

15 [5(R)-methyl-3(S)-(2-[1,8]naphthyridin-2-yl-ethyl)-2-oxo-pyrrolidin-1-yl]-acetic acid ethyl ester (11-10)

A mixture of 11-9 (220 mg, 0.8619 mmol), 1-4, 2-amino-3-formylpyridine (137 mg, 1.12 mmol) and proline (99 mg, 0.8619 mmol) in absolute ethanol (5 mL) was heated at reflux for 12 h.

20 Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:25:5 chloroform/ethyl acetate/MeOH) to give 11-10 as a yellow oil.

TLC R_f = 0.37 (70:25:5 chloroform/ethyl acetate/MeOH).

25 ¹H NMR (300 MHz, CDCl₃) δ 9.08 (m, 1H), 8.16 (dd, J=2Hz, 6 Hz 1H), 8.12 (d, J=8Hz, 1H), 7.46 (m, 2H), 4.33 (d, J=17.5 Hz, 1H), 4.17 (m, 2H), 3.71 (m, 3H), 3.21 (t, J=8.0 Hz, 2H), 2.54 (m, 2H), 2.39 (m, 1H), 2.02 (m, 1H), 1.35 (m, 1H), 1.26 (t, J=7.1 Hz, 3H), 1.21 (d, J=6.3 Hz, 3H).

30 [5(R)-methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl]-acetic acid ethyl ester (11-11)

A mixture of 1-10 (250 mg, 0.7323 mmol) and 10% Pd/carbon (250 mg) in EtOH (5 mL) was stirred under a balloon of hydrogen for 20 h. Following filtration and evaporative removal of the

- 142 -

solvent, the residue was chromatographed (silica gel, 70:25:5 chloroform/ethyl acetate/MeOH) to give 11-11 as a colorless oil. TLC Rf = 0.25 (70:25:5 chloroform/ethyl acetate/MeOH).

¹H NMR (300 MHz, CDCl₃) δ 7.05 (d, 1H, J=7.3 Hz), 6.39 (d, 1H, J=7.3 Hz), 4.77 (br s, 1H), 4.17 (d, 1H, J=17.5 Hz), 4.15 (m, 2H), 3.71 (m, 2H), 3.39 (m, 2H), 2.64 (m, 4H), 2.46 (m, 2H), 2.30 (m, 1H), 1.91 (m, 2H), 1.88 (m, 1H), 1.26 (t, 3H, J=6.1 Hz), 1.23 (m, 1H), 1.19 (d, J=6.4 Hz, 3H).

10 {5(R)-methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-acetic acid hydrochloride (11-12)

A mixture of 11-11 (185 mg, 0.5356 mmol) and 6N HCl (10 mL) was heated at 60° C for 1 h. Evaporative removal of the solvent gave 11-12 as a yellow solid.

15 ¹H NMR (300 MHz, CD₃OD) δ 7.59 (d, 1H, J=7.3 Hz), 6.66 (d, 1H, J=7.3 Hz), 4.17 (d, 12H, J=17.8, Hz), 3.90 (d, 1H, J=17.8, Hz), 3.77 (m, 1H), 3.50 (t, J=5.4 Hz, 2H), 3.31 (m, 4H), 2.52 (m, 2H), 2.25 (m, 1H), 1.95 (t, 2H, J=6.6 Hz), 1.80 (m, 1H), 1.34 (m, 1H), 1.25 (d, J=6.3 Hz, 3H)

20

2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-ethyl]pyrrolidin-1-yl}acetyl-3(S)-alkynyl-β-alanine ethyl ester (11-13)

A mixture of 11-12 (350 mg, 0.9892 mmol), 2-9 (193 mg, 1.09 mmol), EDC (378 mg, 1.98 mmol), HOBT (134 mg, 0.9892 mmol) and NMM (1.10 mL, 7.91 mmol) in CH₃CN (5 mL) was stirred for 20 h. The mixture was diluted with ethyl acetate, washed with sat. NaHCO₃, brine, and dried over MgSO₄. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:25:5 chloroform/ethyl acetate/MeOH) to give 11-13 as a colorless foam.

25 TLC Rf = 0.15 (70:25:5 chloroform/ethyl acetate/MeOH).

30 ¹H NMR (300 MHz, CDCl₃) δ 7.05 (m, 2H), 6.39 (d, 1H, J=7.3 Hz), 5.04 (m, 1H), 4.16 (q, 2H, J=7.1 Hz), 3.90 (s, 2H), 3.64 (m, 1H), 3.39 (m, 2H), 2.69 (m, 6H), 2.47 (m, 2H), 2.30 (m, 1H), 1.90 (m, 2H), 1.64 (m, 2H), 1.20 (m, 7H).


- 143 -

2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethylpyrrolidin-1-yl]acetyl-3(S)-alkynyl- β -alanine (11-14)


- To a solution of 11-13 (70 mg, 0.1589 mmol) in EtOH (1 mL) was added 1N NaOH (0.175 ml, 0.164 mmol). After stirring for 1 h, the solvents were evaporated and the residue was chromatographed (silica gel, 25:10:1:1 to 15:10:1:1 ethyl acetate/EtOH/water/NH₄OH to give 11-14 as a colorless foam.
- TLC R_f = 0.21 (10:10:1:1 ethyl acetate/EtOH/water/NH₄OH).
- ¹H NMR (300 MHz, CD₃OD) δ 7.42 (d, 1H, J=7.3 Hz), 6.49 (d, 1H, J=7.3 Hz), 4.35 (d, J=17.1 Hz, 1H), 3.64 (m, 1H), 3.50 (m, 3H), 3.18 (m, 2H), 2.77 (t, J=5.6 Hz, 2H), 2.55 (m, 5H), 2.23 (m, 1H), 1.91 (m, 4H), 1.41 (m, 1H) 1.28 (d, J=6.3 Hz, 3H).

- 144 -


SCHEME 12



- 145 -

SCHEME 12 (CONT'D)

- 146 -

SCHEME 12 (CONT'D)

- 147 -

1-benzyl-5(S)-methyl-p-toluenesulfonate-3(S)-[2-(2-methyl-
[1,3]dioxolan-2-yl)-ethyl]-pyrrolidin-2-one (12-1)

To a stirred solution of 11-4 (1.8 g, 5.63 mmol) and THF (30 mL) at 0°C was added NaH (248 mg, 6.19 mmol). After 30
5 minutes, TosCl was added followed by the removal of the cooling bath. After 1.0 hour, the reaction was diluted with EtOAc and then washed with H₂O, sat NaHCO₃, brine, dried (MgSO₄) and concentrated. Flash chromatography (silica, 40 → 60% EtOAc/hexanes) gave 12-1 as an oil. TLC R_f = 0.75 (silica, EtOAc)
10 ¹H NMR (300 MHz, CDCl₃) δ 7.72 (d, J=8.30 Hz, 2H), 7.35 (d, J=7.3 Hz, 2H), 7.25 (m, 3H), 7.09 (m, 2H), 4.94 (d, J=14.9 Hz, 1H), 4.01 (m, 1H), 3.94 (m, 5H), 3.83 (d, J=15.1 Hz, 1H), 3.54 (m, 1H), 2.46 (s, 3H), 2.42 (m, 1H), 2.21 (m, 1H), 2.01 (m, 1H), 1.72 (m, 2H), 1.43 (m, 2H), 1.32 (s, 3H).

15 1-benzyl-5(S)-benzyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-
pyrrolidin-2-one (12-2)

To a stirred suspension of CuI (2.57 g, 13.5 mmol) and Et₂O (10 mL) at 0°C was added PhLi (14.2 mL, 25.6 mmol, 1.8 M cyclohexane-ether) dropwise over a 1.0 hour period. After an
20 additional hour, 12-1 (1.4 g, 2.96 mmol), dissolved in 10 mL Et₂O, was added. The reaction was stirred at -15°C for 96 hours. The reaction was diluted with EtOAc and then washed with sat NH₄Cl, sat NaHCO₃, brine, dried (MgSO₄) and concentrated. Flash chromatography (silica,
25 30 → 60% EtOAc/hexanes) gave 12-2 as an oil. TLC R_f = 0.29 (silica, 50% EtOAc/hexanes)
¹H NMR (300 MHz, CDCl₃) δ 7.13 → 7.36 (m, 8H), 7.02 (d, J=7.6 Hz, 2H), 5.06 (d, J=14.9 Hz, 1H), 4.14 (d, J=15.1 Hz, 1H), 3.95 (m, 4H), 3.55 (m, 1H), 3.18 (dd, J=4.2, 17.0 Hz, 1H), 2.35 (m, 2H), 2.04 (m, 2H),
30 1.66 (m, 2H), 1.32 (m, 5H).

- 148 -

5(S)-benzyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-pyrrolidin-2-one (12-3)

Into a 3-necked 500mL flask at -78°C was condensed 100 mL of ammonia. Next, 12-2 (470 mg, 1.24 mmol), dissolved in 20mL of THF, was added. Lithium (19 mg, 2.48 mmol) was washed with MeOH, then THF and then added to the ammonia. After 20 minutes, the reaction was quenched with NH₄Cl; 200 mL of THF was added, the cooling bath was removed and the solution purged with argon for 30 minutes to remove the ammonia. The solution was dried (MgSO₄) and concentrated. Flash chromatography (silica, EtOAc) gave 12-3 as an oil. TLC R_f = 0.22 (silica, EtOAc)
¹H NMR (300 MHz, CDCl₃) δ 7.18 → 7.35 (m, 5H), 5.43 (br s, 1H), 3.95 (s, 4H), 3.92 (m 1H), 2.88 (dd, J=5.3, 18.6 Hz, 1H), 2.41 (m, 2H), 2.03 (m, 1H), 1.71 (m, 2H), 1.43 (m, 2H), 1.33 (s, 3H).

{5(S)-benzyl-3(S)-[2-(2-methyl-[1,3]dioxolan-2-yl)-ethyl]-2-oxo-pyrrolidin-1-yl}-acetic acid ethyl ester (12-4)

To a stirred solution of 12-3 (210 mg, 0.7257 mmol) and THF (5 mL) at -78°C was added NaN(TMS)₂ (0.943 mL, 0.943 mmol, 1.0 M in THF). After 30 min, ethyl bromoacetate (0.104 mL, 0.9434 mmol) was added and the reaction was warmed to 0°C. After 1.0 hour, the reaction mixture was diluted with EtOAc and then washed with sat NaHCO₃, brine, dried (MgSO₄), and concentrated to give 12-4 as a yellow oil.

TLC R_f = 0.64 (silica, EtOAc)

¹H NMR (300 MHz, CDCl₃) δ 7.15 → 7.33 (m, 5H), 4.40 (d, J=17.8 Hz, 1H), 4.15 (m, 2H), 3.93(m, 5H), 3.77 (d, J=17.8, 1H), 3.07 (dd, J=5.0, 18.6 Hz, 1H), 2.56 (m, 1H), 2.39 (m, 1H), 2.20 (m, 1H), 2.05 (m, 1H), 1.69 (m, 2H), 1.23 → 1.46 (m, 8H).

[5(S)-benzyl-2-oxo-3(S)-(3-oxo-butyl)-pyrrolidin-1-yl]-acetic acid ethyl ester (12-5)

A solution of 12-4 (260 mg, 0.6925 mmol), p-TSA (10 mg) and acetone (20 mL) was heated at reflux for 1 hr. NaHCO₃ was added

- 149 -

to the cooled reaction mixture and then the mixture was concentrated. The residue was diluted with CHCl_3 and then washed with brine, dried (MgSO_4), and concentrated to afford 12-5 as an oil.

TLC R_f = 0.66 (silica, 75%EtOAc/hexanes)

- 5 ^1H NMR (300 MHz, CDCl_3) δ 7.22 \rightarrow 7.36 (m, 3H), 7.15 (d, $J=6.5$ Hz, 2H), 4.37 (d, $J=17.6$ Hz, 1H), 4.18 (m, 2H), 3.97 (m, 1H), 3.77 (d, $J=17.8$ Hz, 1H), 3.06 (dd, $J=5, 18$ Hz, 1H), 2.60 (m, 3H), 2.42 (m, 1H), 2.17 (m, 1H), 2.14 (s, 3H), 1.96 (m, 1H), 1.74 (m, 1H) 1.27 (m, 4H).

- 10 [5(S)-benzyl-3(S)-(2-[1,8]naphthyridin-2-yl-ethyl)-2-oxo-pyrrolidin-1-yl]-acetic acid ethyl ester (12-6)

A mixture of 12-5 (230 mg, 0.6940 mmol), 1-4, (2-amino-3-formylpyridine, 110 mg, 0.9022 mmol) and proline (80 mg, 0.6940 mmol) in absolute ethanol (10 mL) was heated at reflux for 18 h.

- 15 Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:28:2 chloroform/ethyl acetate/MeOH) to give 12-6 as a yellow oil.

TLC R_f = 0.38 (70:25:5 chloroform/ethyl acetate/MeOH).

- 20 ^1H NMR (300 MHz, CDCl_3) δ 9.08 (m, 1H), 8.16 (dd, $J=2$ Hz, 10 Hz, 1H), 8.09 (d, $J=8.3$ Hz, 1H), 7.44 (m, 2H), 7.28 (m, 2H), 7.16 (d, $J=8$ Hz, 2H), 4.37 (d, $J=17.6$ Hz, 1H), 4.16 (m, 2H), 3.96 (m, 1H), 3.80 (d, $J=17.6$ Hz, 1H), 3.15 (m, 2H), 3.06 (dd, $J=5.3, 18.5$, 1H), 2.26 \rightarrow 2.63 (m, 4H), 1.97 (m, 1H), 1.47 (m, 1H), 1.25 (t, $J=7.1$ Hz, 3H).

- 25 [5(S)-benzyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl]-acetic acid ethyl ester (12-7)

A mixture of 12-6 (220 mg, 0.5270 mmol) and 10% Pd/carbon (100 mg) in EtOH (4 mL) was stirred under a balloon of hydrogen for 2 h. Following filtration and evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:25:5 chloroform/ethyl acetate/MeOH) to give 12-7 as a colorless oil.

30 TLC R_f = 0.25 (70:25:5 chloroform/ethyl acetate/MeOH).

- ^1H NMR (300 MHz, CDCl_3) δ 7.26 (m, 3H), 7.16 (d, $J=8.1$ Hz, 2H), 7.04 (d, $J=7.3$ Hz, 1H), 6.36 (d, $J=7.3$ Hz, 1H), 4.74 (br s, 1H), 4.39 (d,

- 150 -

J=17.8 Hz, 1H), 4.15 (m, 2H), 3.90 (m, 1H), 3.77 (d, J=17.5 Hz, 1H), 3.38 (m, 2H), 3.06 (dd, J=2.4, 18.8 Hz, 1H), 2.65 (m, 5H), 2.43 (m, 1H), 2.22 (m, 3H), 1.89 (m, 1H) 1.36 (m, 1H), 1.26 (t, J=7.1 Hz, 3H).

5 (5(s)-benzyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]pyrrolidin-1-yl)-acetic acid hydrochloride (12-8)

A mixture of 12-7 (150 mg, 0.3559 mmol) and 6N HCl (10 mL) was heated at 60° C for 1 h. Evaporative removal of the solvent gave 12-8 as a yellow solid.

10 ¹H NMR (300 MHz, CD₃OD) δ 7.57 (d, J=7.3 Hz, 1H), 7.24 (m, 5H), 6.60 (d, J=7.3 Hz, 1H), 4.24 (d, J=17.8, Hz, 1H), 4.03 (m, 2H), 3.49 (t, J=5.6 Hz, 2H), 3.15 (dd, J=4.4, 17.6 Hz, 1H), 2.71 (m, 5H), 2.46 (m, 1H), 2.21 (m, 1H), 1.97 (m, 3H), 1.64 (m, 1H), 1.45 (m, 1H)

15 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester (12-9)

A mixture of 12-8 (150 mg, 0.3559 mmol), 2-10 (60 mg, 0.2135 mmol), EDC (132 mg, 0.7118 mmol), HOBT (48 mg, 0.3559 mmol) and NMM (0.4 mL, 2.85 mmol) in DMF (4 mL) was stirred for 20 h. The mixture was diluted with ethyl acetate, washed with sat. NaHCO₃, brine, and dried over MgSO₄. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 70:20:10 chloroform/ethyl acetate/MeOH to give 12-9 as a colorless foam.

25 TLC R_f = 0.15 (70:25:5 chloroform/ethyl acetate/MeOH).
¹H NMR (300 MHz, CD₃OD) δ 8.55 (s, 1H), 8.44 (m, 1H), 7.82 (m, 1H), 7.6-7.1 (m, 7H), 6.33 (d, J=7.5 Hz, 1H), 5.40 (t, J=8 Hz, 1H), 4.2-3.8 (m, 6H), 3.38 (m, 1H), 3.17 (m, 2H), 2.90 (m, 2H), 2.67 (m, 2H), 2.54 (m, 2H), 2.12 (m, 2H), 1.84 (m, 2H), 1.43 (m, 2H) 1.18 (m, 3H).

30 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine (12-13)

To a solution of 12-9 (70 mg, 0.1229 mmol) in EtOH (1 mL) was added 1N NaOH (0.150 ml, 0.150 mmol). After stirring for


- 151 -

1.5 h, the solvents were evaporated and the residue was chromatographed (silica gel, 25:10:1:1 to 15:10:1:1 ethyl acetate/EtOH/water/NH₄OH to give 12-10 as a colorless foam. TLC R_f = 0.21 (10:10:1:1 ethyl acetate/EtOH/water/NH₄OH).






- 5 ¹H NMR (300 MHz, CD₃OD) δ 8.62 (s, 1H), 8.36 (m, 1H), 7.92 (m, 1H), 7.45-7.2 (m, 7H), 6.49 (d, J=7.1 Hz, 1H), 5.27 (m, 1H), 4.31 (d, J=17.3 Hz, 1H), 3.93 (m, 1H), 3.72 (d, J=17.5 Hz, 1H), 3.30 (m, 3H), 2.92 -2.52 (m, 8H), 2.36 (m, 2H), 1.90 (m, 3H), 1.57 (m, 1H).

- 152 -





SCHEME 13



- 153 -

SCHEME 13. CONT'DEDC, HOBT
NMM

- 154 -

SCHEME 13 (CONT'D)

- 155 -

1-benzyl-5(S)-hydroxymethyl-pyrrolidin-2-one (13-1)

A mixture of 11-2 (5.0 g, 24.6 mmol), 10% Pd/C (2.5 g), and ethanol (80 mL) was stirred at ambient temperature under a hydrogen atmosphere (1 atm) for 5 hr. The catalyst was removed by
5 filtration through a celite pad and the filtrate concentrated to give 13-1 as a colorless oil.

TLC RF = 0.55 (silica, 70:20:10 CHCl₃/EtOAc/CH₃OH);

¹H NMR (300 MHz, CDCl₃) δ 7.29 (m, 5H), 4.83 (d, 2H, J=15Hz), 4.25 (d, 1H, J=15Hz), 3.77 (m, 1H), 3.51 (m, 2H), 2.54 (m, 1H), 2.40 (m,
10 1H), 1.92 (m, 2H).

1-benzyl-5(S)-iodomethyl-pyrrolidin-2-one (13-2)

To a solution of 13-1 (18.5 g, 90.1 mmol), triphenylphosphine (40.1 g, 153 mmol), and imidazole (11.03 g, 162
15 mmol) in 225 mL of acetonitrile and 150 mL of ether at 0°C was added iodine (34.3 g, 135 mmol) in 5 portions over 5 minutes. After 10 minutes, the reaction was heated to 50°C, and a stream of argon passed over the reaction to purge the evaporating ether. After an additional 30 minutes, the mixture was diluted with ether, the organic layer washed
20 with NaHCO₃ (sat.) and brine, dried over K₂CO₃, and the solvent evaporated. Flash chromatography of the residue (silica, 7-15% EtOAc/CHCl₃) gave 13-2 as a yellow oil.

TLC R_f = 0.53 (silica, 30% EtOAc/CHCl₃)

¹H NMR (300 MHz, CDCl₃) δ 7.31 (m, 5H), 5.05 (d, 1H, J=15Hz), 3.92
25 (d, 1H, J=15Hz), 3.41 (m, 1H), 3.26 (m, 2H), 2.62 (m, 1H), 2.43 (m, 1H), 2.16 (m, 1H), 1.81 (m, 1H).

1-benzyl-5(R)-methyl-pyrrolidin-2-one (13-3)

To a solution of 13-2 (22.1 g, 70 mmol) in 200 mL of
30 hexamethylphosphorous triamide at 0°C was added NaBH₄ (5.25 g, 140 mmol) in 5 portions over 5 minutes. After 10 minutes, the reaction was allowed to warm to ambient temperature and stirred for 2 h. The mixture was diluted with 1:1 ether/hexanes, quenched by the careful

- 156 -

addition of 300 mL 10% KHSO₄ (aq), separated, the organics dried over K₂CO₃, and the solvent evaporated to give 13-2 as a yellow oil
TLC R_f = 0.45 (silica, 30% EtOAc/CHCl₃)

¹H NMR (300 MHz, CDCl₃) δ 7.32 (m, 5H), 4.95 (d, 1H, J=15Hz), 4.00
5 (d, 1H, J=15Hz), 3.52 (m, 1H), 2.46 (m, 2H), 2.15 (m, 1H), 1.60 (m, 1H), 1.16 (d, 3H, J=6.0Hz)).

3(R)-azido-1-benzyl-5(R)-methyl-pyrrolidin-2-one (13-4)

To a solution of 13-3 (2.2 g, 11.6 mmol) in THF (45 mL)
10 at -78° C was added a solution of LDA (6.39 mL, 12.8 mmol;
2M/THF,ethylbenzene). The mixture was warmed to -15°C for 20
minutes, then recooled to -78°C, and 2,4,6-triisopropylbenzenesulfonyl
azide (4.31 g, 13.9 mmol, prepared as described in Harmon, *et al*, *J.*
Org. Chem. 1973, 38, 11-16.) was added rapidly as a solution in 40 mL
15 THF at -78°C. After 10 minutes, glacial acetic acid (2.67 mL, 47 mmol)
was added, and the resultant viscous liquid mixture allowed to warm to
ambient temperature and stir for 1 hour. The solvent was then
evaporated, the residue dissolved in CHCl₃, washed with NaHCO₃ (sat.),
and dried over magnesium sulfate. Following evaporative removal of
20 the solvent, the residue was chromatographed (silica gel, 25% ethyl
acetate/hexanes) to give 13-4 as a colorless oil.
TLC R_f = 0.38 (25% ethyl acetate/hexanes).
¹H NMR (300 MHz, CHCl₃) δ 7.32 (m, 5H), 5.00 (d, 1H, J=15 Hz),
4.27 (t, 1H, J=7.5 Hz), 3.98 (d, 1H, J=15 Hz), 3.54 (m, 1H), 1.97 (m,
25 2H), 1.16 (d, 3H, J=6.0 Hz).

3(S)-azido-1-benzyl-5(R)-methyl-pyrrolidin-2-one

To a solution of 13-4 (2.17 g, 9.42 mmol) in EtOH (50
mL) was added a solution of NaOEt (3.52 mL, 9.42 mmol; 2.68
30 M/EtOH). The mixture was stirred for 90 minutes, then quenched by
the addition of glacial acetic acid (3 mL). The solvent was then
evaporated, the residue slurried in EtOAc, washed with NaHCO₃ (sat.),
and dried over K₂CO₃. Following evaporative removal of the solvent,

- 157 -

the residue was chromatographed (silica gel, 17% ethyl acetate/hexanes) to give 13-5 as a colorless oil and 13-4 as a colorless oil.

TLC Rf = 0.44 (25% ethyl acetate/hexanes).

¹H NMR (300 MHz, CHCl₃) δ 7.32 (m, 5H), 4.97 (d, 1H, J=15 Hz), 4.17 (t, 1H, J=7.5 Hz), 4.05 (d, 1H, J=15 Hz), 3.44 (m, 1H), 2.48 (m, 2H), 1.50 (m, 1H), 1.22 (d, 3H, J=6.6 Hz).

(1-benzyl-5(R)-methyl-2-oxo-pyrrolidin-3(S)-yl)-carbamic acid tert-butyl ester (13-6)

10 A mixture of 13-5 (2.38 g, 10.3 mmol), 10% Pd/C (1.0 g), TFA (10 mL), THF (80 mL) and methanol (100 mL) was stirred at ambient temperature under a hydrogen atmosphere (1 atm) for 3 hr. The catalyst was removed by filtration through a celite pad and the filtrate concentrated to give the intermediate amine salt as a colorless
15 oil. To a solution of the crude amine salt in THF (50 mL) at 0°C was added NEt₃ (2.88 mL, 20.7 mmol) and di-tert-butyl dicarbonate (2.59 g, 11.9 mmol). The mixture was allowed to warm to ambient temperature and stir for 4 hours. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 40% ethyl acetate/hexanes) to
20 give 13-6 as a colorless oil.

TLC RF = 0.44 (silica, 40% ethyl acetate/hexanes);

¹H NMR (300 MHz, CHCl₃) δ 7.31 (m, 5H), 5.17 (br s, 1H), 4.94 (d, 1H, J=15 Hz), 4.20 (m, 1H), 4.07 (d, 1H, J=15 Hz), 3.44 (m, 1H), 2.77 (m, 1H), 1.45 (s, 9H), 1.20 (d, 3H, J=7 Hz).

25 (5(R)-methyl-2-oxo-pyrrolidin-3(S)-yl)-carbamic acid tert-butyl ester (13-7)

To a blue solution of lithium metal (0.237 g, 34.2 mmol) in NH₃ (l) (200 mL) at -78°C was added a solution of 13-6 (2.60 g, 8.54
30 mmol) in THF (15 mL). The mixture was stirred for 15 minutes, then quenched by the addition of ammonium chloride until the blue color dispersed. An additional 30 mL of THF was added, and the mixture warmed to 35°C to evaporate the ammonia. MgSO₄ was added, the mixture was filtered through a celite pad. Following evaporative

- 158 -

removal of the solvent, the residue was chromatographed (silica gel, 70:20:10 chloroform/ethyl acetate/MeOH) to give 13-7 as a colorless oil
TLC R_f = 0.45 (70:20:10 chloroform/ethyl acetate/MeOH).

¹H NMR (300 MHz, CHCl₃) δ 6.97 (br s, 1H), 5.24 (d, 1H, J=7.6 Hz),
5 4.32 (br s, 1H), 3.66 (m, 1H), 2.79 (m, 1H), 1.45 (s, 9H), 1.25 (d, 3H,
J=6.0 Hz).

(3(S)-tert-butoxycarbonylamino-5(R)-methyl-2-oxo-pyrrolidin-1-yl)-
acetic acid ethyl ester (13-8)

10 To a solution of 13-7 (1.83 g, 8.4 mmol) in THF (22 mL)
at -78° C was added sodium bis(trimethylsilyl)amide (9.4 mL, 9.4
mmol; 1M/ THF) dropwise. After an additional 20 min, ethyl
bromoacetate (1.13 mL, 10.3 mmol) was added dropwise. After an
additional 20 minutes, the mixture was allowed to warm to 0°C, and 20
15 mL sat. aqueous NH₄Cl was added. The layers were separated, the
aqueous layer washed with EtOAc, and the combined organic extracts
were dried over K₂CO₃. Following evaporative removal of the solvent,
the residue was chromatographed (silica gel, 40% ethyl acetate/hexanes)
to give 13-8 as a colorless oil.

20 TLC R_f = 0.39 (40% ethyl acetate/hexanes).
¹H NMR (300 MHz, CHCl₃) δ 5.20 (br s, 1H), 4.38 (d, 1H, J=18 Hz),
4.21 (m, 3H), 3.77 (m, 2H), 2.83 (m, 1H), 1.44 (s, 9H), 1.23 (m, 6H).

(3(S)-tert-butoxycarbonylamino-5(R)-methyl-2-oxo-pyrrolidin-1-yl)-
25 acetic acid (13-9)

To a solution 13-8 (527 mg, 1.75 mmol) in EtOH was
added 1N NaOH (1.93 mL, 1.925 mmol). After stirring for 1 h, the
solvents were evaporated, the mixture was diluted with EtOAc, acidified
with 10% KHSO₄, washed with brine, dried over MgSO₄, and
30 evaporated to give 13-9 as a white solid.

TLC R_f = 0.48 (silica, 9.5/0.5/0.5 CH₂Cl₂/MeOH/AcOH)

¹H NMR (300 MHz, CD₃OD) δ 4.21 (m, 2H), 3.85 (d, 1H, J=18 Hz),
3.74 (m, 1H), 2.58 (m, 1H), 1.52 (m, 1H), 1.44 (s, 9H), 1.25 (d, J=6.3
Hz, 3H).

- 159 -

(3(S)-tert-butoxycarbonylamino-5(R)-methyl-2-oxo-pyrrolidin-1-yl)-acetyl-3(S)-alkynyl-β-alanine ethyl ester (13-10)

5 A mixture of 13-9 (440 mg, 1.62 mmol), 2-9 (290 mg, 1.62 mmol), EDC (373 mg, 1.94 mmol), HOBT (262 mg, 1.94 mmol) and NMM (1.20 mL, 11.34 mmol) in CH₃CN (5 mL) was stirred for 20 h. The mixture was diluted with EtOAc, washed with sat. NaHCO₃, brine, and dried over MgSO₄. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, EtOAc) to give
10 13-10 as a colorless foam.

TLC R_f = 0.20 (silica, EtOAc).

¹H NMR (300 MHz, CDCl₃) δ 7.31 (bd, 1H), 5.33 (bd, 1H), 5.21 (m, 1H), 4.16 (m, 5H), 3.64 (m, 2H), 2.72 (m, 2H), 2.45 (d, J=2.2 Hz, 1H), 1.52 (m, 1H), 1.46 (s, 9H), 1.27 (m, 6H).

15 (3(S)-amino-5(R)-methyl-2-oxo-pyrrolidin-1-yl)-acetyl-3(S)-alkynyl-β-alanine ethyl ester hydrochloride (13-11)

To a solution of 13-10 (550 mg, 1.39 mmol) in EtOAc at 0°C was bubbled HCl gas for 5 minutes. The reaction was stirred an
20 additional 5 minutes, followed by removal of the cooling bath and then purged with Argon for 20 minutes. Evaporative removal of the solvent gave 13-11 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 5.02 (m, 1H), 4.12 (m, 4H), 3.83 (m, 2H), 2.77 (m, 2H), 1.59 (m, 1H), 1.25 (m, 6H).

25 5(R)-methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester (13-12)

To a solution of 13-11 (450 mg, 1.39 mmol) and 5,6,7,8-tetrahydro-[1,8]naphthyridine-2-carbaldehyde (225 mg, 1.39 mmol) in dichloroethane at 0°C was added Na(OAc)₃BH. After 1.5 h the reaction
30 was quenched with sat. NaHCO₃, diluted with EtOAc, washed with sat. NaHCO₃, brine and dried over MgSO₄. Following evaporative removal

- 160 -

of the solvent, the residue was chromatographed (silica gel, 70:20:10 chloroform/ethyl acetate/MeOH) to give 13-12 as a colorless foam.

TLC R_f = 0.17 (70:15:15 chloroform/ethyl acetate/MeOH).

¹H NMR (300 MHz, CDCl₃) δ 7.09 (d, J=6.8 Hz, 1H), 7.03 (d, J=8.5 Hz, 1H), 6.48 (d, J=7.3 Hz, 1H), 5.05 (m, 1H), 4.83 (bs, 1H), 4.17 (q, J=6.4, 1H), 3.92 (m, 2H), 3.74 (m, 2H), 3.58 (m, 4H), 3.40 (m, 2H), 2.70 (m, 4H), 2.54 (m, 1H), 2.26 (s, 1H), 1.90 (m, 2H), 1.55 (m, 2H), 1.25 (m, 6H).


- 10 5(R)-methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine (13-13)

To a solution of 13-12 (108 mg, 0.24 mmol) in EtOH (2 mL) was added 1N NaOH (0.270 ml, 0.264 mmol). After stirring for 1 h, the solvents were evaporated and the residue was chromatographed (silica gel, 25:10:1:1 to 15:10:1:1 ethyl acetate/EtOH/water/NH₄OH) to give 13-13 as a colorless foam.

TLC R_f = 0.23 (12:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

¹H NMR (300 MHz, D₂O) δ 7.53 (d, 1H, J=7.3 Hz), 6.69 (d, 1H, J=7.3 Hz), 4.41 (m, 2H), 3.71 (m, 2H), 3.45 (t, J=5.8 Hz, 2H), 2.79 (t, J=5.8 Hz, 2H), 2.6 (m, 4H), 1.92 (m, 3H), 1.50 (m, 1H), 1.19 (m, 3H).

SCHEME 14



- 161 -

2-Dimethoxymethyl-[1.8]naphthyridine (14-1)

A mixture containing 1-4 (30 g, 0.245 mol), pyruvaldehyde dimethylacetal (87 g, 0.737 mol), and L-proline (7.0g, 0.062 mol) in MeOH (300 mL) was refluxed under argon for 16 h.

5 The cooled solution was filtered, evaporated and the residue dissolved in CH₂Cl₂ (500 mL) and washed with water and brine then dried and concentrated to a volume of ca. 100 mL. Hexane (300 mL) was added and the mixture was kept at 0°C for 3 h, then filtered affording 14-1 as an off-white crystalline solid.

10 ¹H NMR (300 MHz, CDCl₃) δ, 9.14 (d, J = 2.2 Hz, 1H); 8.26 (d, J = 8.7 Hz, 1H); 8.21 (dd, J = 8.7, 2.2 Hz, 1H); 7.8 (d, J = 8.3 Hz, 1H); 7.5 (m, 1H); 5.48 (s, 1H); 3.53 (s, 6H).

2-Dimethoxymethyl-5,6,7,8-tetrahydro-[1.8]naphthyridine (14-2)

15 A solution 14-1 (10 g, 0.049 mol) in MeOH, (100 ml) was treated with 10% Pd on C (1.5 g) and the resulting mixture stirred under a H₂ filled balloon for 12.5 h. The catalyst was removed by filtration through celite and the solution concentrated to afford 14-2 as a yellow crystalline solid.

20 ¹H NMR (300 MHz, CDCl₃) δ 7.18 (d, J = 7.12 Hz, 1H); 6.71 (d, J = 7.12 Hz, 1H); 5.18 (s, 1H); 4.96 (br, s, 1H); 3.43 (s, 6H); 3.4 (m, 2H); 2.65 (m, 2H); 1.91 (m, 2H).


5,6,7,8-tetrahydro-[1.8]naphthyridine-2-carboxaldehyde (14-3)

25 14-2 (10 g, 0.048 mol) was trifluoroacetic acid (50 mL) and the resulting solution stirred under argon for 12.5 h. The TFA was removed at reduced pressure and the residue partitioned between sat. NaHCO₃ and CH₂Cl₂. The organic layer was dried, concentrated and passed through a 3 in. pad of silica gel (10% acetone/CH₂Cl₂) and concentrated to afford 14-3 as a yellow crystalline solid.

30 ¹H NMR (300 MHz, CDCl₃) δ 9.80 (s, 1H); 7.31 (d, J = 7.32 Hz, 1H); 7.16 (d, J = 7.32 Hz, 1H); 5.31 (br, s, 1H); 3.48 (m, 2H); 2.81 (m, 2H); 1.94 (m, 2H).


- 162 -

SCHEME 15



- 163 -

SCHEME 15 (CONT'D)



- 164 -

1-Bromo-3-(2,2-diethoxy-ethoxy)-benzene (15-2)

To a suspension of NaH (2.77 g, 115.6 mmol) in DMF (100 mL) at 0 °C was added a solution of 3-bromophenol 15-1 in DMF (40 mL) over 40 min. After the addition was complete, the solution was stirred for an additional 30 min. The solution was then treated with neat bromoacetaldehyde diethyl acetal (17.36 g, 115.6 mmol). The solution was heated at 100 °C for 8 h, cooled to room temperature, and extracted with Et₂O (3 x 200 mL). The combined organic extracts were washed with 10% aq NaOH (100 mL) and brine (100 mL), dried over MgSO₄, filtered and concentrated to give 15-2 as a yellow oil. TLC R_f = 0.4 (10% ethyl acetate/hexanes). ¹H NMR (300 MHz, CHCl₃) δ 7.19-7.05 (m, 3H), 6.85 (d, 1H), 4.81 (t, 1H, J=6.8 Hz), 3.99 (d, 2H, J=6.8 Hz), 3.71 (m, 4H), 1.22 (t, 6H, J=7.1 Hz) ppm.

6-Bromo-benzofuran (15-3)

To a solution of the acetal 15-2 in toluene (200 mL) was added polyphosphoric acid (20 g). The biphasic mixture was heated to 100 °C and stirred at this temperature for 4 h. The mixture was cooled to room temperature, poured onto ice, and extracted with Et₂O (2 x 200 mL). The combined organic extracts were washed with saturated aq NaHCO₃ and brine. The solution was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (100% hexanes) to give the product 15-3 as a yellow oil. TLC R_f = 0.3 (100% hexanes). ¹H NMR (300 MHz, CHCl₃) δ 7.68 (s, 1H), 7.60 (d, 1H, J=2.1 Hz), 7.46 (d, 1H, J=8.4 Hz), 7.36 (dd, 1H, J=8.1, 1.5 Hz), 6.75 (dd, 1H, J=7.1, 0.9 Hz) ppm.

3-Benzofuran-6-yl-acrylic acid ethyl ester (15-4)

A mixture of the 6-bromobenzofuran 15-3 (1.74 g, 8.79 mmol), ethyl acrylate (1.09 g, 10.98 mmol), Pd(OAc)₂ (0.099 g, 0.44 mmol), tri-*o*-tolylphosphine (0.268 g, 0.880 mmol), and sodium acetate (3.60 g, 43.9 mmol) in DMF (10 mL) was heated to 100 °C in a sealed

- 165 -

tube for 4 h. The mixture was cooled to room temperature, diluted with water, and extracted with Et₂O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL), dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (10% ethyl acetate/hexanes) to give the ester 15-4 as an off-white solid.

TLC R_f = 0.3 (10% ethyl acetate/hexanes).

¹H NMR (300 MHz, CHCl₃) δ 7.78 (d, 1H, J=15.9 Hz), 7.68 (d, 1H, J=2.4 Hz), 7.66 (s, 1H), 7.59 (d, 1H, J=8.4 Hz), 7.43 (dd, 1H, J=9.0, 1.5 Hz), 6.78 (m, 1H), 6.47 (d, 1H, J=15.9 Hz), 4.27 (q, 2H, J=7.2 Hz), 1.34 (t, 3H, J=7.2 Hz) ppm.

3-(S)-Benzofuran-6-yl-3-[benzyl-(1(R)-phenyl-ethyl)-amino]-propionic acid ethyl ester (15-5)

15 A solution of benzyl-α-(R)-methylbenzylamine (1.32 g, 6.30 mmol) in THF (25 mL) at 0 °C was treated with n-BuLi (2.52 mL of a 2.5 M soln in hexanes). The resulting solution was stirred at 0 °C for 30 min and then cooled to -78 °C. A solution of acrylate 15-4 (0.681 g, 3.15 mmol) in THF (5 mL) was added. After stirring for 15 min at -78 °C, satd aq NH₄Cl soln (5 mL) was added and the cold bath removed. The mixture was warmed to room temperature, and extracted with Et₂O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL), dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (10% ethyl acetate/hexanes) to give the β-aminoester 15-5 as a yellow oil.

TLC R_f = 0.8 (10% ethanol/dichloromethane).

¹H NMR (300 MHz, CHCl₃) δ 7.58 (m, 3H), 7.41 (m, 2H), 7.22 (m, 9H), 7.59 (s, 1H), 4.58 (m, 1H), 4.05 (m, 1H), 3.91 (q, 2H, J=7.1 Hz), 3.72 (m, 2H), 2.62 (m, 2H), 1.21 (d, 3H, J=7.2 Hz), 1.03 (t, 3H, J=7.1 Hz) ppm.

3(S)-Amino-3-(2,3-dihydro-benzofuran-6-yl)-propionic acid ethyl ester (15-6)

- 166 -

A mixture of the dibenzylamine 15-5 (1.19 g, 2.78 mmol) in EtOH/H₂O/AcOH (26 mL/3 mL/1.0 mL) was degassed with argon and treated with Pd(OH)₂ (1.19 g). The mixture was placed under 1 atm of H₂. After stirring for 18 h, the mixture was diluted with EtOAc, and filtered through celite. The filtrate was concentrated and the residue purified by flash chromatography (10% ethyl acetate/dichloromethane) to give the ester 15-6 as a white solid. TLC R_f = 0.25 (10% ethanol/dichloromethane).

¹H NMR (300 MHz, CD₃OD) as the trifluoroacetate salt: δ 7.25 (d, 1H, J=8.1 Hz), 6.88 (m, 1H), 7.66 (s, 1H), 6.82 (s, 1H), 4.58 (m, 3H), 4.12 (m, 2H), 3.30 (m, 1H), 3.19 (m, 2H), 2.98 (m, 2H), 1.11 (t, 3H, J=7.2 Hz) ppm.

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-acetylamino)-propionic acid ethyl ester (15-7)

A solution of the amine 15-6 (0.100 g, 0.425 mmol), acid 3-11 (0.155 g, 0.511 mmol), EDC (0.098 g, 0.511 mmol), NMM (0.103 g, 1.02 mmol), and HOAT (0.069 g, 0.511 mmol) in DMF (6 mL) was stirred at room temperature for 48 h. The solution was diluted with satd aq NaHCO₃ (3 mL) and extracted with EtOAc (2 x 10 mL). The combined organic extracts were washed with brine (10 mL), dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (8% ethanol/dichloromethane) to give the ester 15-7 as a yellow oil.

TLC R_f = 0.3 (10% ethanol/dichloromethane).
¹H NMR (300 MHz, CHCl₃) δ 7.12 (m, 2H), 6.78 (m, 1H), 6.65 (s, 1H), 6.39 (m, 1H), 5.36 (m, 1H), 4.99 (br s, 1H), 4.55 (t, J=7.2 Hz, 2H), 4.11 (m, 2H), 3.91 (m, 2H), 3.39 (m, 2H), 3.19 (m, 2H), 2.79 (m, 2H), 2.70 (m, 2H), 2.51 (m, 1H), 2.28 (m, 2H), 1.85 (m, 3H), 1.18 (m, 3H) ppm.

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-acetylamino)-propionic acid (15-8)


- 167 -

A solution of the ester 15-7 (0.038 g, 0.073 mmol) in EtOH/H₂O (4.5 mL/0.5 mL) was treated with LiOH (0.009 g, 0.365 mmol) and the homogeneous solution stirred at room temperature for 4 h. The solution was concentrated to a solid residue which was dissolved in H₂O and purified by preparative HPLC (gradient conditions: 95:05 to 50:50 H₂O/MeCN with 0.1% TFA) to give the acid 15-8 as a white solid (as the *bis*-trifluoroacetate salt).




MS (LR, FAB) M +1 calcd 493, found 493.39.

¹H NMR (300 MHz, CHCl₃) δ 7.91 (m, 1H), 7.35 (m, 1H), 7.09 (m, 1H), 6.76 (m, 1H), 6.68 (s, 1H), 6.43 (m, 2H), 5.28 (m, 1H), 4.53 (m, 2H), 4.41 (m, 1H), 3.38 (m, 7H), 3.14 (m, 3H), 2.81 (m, 5H), 2.60 (m, 1H), 2.28 (m, 1H), 2.05 (m, 3H) ppm.


- 168 -

SCHEME 16


- 169 -

SCHEME 16 (cont.)

- 170 -

3-Benzoyloxycarbonylamino-6-methyl-2-pyridinone (16-2).

- 5 DPPA (35.6 ml, 165 mmol) was added to a stirred solution of 2-hydroxy-6-methylpyridine-3-carboxylic acid (16-1; Aldrich; 22.97 g, 165 mmol) and triethylamine (23.0 ml, 165 mmol) in dry dioxane (300 ml) and the resulting solution was heated to reflux. After 16 h more triethylamine (23.0 ml, 165 mmol) and benzyl alcohol (17.1 ml, 165 mmol) were added and the solution was refluxed for a further 24 h.
- 10 The reaction was concentrated *in vacuo* to remove most of the volatiles. The residue was partitioned between methylene chloride (500 ml) and brine (500 ml), acidified to pH 1 with 1 M HCl (165 ml). The aqueous layer was extracted methylene chloride (two times) and the combined
- 15 organic layers were washed with sodium hydrogen carbonate solution and brine, dried (Na₂SO₄) and evaporated *in vacuo* to a brown solid. This was recrystallized from methanol, to give the title compound 16-2 as a tan solid: ¹H NMR (300 MHz, CDCl₃) δ 2.29 (s, 3H, CH₃), 5.20 (s, 2H, PhCH₂), 6.06 (d, J=7.6 Hz, pyridinone-5-H), 7.32-7.43 (m, 5H, Ph), 7.67 (br s, 1H, CbzNH), 8.03 (br d, pyridinone-4-H).
- 20

2-[6-methyl-2-oxo-3-(benzyloxycarbonylamino)-2H-pyridin-1-yl]acetic acid t-butyl ester (16-3).

25


Sodium hydride (5.3 g, 0.22 mol) was added to a stirred slurry of 3-benzyloxycarbonylamino-6-methyl-2-pyridinone (16-2; 53.2 g, 0.20 mol) in THF at 0°C. t-Butylbromoacetate (45 ml, 0.27 mol) was added

- 171 -

to the resulting solution and a precipitate rapidly forms. The reaction was warmed to rt over 1h and after 2h the solvent was evaporated *in vacuo* and the residue was partitioned between 1:1 water/brine (200 ml) and 6:1 THF/methylene chloride (700 ml). The organic layer was dried (Na₂SO₄) and evaporated *in vacuo* to a solid which was triturated with hexane to give the title compound **16-3** as a crystalline solid:
¹H NMR (400 Mz, CDCl₃) δ 1.47 (s, 9H), 2.25 (s, 3H), 4.75 (s, 2 H), 5.19 (s, 2H), 6.09 (d, J=7.8 Hz), 7.30-7.40 (m, 5H), 7.75 (br s, 1H), 7.94 (br d, 1H).

10

2-[6-methyl-2-oxo-3-(benzyloxycarbonylamino)-2H-pyridin-1-yl]acetic acid (**16-4**).




15

HCl gas was bubbled through a stirred suspension of 2-[6-methyl-2-oxo-3-(benzyloxycarbonylamino)-2H-pyridin-1-yl]acetic acid t-butyl ester (**16-3**; 12.3 g, 33 mmol) in ethyl acetate (250 ml) at -15°C for 20 min. The resulting solution was allowed to warm to room temperature and was then stirred there for 3 h. After purging with argon, the bulk of the solvent was rotavapped off and ether added to the residue. The solid which precipitated was filtered off and washed with ether. The title compound **16-4** was thus obtained as a white fluffy powder: ¹H NMR (CD₃OD) δ 2.32 (s, 3 H), 4.86 (s, 2 H), 5.18 (s, 2 H), 6.24 (d, J = 7.9 Hz, 1 H), 7.31-7.41 (m, 6 H), 7.94 (br s, 1 H).

25


3-(2-(6-methyl-2-oxo-3-(benzyloxycarbonylamino)-2H-pyridin-1-yl)-acetylamin)-3(S)-pyridin-3-yl-propionic acid ethyl ester (**16-5**).

- 172 -



To a solution of the acid 2-4 (150 mg, 0.47 mmol) and the amine 1-9 (Rico *et al*; *J. Org. Chem.*, 1993, 58, 7948; 139 mg, 0.52 mmol) in DMF (3 mL) was added HOBT (77 mg, 0.57 mmol) then Et₃N (200 μ L, 1.42 mmol). After 15 minutes, EDC (136 mg, 0.71 mmol) was added and the mixture was stirred for 16 hours. The solution was poured into EtOAc, washed with saturated NaHCO₃ then brine, dried (MgSO₄), and evaporated to give the title compound 16-5 as a white solid which was used as such in the next step. ¹H NMR (CDCl₃) δ 1.14 (3H, t), 2.40 (3H, s), 2.8-2.9 (2H, m), 4.05 (2H, q), 4.78 (2H, m), 5.22 (2H, s), 5.4 (1H, q), 6.17 (1H, d), 7.22 (1H, m), 7.3-7.45 (4H, m), 7.59 (1H, m), 7.7-7.8 (2H, m), 8.0 (1H, m), 8.52 (2H, m).

3-(2-{6-methyl-2-oxo-3-amino-2H-pyridin-1-yl}-acetyl-amino)-3(S)-pyridin-3-yl-propionic acid ethyl ester (16-6).




To a degassed solution of the pyridone 16-5 (243 mg; 0.49 mmol) in EtOH (20 mL) was added 10% Pd on carbon (25 mg) and this was then stirred under an atmosphere of hydrogen gas (balloon) for 3 hours. The mixture was filtered through a pad of celite and the solvent removed to give the title compound (16-6) as a viscous oil which was used as such in the next step.

- 173 -

3-(2-{6-methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl}-acetylamino)-3(S)-pyridin-3-yl-propionic acid ethyl ester (16-7).

5



To a solution of the amine 16-6 (155 mg, 0.433 mmol), the aldehyde 14-3 (70 mg, 0.433 mmol) in $\text{CH}_2\text{ClCH}_2\text{Cl}$ was added crushed 4Å sieves, AcOH (20 μL) and then $\text{NaB}(\text{OAc})_3\text{H}$ (184 mg, 0.866 mmol).

- 10 After stirring for 48 hours, the mixture was filtered through celite, poured into EtOAc and washed with saturated NaHCO_3 then brine. The dried (MgSO_4) solution was concentrated in vacuo to give a foam type solid. Column chromatography (5% MeOH in CHCl_3) afforded the title compound 16-7 as a foam type solid.


- 15 Analysis calculated for $\text{C}_{27}\text{H}_{32}\text{N}_6\text{O}_4 \cdot 0.25\text{CHCl}_3$

C, 61.24; H, 6.08; N, 15.73

found C, 61.33; H, 6.09; N, 15.85.

FAB mass spectrum. $m/z = 505.34$ (M+H)

- 20 3-(2-{6-methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl}-acetylamino)-3(S)-pyridin-3-yl-propionic acid bis trifluoroacetate (16-8).



25

- 174 -


The ester 16-7 (120 mg, 0.238 mmol) was dissolved in H₂O (1 mL) and THF (1 mL) and then 1N LiOH (1 mL, 1 mmol) was added. After 2 hours, the mixture was purified by reverse phase HPLC (Waters PrepPak C18 column eluting with H₂O/acetonitrile gradient) to give, after lyophilization, the title compound 16-8 as a powder

5 Analysis calculated for C₂₅H₂₈N₆O₄•2.5TFA•0.55H₂O
C, 46.70; H, 4.13; N, 10.89
found C, 46.70; H, 4.14; N, 11.04.
FAB mass spectrum. m/z = 477.2 (M+H)

10


- 175 -

SCHEME 17




- 176 -


SCHEME 17 (cont.)



- 177 -

 β -N,N-Dimethylaminoethenylcyclopropyl ketone (17-2).


5 A mixture of cyclopropyl methyl ketone (5.88 ml, 59 mmol) and N,N-dimethylformaldehyde dimethyl acetal (7.83 ml, 59 mmol) was heated in the presence of a catalytic quantity of p-toluenesulfonic acid for 48 hours. The resulting crude sample of the title compound (17-2, a pale yellow oil) was used in subsequent
10 reactions without further purification: ^1H NMR (CDCl_3) δ 0.74 (m, 2 H), 1.00 (m, 2 H), 1.75 (m, 1 H), 3.48 (s, 3 H), 3.50 (s, 3 H), 5.20 (d, 1 H), 7.55 (d, 1 H).

6-Cyclopropyl-3-nitro-2-(1H)-pyridinone (17-3).

15 A mixture of crude β -N,N-dimethylaminoethenylcyclopropyl ketone (17-2; 12 g, < 86 mmol),
20 nitroacetamide (9 g, 86 mmol) and aqueous piperidinium acetate (10 ml) [prepared from glacial acetic acid (42 ml), water (100 ml) and piperidine (72 ml)] was stirred at room temperature overnight. Following dilution with water (20 ml), the yellow precipitate was isolated via filtration and drying *in vacuo* to yield the title compound
25 17-3: ^1H NMR (CDCl_3) δ 1.15 (m, 2 H), 1.36 (m, 2 H), 2.10 (m, 1 H), 6.02 (br d, $J = 8.0$ Hz, 1 H), 8.41 (d, $J = 8.0$ Hz, 1 H).

3-Nitro-6-cyclopropyl-1-(t-butyl-carboxymethylene)-2-pyridinone (17-4)


- 178 -



5 Solid 6-cyclopropyl-3-nitro-2-(1H)-pyridinone (17-3; 1.4 g, 7.78 mmol) was added in small portions to a suspension of sodium hydride (260 mg, 10.8 mmol) in THF (30 ml) at room temperature. After stirring the resulting solution for 20 min, *tert*-butylbromoacetate (4 ml, 27 mmol) was added. The mixture was stirred for an additional 30 min. then heated at 55°C for 15 hrs. After cooling to room
10 temperature the THF was evaporated *in vacuo* and ice carefully added to the residue to destroy any excess sodium hydride. The resulting mixture was extracted with 2:1:1 ethyl acetate:ether:chloroform and the extracts dried over magnesium sulfate. Filtration and evaporation of the filtrate gave a yellow oil as a 3:1 mixture of N and O-alkylated products
15 respectively. Flash column chromatography eluting with 1:1 hexane/ethyl acetate gave the title compound 17-4 as a yellow crystalline solid: ¹H NMR (CDCl₃) δ 0.94 (m, 2 H), 1.18 (m, 2 H), 1.49 (s, 9 H), 1.79 (m, 1 H), 5.04 (s, 2 H), 6.10 (d, J = 8.1 Hz, 1 H), 8.33 (d, J = 8.1 Hz, 1 H).

20

3-Amino-6-cyclopropyl-1-(t-butyl-carboxymethylene)-2-pyridinone
(17-5)




25

A mixture of 3-nitro-6-cyclopropyl-1-(t-butyl-methylene-carboxy)-2-pyridinone (17-4; 760 mg, 2.58 mmol) and platinum oxide (250 mg) in ethanol (30 ml) was stirred at 0° C under an atmosphere of

- 179 -


- hydrogen for 3 hours. Following removal of most of the catalyst by filtration through a bed of Celite, the filtrate was concentrated and the residue purified by flash column chromatography eluting with 2:1 hexane/ethyl acetate. This yielded the title product 17-5 as a viscous orange gum: ^1H NMR (CDCl_3) δ 0.67 (m, 2 H), 0.89 (m, 2 H), 1.49 (s, 9 H), 1.63 (m, 1 H), 4.07 (br s, 2 H), 4.99 (s, 2 H), 5.91 (dd, $J = 1.2$ and 7.4 Hz, 1 H), 6.47 (d, $J = 7.4$ Hz, 1 H).

- 10 [6-Cyclopropyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl] -amino]-2H-pyridin-1-yl]-acetic acid *tert*-butyl ester (17-6)



- 15 Following the procedure described for the synthesis of 16-7, the amine 17-5 was coupled with 14-3 to yield the title compound 17-6 as an oil. R_f (silica gel; 5%MeOH in CHCl_3) = 0.39

- 20 [6-Cyclopropyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl] -amino]-2H-pyridin-1-yl]-acetic acid (17-7).




- 25 Following the procedure described for the preparation of 16-8, the ester 17-6 was hydrolysed to give the title compound 17-7. ^1H NMR (CD_3OD) δ 0.66 (m, 2 H), 0.9 (m, 2 H), 1.78 (m, 1 H), 1.9 (m, 2H),

- 180 -

2.75 (m, 2H), 3.4 (m, 2H), 4.6 (br s, 2 H), 6.02 (d, 1H), 6.19 (d, 1H), 6.58 (m, 1 H), 7.27 (m, 1H).

5 Ethyl 3-amino-3(S)-(3-fluorophenyl)propionate hydrochloride (17-8).




10 The title compound was prepared starting from 3-fluorobenzaldehyde (Aldrich) by conversion to ethyl 3-fluorocinnamate and employing the asymmetric addition/hydrogenolysis methodology described by Rico *et al*; *J. Org. Chem.*, 1993, 58, 7948.

¹H NMR (CD₃OD) δ 1.21 (t, 3H), 3.03 (dd, 1H), 3.13 (dd, 1H), 4.15 (q, 2H), 4.77 (t, 1H), 7.19 (m, 1H), 7.3 (m, 2H), 7.5 (m, 1H).

15

3-(2-[6-Cyclopropyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl]-acetylamino)-3(S)-(3-fluorophenyl)-propionic acid ethyl ester (17-9).

20



Following the procedure described for the preparation of 16-5, the acid 17-7 was coupled with the amine 17-8 to yield the title compound 17-9.
25 Analysis calculated for C₃₀H₃₄N₅O₄ F1•0.25H₂O


C, 65.26; H, 6.30; N, 12.69

found C, 65.20; H, 6.04; N, 13.00.

- 181 -

FAB mass spectrum. $m/z = 548.12$ (M+H)

3-(2-{6-Cyclopropyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl}-acetylamino)-3(S)-(3-fluorophenyl)-
 5 propionic acid ditrifluoroacetate (17-10).



Following the procedure described for the preparation of 16-8, the ester
 10 17-9 was hydrolysed to give the title compound 17-10.

Analysis calculated for $C_{28}H_{30}N_5O_4 \cdot F_1 \cdot 2.15TFA \cdot 0.5H_2O$


C, 55.16; H, 4.91; N, 10.62

found C, 55.19; H, 4.91; N, 10.89.

FAB mass spectrum. $m/z = 520.05$ (M+H)

15

3-(2-{6-Cyclopropyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl}-acetylamino)-3(S)-(3-pyridyl)-
propionic acid ditrifluoroacetate (17-11).



20


Following the procedures described for Scheme 16, the acid 17-7 was
 coupled with the amine 1-9 followed by saponification of the ester to
 afford the title compound 17-11.

25 Analysis calculated for $C_{27}H_{30}N_6O_4 \cdot 2.5TFA \cdot 0.6H_2O$

- 182 -

C, 48.13; H, 4.25; N, 10.53
 found C, 48.11; H, 4.23; N, 10.64.
 FAB mass spectrum. $m/z = 503.25$ (M+H)

- 5 3-(2-{6-Cyclopropyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl}-acetylamino)-3(S)-(alkynyl)-propionic acid ethyl ester (17-12).
-




10

Following the procedures described for Scheme 16, the acid 17-7 was coupled with 3-amino-3(S)-(alkynyl)-propionic acid ethyl ester 2-9 (Zablockie *et al*, *J. Med. Chem.*, 1995, 38, 2378) to afford the title compound 17-12.

- 15 Analysis calculated for $C_{26}H_{31}N_5O_4 \cdot 0.35H_2O$

C, 64.53; H, 6.60; N, 14.47
 found C, 64.52; H, 6.71; N, 14.54.
 FAB mass spectrum. $m/z = 478.35$ (M+H)

- 20 3-(2-{6-Cyclopropyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyridin-1-yl}-acetylamino)-3(S)-(alkynyl)-propionic acid (17-13).
-



25


- 183 -

Following the procedure described for the preparation of 16-8, the ester 17-12 was hydrolysed to give the title compound 17-13.


FAB mass spectrum. $m/z = 450.23$ (M+H)

- 184 -


SCHEME 18




- 185 -

SCHEME 18 (cont.)

- 186 -

Benzyl-N-(1-cyanoethyl)glycine hydrochloride (18-2).

- 5 TMSCN (18.8 mL, 141 mmol) was added cautiously to a stirred solution of benzylglycine free base (23.3 g 141 mmol - from the HCl salt by partition between EtOAc and brine basified with saturated Na₂CO₃ solution) and acetaldehyde (7.88 mL, 141 mmol) in CH₂Cl₂ (50 mL). After 4 h the volatiles were removed *in vacuo* and the residue was taken
- 10 up in EtOAc and washed with brine, dried (Na₂SO₄) and evaporated *in vacuo* to an oil. The oil was redissolved in EtOAc and 9.9 M HCl in EtOH (15.25 mL, 151 mmol) was added to give a crystalline precipitate which was isolated by filtration, washing with EtOAc and Et₂O to give the title compound (18-2):
- 15 ¹H NMR (CD₃OD) δ 1.70 (d, 3H), 4.16 (d, 1H), 4.21 (d, 1H), 4.64 (q, 1H), 5.31 (s, 2H), 7.35-7.44 (m, 5H).

1-Benzyloxycarbonylmethyl-3,5-dichloro-6-methylpyrazinone (18-3).


20

- A stirred mixture of oxalyl chloride (40.4 mL, 463 mmol) and 18-2 (29.51 g, 116 mmol) in 1,2-dichlorobenzene (110 mL) was heated to 100°C for 15 h. The volatiles were removed *in vacuo* and the residue
- 25 was purified by chromatography (silica gel; hexanes followed by 30%EtOAc in hexanes) to give a solid which was heated in EtOAc/hexanes 2:5 (140 mL), cooled and collected by filtration to give the title compound 18-3 as a pale green solid:

- 187 -

¹H NMR (CDCl₃) δ 2.35 (s, 3H), 4.88 (s, 2H), 5.24 (s, 2H), 7.38 (m, 5H).


5 3-(4-Methoxybenzylamino)-5-chloro-6-methyl-1-benzyloxycarbonylmethyl-pyrazinone (18-4).



10 A solution of the pyrazinone 18-3 (5 g, 15.3 mmol) and 4 methoxybenzylamine (6.0 mL, 45.9 mmol) in EtOAc (60 mL) was heated at 80°C for 2 h. The solution was cooled and filtered. The filtrate was concentrated *in vacuo*, the residue swished with MeOH and filtered to afford the title compound as a solid:

15 ¹H NMR (CDCl₃) δ 2.23 (s, 3H), 3.82 (s, 3H), 4.5 (d, 2H), 4.81 (s, 2H), 5.22 (s, 2H), 6.25 (t, 1H), 6.85 (m, 2H), 7.27 (m, 2H), 7.38 (m, 5H).

3-(4-Methoxybenzylamino)-5-chloro-6-methyl-1-carboxymethyl pyrazinone (18-5).



20

A solution of the benzyl ester 18-4 (1.06 g, 2.48 mmol) in toluene (60 mL) was degassed with argon and then 150 mg 10% palladium on carbon was added. The mixture was stirred under an atmosphere of


- 188 -

hydrogen gas for 16 h. The solution was filtered through celite and the solvent evaporated to give the title compound 18-5 as a white solid:

¹H NMR (CD₃OD) δ 2.25 (s, 3H), 3.78 (s, 3H), 4.45 (s, 2H), 4.81 (s, 2H), 4.90 (s, 2H), 6.85 (d, 2H), 7.28 (d, 2H).

5

3-(4-Methoxybenzylamino)- 6-methyl-1-carboxymethyl pyrazinone (18-6).



10

The acid 18-5 (810 mg) was dissolved in 40 mL 1 N NaOH and 40 mL MeOH at room temperature and was treated with Raney nickel suspension (~5 g). A second charge of Raney nickel (~5 g) and 1 N NaOH (20 mL) was added after 3 h. After 6 h, the suspension was filtered through celite washing with water and MeOH. The volatiles were removed *in vacuo* and the residue then taken up 1 N HCl (~5 mL). Saturated NaHCO₃ solution was added until pH~7-8 and the solution was extracted exhaustively with EtOAc/THF. After drying (MgSO₄), the solvent was removed to give the title compound 18-6 as a solid which was used as such:


15

20

¹H NMR (CD₃OD) δ 2.16 (s, 3H), 3.76 (s, 3H), 4.46 (s, 2H), 4.64 (s, 2H), 4.86 (s, 2H), 6.65 (s, 1H), 6.85 (d, 2H), 7.25 (d, 2H).

3-Amino-6-methyl-1-carboxymethylpyrazinone (18-7).

25




- 189 -

The pyrazinone 18-6 (900 mg) was heated at reflux in trifluoroacetic acid (20 mL) for 7 h. The volatiles were removed *in vacuo* and the residue was azeotroped with CH₂Cl₂, then EtOAc then MeOH. MeOH was added to the residue and the solution filtered to remove impurities.

- 5 Removal of the methanol then afforded the title compound 18-7 which was used as such:


¹H NMR (CD₃OD) δ 2.22 (s, 3H), 4.82 (s, 2H), 6.58 (s, 1H).

- 10 3-(3-Fluorophenyl)-3-(2-{6-methyl-2-oxo-3-amino}-2H-pyrazin-1-yl)-acetylaminopropionic acid ethyl ester (18-8).



- 15 Following the procedure described for the preparation of 16-6, the acid 18-7 was coupled with the amine 17-8 to yield the title compound 18-8.
¹H NMR (CDCl₃) δ 1.15 (t, 3H), 2.23 (s, 3H), 2.78 (dd, 1H), 2.84 (dd, 1H), 4.05 (q, 2H), 4.68 (ABq, 2H), 5.30 (br s, 2H), 5.35 (m, 1H), 6.68 (s, 1H), 6.9-7.1 (m, 3H), 7.27 (m, 1H), 7.57 (d, 2H).

- 20 3-(3-Fluorophenyl)-3-(2-{6-methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]-2H-pyrazin-1-yl}-acetylaminopropionic acid ditrifluoroacetate (18-9).



- 190 -

Following the procedure described for the synthesis of 16-9, the amine 18-8 was coupled with 14-3 and the product hydrolyzed to yield the title compound 18-9.


Analysis calculated for $C_{25}H_{27}N_6O_4F \cdot 2.25TFA \cdot 0.85H_2O$

5 C, 46.23; H, 4.07; N, 10.97






found C, 46.22; H, 4.00; N, 11.12.

FAB mass spectrum. $m/z = 495.26$ (M+H)




- 191 -

SCHEME 19


- 192 -

SCHEME 19 (Cont'd)

- 193 -

SCHEME 19 (Cont'd)

NaOH



- 194 -

The test procedures employed to measure $\alpha v \beta 3$ binding and the bone resorption inhibiting activity of the compounds of the present invention are described below.

5 BONE RESORPTION-PIT ASSAY

When osteoclasts engage in bone resorption, they will literally cause the formation of pits in the surface of bone that they are acting upon. Therefore, when testing compounds for their ability to inhibit osteoclasts, it is useful to measure the ability of osteoclasts to excavate these resorption pits when the inhibiting compound is present.

Consecutive 200 micron thick cross sections from a six mm cylinder of bovine femur diaphysis were cut with a low speed diamond saw (Isomet, Beuler, Ltd., Lake Bluff, IL). Bone slices were pooled, placed in a 10% ethanol solution and refrigerated until further use.

Prior to experimentation, bone slices were ultrasonicated twice, 20 minutes each in H₂O. Cleaned slices were placed in 96 well plates such that two control lanes and one lane for each drug dosage are available. Each lane represents either triplicate or quadruplicate cultures. The bone slices in 96 well plates were sterilized by UV irradiation. Prior to incubation with osteoclasts, the bone slices were hydrated by the addition of 0.1 ml Medium 199, pH 6.9 containing 15% fetal bovine serum and 1% penicillin/streptomycin.

Osteoclasts were isolated from the long bones of 1 to 3 day old rat pups (Sprague-Dawley) by modifications of Chambers et al., (J. Cell. Science, 66:383-399). The resulting suspension (0.75 ml/bone) was gently triturated 90-120 times using a wide bore transfer pipet. The cellular population was separated from bone fragments by a cell strainer with a 100 micron nylon mesh. 100 μ l of the cell suspension was placed onto each bone slice. Test compounds were then added at the desired experimental concentrations.

Bone slices exposed to osteoclasts for 20-24 hrs were processed for staining. Tissue culture media was removed from each bone slice. Each well was washed with 200 μ l of H₂O, and the bone slices were then fixed for 20 minutes in 2.5% glutaraldehyde, 0.1 M

- 195 -

5 cacodylate, pH 7.4. After fixation, any remaining cellular debris was removed by 2 min. ultrasonication in the presence of 0.25 M NH₄OH followed by 2 X 15 min ultrasonication in H₂O. The bone slices were immediately stained for 6-8 min with filtered 1% toluidine blue and 1% borax.

10 After the bone slices have dried, resorption pits were counted in test and control slices. Resorption pits were viewed in a Microphot Fx (Nikon) fluorescence microscope using a polarizing Nikon IGS filter cube. Test dosage results were compared with controls and resulting IC₅₀ values were determined for each compound tested.

15 The appropriateness of extrapolating data from this assay to utility and use in mammalian (including human) disease states is supported by the teaching found in Sato, M., *et al.*, Journal of Bone and Mineral Research, Vol. 5, No. 1, 1990. That article teaches that certain bisphosphonates have been used clinically and appear to be effective in the treatment of Paget's disease, hypercalcemia of malignancy, osteolytic lesions produced by bone metastases, and bone loss due to immobilization or sex hormone deficiency. These same bisphosphonates are then tested in the resorption pit assay described above to confirm a correlation between their known utility and positive performance in the assay.

EIB ASSAY

25 Duong *et al.*, J. Bone Miner. Res., 8:S 378, describe a system for expressing the human integrin $\alpha\beta$ 3. It has been suggested that the integrin stimulates attachment of osteoclasts to bone matrix, since antibodies against the integrin, or RGD-containing molecules, such as echistatin (European Publication 382 451), can effectively block bone resorption.

30

Reaction Mixture:

1. 175 μ l TBS buffer (50 mM Tris-HCl pH 7.2, 150 mM NaCl, 1% BSA, 1 mM CaCl₂, 1 mM MgCl₂).

- 196 -

2. 25 μ l cell extract (dilute with 100 mM octylglucoside buffer to give 2000 cpm/25 μ l).
3. 125 I-echistatin (25 μ l/50,000 cpm) (see EP 382 451).
- 5 4. 25 μ l buffer (total binding) or unlabeled echistatin (non-specific binding).

10 The reaction mixture was then incubated for 1 h at room temp. The unbound and the bound $\alpha_v\beta_3$ were separated by filtration using a Skatron Cell Harvester. The filters (prewet in 1.5% polyethyleneimine for 10 mins) were then washed with the wash buffer (50 mM Tris HCl, 1mM $\text{CaCl}_2/\text{MgCl}_2$, pH 7.2). The filter was then counted in a gamma counter.

15 SPA ASSAY

MATERIALS:

1. Wheatgerm agglutinin Scintillation Proximity Beads (SPA): Amersham
- 20 2. Octylglucopyranoside: Calbiochem
3. HEPES: Calbiochem
4. NaCl: Fisher
5. CaCl_2 : Fisher
6. MgCl_2 : SIGMA
- 25 7. Phenylmethylsulfonylfluoride (PMSF): SIGMA
8. Optiplate: PACKARD
9. Compound 8-10 (specific activity 500-1000 Ci/mmole)
10. test compound
- 30 11. Purified integrin receptor: $\alpha_v\beta_3$ was purified from 293 cells overexpressing $\alpha_v\beta_3$ (Duong *et al.*, *J. Bone Min. Res.*, 8:S378, 1993) according to Pytela (*Methods in Enzymology*, 144:475, 1987)
12. Binding buffer: 50 mM HEPES, pH 7.8, 100 mM NaCl, 1 mM $\text{Ca}^{2+}/\text{Mg}^{2+}$, 0.5 mM PMSF

- 197 -

13. 50 mM octylglucoside in binding buffer: 50-OG buffer

PROCEDURE:

1. Pretreatment of SPA beads:
5 500 mg of lyophilized SPA beads were first washed four times with 200 ml of 50-OG buffer and once with 100 ml of binding buffer, and then resuspended in 12.5 ml of binding buffer.
2. Preparation of SPA beads and receptor mixture
10 In each assay tube, 2.5 μ l (40 mg/ml) of pretreated beads were suspended in 97.5 μ l of binding buffer and 20 μ l of 50-OG buffer. 5 μ l (~30 ng/ μ l) of purified receptor was added to the beads in suspension with stirring at room temperature for 30 minutes. The mixture was then centrifuged at 2,500 rpm in a
15 Beckman GPR Benchtop centrifuge for 10 minutes at 4°C. The pellets were then resuspended in 50 μ l of binding buffer and 25 μ l of 50-OG buffer.
3. Reaction
20 The following were sequentially added into Optiplate in corresponding wells:
 - (i) Receptor/beads mixture (75 μ l)
 - (ii) 25 μ l of each of the following: compound to be tested, binding buffer for total binding or 8-8 for non-specific
25 binding (final concentration 1 μ M)
 - (iii) 8-10 in binding buffer (25 μ l, final concentration 40 pM)
 - (iv) Binding buffer (125 μ l)
 - (v) Each plate was sealed with plate sealer from PACKARD and incubated overnight with rocking at 4°C
30
4. Plates were counted using PACKARD TOPCOUNT
5. % inhibition was calculated as follows:
A = total counts

- 198 -

B = nonspecific counts

C = sample counts

$$\% \text{ inhibition} = [(A-B)-(C-B)]/(A-B)/(A-B) \times 100$$

5 OCFORM ASSAY

Osteoblast-like cells (1.8 cells), originally derived from mouse calvaria, were plated in CORNING 24 well tissue culture plates in α MEM medium containing ribo- and deoxyribonucleosides, 10% fetal bovine serum and penicillin-streptomycin. Cells were seeded at 40,000/well in the morning. In the afternoon, bone marrow cells were prepared from six week old male Balb/C mice as follows:

Mice were sacrificed, tibiae removed and placed in the above medium. The ends were cut off and the marrow was flushed out of the cavity into a tube with a 1 mL syringe with a 27.5 gauge needle. The marrow was suspended by pipetting up and down. The suspension was passed through $>100 \mu\text{m}$ nylon cell strainer. The resulting suspension was centrifuged at $350 \times g$ for seven minutes. The pellet was resuspended, and a sample was diluted in 2% acetic acid to lyse the red cells. The remaining cells were counted in a hemacytometer. The cells were pelleted and resuspended at 1×10^6 cells/mL. 50 μL was added to each well of 1.8 cells to yield 50,000 cells/well and 1,25-dihydroxy-vitamin D3(D3) was added to each well to a final concentration of 10 nM. The cultures were incubated at 37°C in a humidified, 5% CO_2 atmosphere. After 48 h, the medium was changed. 72 h after the addition of bone marrow, test compounds were added with fresh medium containing D3 to quadruplicate wells. Compounds were added again after 48 h with fresh medium containing D3. After an additional 48 h the medium was removed, cells were fixed with 10% formaldehyde in phosphate buffered saline for 10 minutes at room temperature, followed by a 1-2 minute treatment with ethanol:acetone (1:1) and air dried. The cells were then stained for tartrate resistant acid phosphatase as follows:

The cells were stained for 10-15 minutes at room temperature with 50 mM acetate buffer, pH 5.0 containing 30 mM

- 199 -

sodium tartrate, 0.3 mg/mL Fast Red Violet LB Salt and 0.1 mg/mL Naphthol AS -MX phosphate. After staining, the plates were washed extensively with deionized water and air dried. The number of multinucleated, positive staining cells were counted in each well.

5

α v β 5 ATTACHMENT ASSAY

Duong et al., J. Bone Miner. Res., 11:S 290, describe a system for expressing the human α v β 5.

10 Materials:

1. Media and solutions used in this assay are purchased from BRL/Gibco, except BSA and the chemicals are from Sigma.
2. Attachment medium: HBSS with 1 mg/ml heat-inactivated fatty acid free BSA and 2 mM CaCl_2 .
- 15 3. Glucosaminidase substrate solution: 3.75 mM p-nitrophenyl-N-acetyl-beta-D-glucosaminide, 0.1 M sodium citrate, 0.25% Triton, pH 5.0.
4. Glycine-EDTA developing solution: 50 mM glycine, 5 mM EDTA, pH 10.5.

20

Methods:

1. Plates (96 well, Nunc Maxi Sorp) were coated overnight at 4 °C with human vitronectin (3 ug/ml) in 50 mM carbonate buffer (pH 9.6), using 100 μ l/well. Plates were then washed 25 2X with DPBS and blocked with 2% BSA in DPBS for 2h at room temperature. After additional washes (2X) with DPBS, plates were used for cell attachment assay.
2. 293 (α v β 5) cells were grown in MEM media in presence of 10% fetal calf serum to 90% confluence. Cells were then lifted from dishes with 1X Trypsin/EDTA and washed 3X with serum free MEM. Cells were resuspended in 30 attachment medium (3×10^5 cells/ml).
3. Test compounds were prepared as a series of dilutions at 2X concentrations and added as 50 μ l/well. Cell suspension was

- 200 -

then added as 50 ml/well. Plates were incubated at 37 °C with 55 CO₂ for 1 hour to allow attachment.

- 5 4. Non-adherent cells were removed by gently washing the plates (3X) with DPBS and then incubated with glucosaminidase substrate solution (100 µl/well), overnight at room temperature in the dark. To quantitate cell numbers, standard curve of glucosaminidase activity was determined for each experiment by adding samples of cell suspension directly to wells containing the enzyme substrate solution.
- 10 5. The next day, the reaction was developed by addition of 185 µl/well of glycine/EDTA solution and reading absorbance at 405 nm using a Molecular Devices V-Max plate reader. Average test absorbance values (4 wells per test samples) were calculated. Then, the number of attached cells at each drug concentration was quantitated versus the standard curve of cells using the Softmax program.
- 15

EXAMPLE OF A PHARMACEUTICAL FORMULATION

20 As a specific embodiment of an oral composition, 100 mg of compound 3-13 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

25 Representative compounds of the present invention were tested and found to bind to human $\alpha\beta 3$ integrin. These compounds were found to have IC₅₀ values in the range of 0.1 to 100 nM in the SPA assay.

30 Representative compounds of the present invention were tested and found to inhibit $\geq 50\%$ the attachment of $\alpha\beta 5$ expressing cells to plates coated with vitronectin at concentrations of 1 µM.

While the invention has been described and illustrated in reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the


- 201 -

invention. For example, effective dosages other than the preferred doses as set forth hereinabove may be applicable as a consequence of variations in the responsiveness of the mammal being treated for severity of bone disorders caused by resorption, or for other indications
5 for the compounds of the invention indicated above. Likewise, the specific pharmacological responses observed may vary according to and depending upon the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected
10 variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended, therefore, that the invention be limited only by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.


- 202 -

WHAT IS CLAIMED IS:

1. A method of eliciting a vitronectin receptor antagonizing effect in a mammal in need thereof, comprising
 5 administering to the mammal a therapeutically effective amount of a compound of the formula




wherein X is selected from



- 10 a 5- or 6-membered monocyclic aromatic or nonaromatic ring system containing 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S wherein the 5- or 6-membered ring system is either unsubstituted or substituted with one or more groups chosen from R¹, R², R¹⁵ or R¹⁶;
 15 a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted with one or more groups chosen from R¹, R², R¹⁵ or R¹⁶;
 20

Y is selected from

- 203 -



Z is

a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; provided that Z is not a 6-membered monocyclic aromatic ring system, an isoxazoline ring or an isoxazole ring;

R¹, R², R⁴, R⁵, R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from

hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, C₃-8 cycloheteroalkyl, C₃-8 cycloalkyl C₁-6 alkyl, C₃-8 cycloheteroalkyl C₁-6 alkyl, aryl, aryl C₁-8 alkyl, amino, amino C₁-8 alkyl, C₁-3 acylamino, C₁-3 acylamino C₁-8 alkyl, (C₁-6 alkyl)_qamino, (C₁-6 alkyl)_qamino C₁-8 alkyl, C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl, hydroxycarbonyl C₁-6 alkyl, C₁-3 alkoxycarbonyl, C₁-3 alkoxycarbonyl C₁-6 alkyl, hydroxycarbonyl-C₁-6 alkyloxy, hydroxy, hydroxy C₁-6 alkyl, C₁-6 alkyloxy-C₁-6 alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy, trifluoroethoxy, C₁-8 alkyl-S(O)_q, (C₁-8 alkyl)_qaminocarbonyl,

- 204 -

C1-8 alkyloxycarbonylamino, (C1-8 alkyl)qaminocarbonyloxy, oxo, (aryl C1-8 alkyl)qamino, (aryl)qamino, aryl C1-8 alkylsulfonfylamino or C1-8 alkylsulfonylamino;

- 5 R³ is selected from
hydrogen,
aryl,
aryl-(CH₂)_p-,
hydroxyl,
10 C1-5 alkoxy,
aminocarbonyl,
C3-8 cycloalkyl,
amino C1-6 alkyl,
(aryl)qaminocarbonyl,
15 (aryl C1-5 alkyl)qaminocarbonyl,
hydroxycarbonyl C1-6 alkyl,
C1-8 alkyl,
aryl C1-6 alkyl,
(C1-6 alkyl)qamino C1-6 alkyl,
20 (aryl C1-6 alkyl)qamino C1-6 alkyl,
C1-8 alkylsulfonyl,
C1-8 alkoxycarbonyl,
aryloxycarbonyl,
aryl C1-8 alkoxycarbonyl,
25 C1-8 alkylcarbonyl,
arylcarbonyl,
aryl C1-6 alkylcarbonyl,
(C1-8 alkyl)qaminocarbonyl,
aminosulfonyl,
30 C1-8 alkylaminosulfonyl,
(aryl)qaminosulfonylamino,
(aryl C1-8 alkyl)qaminosulfonyl,
C1-6 alkylsulfonyl,
arylsulfonyl,

- 205 -

aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylthiocarbonyl,
arylthiocarbonyl, or
5 aryl C₁₋₆ alkylthiocarbonyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴;

R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from
10 hydrogen,
aryl,
aryl-(CH₂)_p-,
aryl-(CH₂)_n-O-(CH₂)_m-,
aryl-(CH₂)_n-S(O)_q-(CH₂)_m-,
15 aryl-(CH₂)_n-C(O)-(CH₂)_m-,
aryl-(CH₂)_n-C(O)-N(R³)-(CH₂)_m-,
aryl-(CH₂)_n-N(R³)-C(O)-(CH₂)_m-,
aryl-(CH₂)_n-N(R³)-(CH₂)_m-,
halogen,
20 hydroxyl,
C₁₋₈ alkylcarbonylamino,
aryl C₁₋₅ alkoxy,
C₁₋₅ alkoxycarbonyl,
(C₁₋₈ alkyl)_qaminocarbonyl,
25 C₁₋₆ alkylcarbonyloxy,
C₃₋₈ cycloalkyl,
oxo,
(C₁₋₆ alkyl)_qamino,
amino C₁₋₆ alkyl,
30 arylaminocarbonyl,
aryl C₁₋₅ alkylaminocarbonyl,
aminocarbonyl,
aminocarbonyl C₁₋₆ alkyl,
hydroxycarbonyl,

- 206 -

- hydroxycarbonyl C₁₋₆ alkyl,
 C₁₋₈ alkyl, either unsubstituted or substituted, with one or more
 groups selected from: halogen, hydroxyl,
 C₁₋₅ alkylcarbonylamino, aryl C₁₋₅ alkoxy,
 5 C₁₋₅ alkoxycarbonyl, aminocarbonyl, (C₁₋₅ alkyl)qamino-
 carbonyl, C₁₋₅ alkylcarbonyloxy, C₃₋₈ cycloalkyl, oxo,
 (C₁₋₃ alkyl)qamino, amino C₁₋₃ alkyl,
 (aryl)qaminocarbonyl, (aryl C₁₋₅ alkyl)qaminocarbonyl,
 aminocarbonyl, aminocarbonyl C₁₋₄ alkyl,
 10 hydroxycarbonyl or hydroxycarbonyl C₁₋₅ alkyl,
 CH≡C-(CH₂)_s-,
 C₁₋₆ alkyl-C≡C-(CH₂)_s-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,
 aryl-C≡C-(CH₂)_s-,
 15 C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,
 CH₂=CH-(CH₂)_s-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_s-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,
 aryl-CH=CH-(CH₂)_s-,
 20 C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,
 C₁₋₆ alkyl-SO₂-(CH₂)_s-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_s-,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 25 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)qamino C₁₋₆ alkyl,
 (aryl)qamino,
 (aryl)qamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)qamino,
 30 (aryl C₁₋₆ alkyl)qamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)qaminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,

- 207 -

- arylsulfonylamino,
C1-8 alkylsulfonylamino C1-6 alkyl,
arylsulfonylamino C1-6 alkyl,
aryl C1-6 alkylsulfonylamino,
5 aryl C1-6 alkylsulfonylamino C1-6 alkyl,
C1-8 alkoxycarbonylamino,
C1-8 alkoxycarbonylamino C1-8 alkyl,
aryloxycarbonylamino C1-8 alkyl,
aryl C1-8 alkoxycarbonylamino,
10 aryl C1-8 alkoxycarbonylamino C1-8 alkyl,
C1-8 alkylcarbonylamino,
C1-8 alkylcarbonylamino C1-6 alkyl,
arylcarbonylamino C1-6 alkyl,
aryl C1-6 alkylcarbonylamino,
15 aryl C1-6 alkylcarbonylamino C1-6 alkyl,
aminocarbonylamino C1-6 alkyl,
(C1-8 alkyl)qaminocarbonylamino,
(C1-8 alkyl)qaminocarbonylamino C1-6 alkyl,
(aryl)qaminocarbonylamino C1-6 alkyl,
20 (aryl C1-8 alkyl)qaminocarbonylamino,
(aryl C1-8 alkyl)qaminocarbonylamino C1-6 alkyl,
aminosulfonylamino C1-6 alkyl,
(C1-8 alkyl)qaminosulfonylamino,
(C1-8 alkyl)qaminosulfonylamino C1-6 alkyl,
25 (aryl)qaminosulfonylamino C1-6 alkyl,
(aryl C1-8 alkyl)qaminosulfonylamino,
(aryl C1-8 alkyl)qaminosulfonylamino C1-6 alkyl,
C1-6 alkylsulfonyl,
C1-6 alkylsulfonyl C1-6 alkyl,
30 arylsulfonyl C1-6 alkyl,
aryl C1-6 alkylsulfonyl,
aryl C1-6 alkylsulfonyl C1-6 alkyl,
C1-6 alkylcarbonyl,
C1-6 alkylcarbonyl C1-6 alkyl,

- 208 -

- 5 arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
10 (C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,
(aryl)_qaminocarbonyl C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_qaminocarbonyl, or
(aryl C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
15 are attached is itself attached to no more than one heteroatom; and
provided further that the carbon atom to which R¹⁰ and R¹¹ are
attached is itself attached to no more than one heteroatom;

- R¹² is selected from
20 hydrogen,
C₁₋₈ alkyl,
aryl,
aryl C₁₋₈ alkyl,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
25 aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
C₁₋₈ alkylaminocarbonylmethylene, or
C₁₋₈ dialkylaminocarbonylmethylene;

- m, s and t are each independently an integer from 0 to 3;
30 n is an integer from 1 to 3;
p is an integer from 1 to 4;
q is an integer from 0 to 2;
r is an integer from 0 to 6;

and the pharmaceutically acceptable salts thereof.


2. The method of Claim 1, wherein

X is



- 5 a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted with R^1 and R^2 ; and

10

Z is selected from




- 210 -




5 and the pharmaceutically acceptable salts thereof.

3. The method of Claim 2, wherein

Z is selected from




- 211 -




- 212 -

and the pharmaceutically acceptable salts thereof.

4. The method of Claim 3 wherein the compound has the formula



wherein X is selected from



Y is selected from $-(CH_2)_r-$ or $-(CH_2)_m-NR^3-(CH_2)_t-$;

R^3 is selected from

- 10 hydrogen,
 aryl- $(CH_2)_p-$,
 C₁₋₅ alkoxy carbonyl,
 C₃₋₈ cycloalkyl,
 (aryl)_qaminocarbonyl,
 15 (aryl C₁₋₅ alkyl)_qaminocarbonyl,
 C₁₋₈ alkyl,
 aryl C₁₋₆ alkyl,
 C₁₋₈ alkylsulfonyl,
 arylsulfonyl,
 20 aryl C₁₋₆ alkylsulfonyl,
 C₁₋₈ alkoxy carbonyl,
 aryloxy carbonyl,
 aryl C₁₋₈ alkoxy carbonyl,
 C₁₋₈ alkylcarbonyl,
 25 arylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl,
 (C₁₋₈ alkyl)_qaminocarbonyl,

- 213 -

C₁-6 alkylsulfonyl, or
aryl C₁-6 alkylcarbonyl,
wherein any of the alkyl groups may be unsubstituted or
substituted with R¹³ and R¹⁴;

5

R⁴ is selected from
hydrogen, C₁-6 alkyl, C₃-8 cycloalkyl, C₃-8 cycloheteroalkyl,
C₃-8 cycloalkyl C₁-6 alkyl, C₃-8 cycloheteroalkyl C₁-6 alkyl,
aryl or aryl C₁-8 alkyl,

10


R⁸ is selected from
hydrogen,
aryl,
aryl-(CH₂)_p-,
15 CH≡C-(CH₂)_s-,
C₁-6 alkyl-C≡C-(CH₂)_s-,
C₃-7 cycloalkyl-C≡C-(CH₂)_s-,
aryl-C≡C-(CH₂)_s-,
C₁-6 alkylaryl-C≡C-(CH₂)_s-,
20 CH₂=CH-(CH₂)_s-,
C₁-6 alkyl-CH=CH-(CH₂)_s-,
C₃-7 cycloalkyl-CH=CH-(CH₂)_s-,
aryl-CH=CH-(CH₂)_s-,
C₁-6 alkylaryl-CH=CH-(CH₂)_s-,
25 C₁-6 alkyl-SO₂-(CH₂)_s-,
C₁-6 alkylaryl-SO₂-(CH₂)_s-; and

r is an integer from 0 to 3;


30 and the pharmaceutically acceptable salts thereof.

5. The method of Claim 4 wherein the compound has
the formula


- 214 -



wherein Z is selected from



R^8 is selected from
hydrogen,



indolyl-(CH₂)_p-,

CH≡C-(CH₂)_s-,

C₁₋₆ alkyl-C≡C-(CH₂)_s-,

C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,

aryl-C≡C-(CH₂)_s-,

- 215 -

C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,
 CH₂=CH-(CH₂)_s-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_s-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,
 5 aryl-CH=CH-(CH₂)_s-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,
 C₁₋₆ alkyl-SO₂-(CH₂)_s-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_s-; and

10 R¹² is selected from hydrogen or C₁₋₈ alkyl;
 s is an integer from 0 to 3;
 and the pharmaceutically acceptable salts thereof.

6. The method of Claim 5, wherein the compound is
 15 selected from

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperidin-1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

20 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperidin-1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine trifluoroacetate;

25 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

30 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;

- 216 -

- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine ethyl ester;
- 5 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;
- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine ethyl ester;
- 10 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;
- Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- 15 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- 20 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- 25 Ethyl 2-oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;
- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;
- 30 Ethyl 3-(2-{2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}-pyrrolidin-1-yl)-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;

- 217 -

3-(2-{2-Oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-amino]pyrrolidin-1-yl}-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;

5 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-pyridin-3-yl-propionic acid;

10 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-pyridin-3-yl-propionic acid;

2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine ethyl ester;

15 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;

2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine ethyl ester;

20

2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

25 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine ethyl ester;

5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;

30

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-acetylamino)-propionic acid ethyl ester;

- 218 -

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-acetylamino)-propionic acid;

5 3-{2-(2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-acetylamino}-3(S)-quinolin-3-yl-propionic acid;

3-(2-(5(S)-Ethyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-acetylamino)-3-(S)-quinolin-3-yl-propionic
10 acid trifluoroacetate;

3-(2-{6-Methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}-2H-pyridin-1-yl)-acetylamino)-3(S)-pyridin-3-yl-propionic acid bis trifluoroacetate; or
15

3-(2-{6-Methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}-2H-pyridin-1-yl)-acetylamino)-3(S)-pyridin-3-yl-propionic acid ethyl ester;

20 and the pharmaceutically acceptable salts thereof.

7. The method of Claim 6, wherein the compound is selected from

25 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine trifluoroacetate;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;
30

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

- 219 -

- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;
- 5 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;
- 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3-(S)-pyridin-3-yl- β -alanine;
- 10 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3-(S)-pyridin-3-yl- β -alanine;
- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;
- 15 3-(2-{2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino]pyrrolidin-1-yl}-acetyl-amino)-3-(S)-pyridin-3-yl-propionic acid;
- 3-(2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl)-hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetyl-amino)-3-(S)-pyridin-3-yl-propionic acid;
- 20 3-(2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl)-hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetyl-amino)-3-(S)-pyridin-3-yl-propionic acid;
- 25 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;
- 30 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;
- 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)methyl]-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine; or

- 220 -

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-(2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-acetylamino)-propionic acid;

5

and the pharmaceutically acceptable salts thereof.

8. The method of Claim 1, wherein the vitronectin receptor antagonizing effect is an $\alpha v\beta 3$ antagonizing effect.

10

9. The method of Claim 8 wherein the $\alpha v\beta 3$ antagonizing effect is selected from inhibition of: bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth.

15

10. The method of Claim 9, wherein the $\alpha v\beta 3$ antagonizing effect is the inhibition of bone resorption.

11. The method of Claim 1, wherein the vitronectin receptor antagonizing effect is an $\alpha v\beta 5$ antagonizing effect.

20

12. The method of Claim 11 wherein the $\alpha v\beta 5$ antagonizing effect is selected from inhibition of: restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth.

25

13. The method of Claim 1, wherein the vitronectin receptor antagonizing effect is a dual $\alpha v\beta 3/\alpha v\beta 5$ antagonizing effect.

30

14. The method of Claim 13, wherein the dual $\alpha v\beta 3/\alpha v\beta 5$ antagonizing effect is selected from inhibition of: bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation or tumor growth.

- 221 -

15. A method of treating or preventing a condition mediated by antagonism of a vitronectin receptor in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the compound of Claim 1.


5

16. The method of Claim 15, wherein the condition is selected from the group consisting of osteoporosis or cancer.

17. A method of inhibiting bone resorption in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the compound of Claim 1.

18. A method of treating osteoporosis in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the compound of Claim 1.

19. A compound of the formula




wherein X is

20 a 9- to 10-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system contains 0, 1, 2, 3 or 4 heteroatoms selected from N, O or S, and wherein the polycyclic ring system is either unsubstituted or substituted with one or more groups chosen from R¹, R², R¹⁵ or R¹⁶;

25

Y is selected from

- 222 -



Z is

a 5-11 membered aromatic or nonaromatic mono- or polycyclic ring system containing 0 to 6 double bonds, and containing 0 to 6 heteroatoms chosen from N, O and S, and wherein the ring system is either unsubstituted or substituted with one or more groups independently selected from R⁴, R⁵, R⁶ and R⁷; provided that Z is not a 6-membered monocyclic aromatic ring system, an isoxazoline ring or an isoxazole ring;

R¹, R², R⁴, R⁵, R¹³, R¹⁴, R¹⁵ and R¹⁶ are each independently selected from

hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, C₃-8 cycloheteroalkyl, C₃-8 cycloalkyl C₁-6 alkyl, C₃-8 cycloheteroalkyl C₁-6 alkyl, aryl, aryl C₁-8 alkyl, amino, amino C₁-8 alkyl, C₁-3 acylamino, C₁-3 acylamino C₁-8 alkyl, (C₁-6 alkyl)_qamino, (C₁-6 alkyl)_qamino C₁-8 alkyl, C₁-4 alkoxy, C₁-4 alkoxy C₁-6 alkyl, hydroxycarbonyl, hydroxycarbonyl C₁-6 alkyl, C₁-3 alkoxy carbonyl, C₁-3 alkoxy carbonyl C₁-6 alkyl, hydroxycarbonyl C₁-6 alkyloxy, hydroxy, hydroxy C₁-6 alkyl, C₁-6 alkyloxy C₁-6 alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy, trifluoroethoxy, C₁-8 alkyl-S(O)_q, (C₁-8 alkyl)_qaminocarbonyl,

- 223 -

C1-8 alkyloxycarbonylamino, (C1-8 alkyl)qaminocarbonyloxy, oxo, (aryl C1-8 alkyl)qamino, (aryl)qamino, aryl C1-8 alkylsulfonylamino or C1-8 alkylsulfonylamino;

- 5 R³ is selected from
hydrogen,
aryl,
aryl-(CH₂)_p-,
hydroxyl,
10 C1-5 alkoxy,
aminocarbonyl,
C3-8 cycloalkyl,
amino C1-6 alkyl,
(aryl)qaminocarbonyl,
15 (aryl C1-5 alkyl)qaminocarbonyl,
hydroxycarbonyl C1-6 alkyl,
C1-8 alkyl,
aryl C1-6 alkyl,
(C1-6 alkyl)qamino C1-6 alkyl,
20 (aryl C1-6 alkyl)qamino C1-6 alkyl,
C1-8 alkylsulfonyl,
C1-8 alkoxycarbonyl,
aryloxycarbonyl,
aryl C1-8 alkoxycarbonyl,
25 C1-8 alkylcarbonyl,
arylcarbonyl,
aryl C1-6 alkylcarbonyl,
(C1-8 alkyl)qaminocarbonyl,
aminosulfonyl,
30 C1-8 alkylaminosulfonyl,
(aryl)qaminosulfonylamino,
(aryl C1-8 alkyl)qaminosulfonyl,
C1-6 alkylsulfonyl,
arylsulfonyl,

- 224 -

aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylthiocarbonyl,
arylthiocarbonyl, or
5 aryl C₁₋₆ alkylthiocarbonyl,
wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴;

R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are each independently selected from
10 hydrogen,
aryl,
aryl-(CH₂)_p-,
aryl-(CH₂)_n-O-(CH₂)_m-,
aryl-(CH₂)_n-S(O)_q-(CH₂)_m-,
15 aryl-(CH₂)_n-C(O)-(CH₂)_m-,
aryl-(CH₂)_n-C(O)-N(R³)-(CH₂)_m-,
aryl-(CH₂)_n-N(R³)-C(O)-(CH₂)_m-,
aryl-(CH₂)_n-N(R³)-(CH₂)_m-,
halogen,
20 hydroxyl,
C₁₋₈ alkylcarbonylamino,
aryl C₁₋₅ alkoxy,
C₁₋₅ alkoxycarbonyl,
(C₁₋₈ alkyl)_qaminocarbonyl,
25 C₁₋₆ alkylcarbonyloxy,
C₃₋₈ cycloalkyl,
oxo,
(C₁₋₆ alkyl)_qamino,
amino C₁₋₆ alkyl,
30 arylaminocarbonyl,
aryl C₁₋₅ alkylaminocarbonyl,
aminocarbonyl,
aminocarbonyl C₁₋₆ alkyl,
hydroxycarbonyl,

- 225 -

- hydroxycarbonyl C₁₋₆ alkyl,
 C₁₋₈ alkyl, either unsubstituted or substituted, with one or more
 groups selected from: halogen, hydroxyl,
 C₁₋₅ alkylcarbonylamino, aryl C₁₋₅ alkoxy,
 5 C₁₋₅ alkoxycarbonyl, aminocarbonyl, (C₁₋₅ alkyl)_qamino-
 carbonyl, C₁₋₅ alkylcarbonyloxy, C₃₋₈ cycloalkyl, oxo,
 (C₁₋₃ alkyl)_qamino, amino C₁₋₃ alkyl,
 (aryl)_qaminocarbonyl, (aryl C₁₋₅ alkyl)_qaminocarbonyl,
 aminocarbonyl, aminocarbonyl C₁₋₄ alkyl,
 10 hydroxycarbonyl or hydroxycarbonyl C₁₋₅ alkyl,
 CH≡C-(CH₂)_s-,
 C₁₋₆ alkyl-C≡C-(CH₂)_s-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,
 aryl-C≡C-(CH₂)_s-,
 15 C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,
 CH₂=CH-(CH₂)_s-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_s-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,
 aryl-CH=CH-(CH₂)_s-,
 20 C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,
 C₁₋₆ alkyl-SO₂-(CH₂)_s-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_s-,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 25 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_qamino C₁₋₆ alkyl,
 (aryl)_qamino,
 (aryl)_qamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_qamino,
 30 (aryl C₁₋₆ alkyl)_qamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_qaminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,

- 226 -

- arylsulfonylamino,
C₁-8 alkylsulfonylamino C₁-6 alkyl,
arylsulfonylamino C₁-6 alkyl,
aryl C₁-6 alkylsulfonylamino,
5 aryl C₁-6 alkylsulfonylamino C₁-6 alkyl,
C₁-8 alkoxycarbonylamino,
C₁-8 alkoxycarbonylamino C₁-8 alkyl,
aryloxycarbonylamino C₁-8 alkyl,
aryl C₁-8 alkoxycarbonylamino,
10 aryl C₁-8 alkoxycarbonylamino C₁-8 alkyl,
C₁-8 alkylcarbonylamino,
C₁-8 alkylcarbonylamino C₁-6 alkyl,
arylcarbonylamino C₁-6 alkyl,
aryl C₁-6 alkylcarbonylamino,
15 aryl C₁-6 alkylcarbonylamino C₁-6 alkyl,
aminocarbonylamino C₁-6 alkyl,
(C₁-8 alkyl)qaminocarbonylamino,
(C₁-8 alkyl)qaminocarbonylamino C₁-6 alkyl,
(aryl)qaminocarbonylamino C₁-6 alkyl,
20 (aryl C₁-8 alkyl)qaminocarbonylamino,
(aryl C₁-8 alkyl)qaminocarbonylamino C₁-6 alkyl,
aminosulfonylamino C₁-6 alkyl,
(C₁-8 alkyl)qaminosulfonylamino,
(C₁-8 alkyl)qaminosulfonylamino C₁-6 alkyl,
25 (aryl)qaminosulfonylamino C₁-6 alkyl,
(aryl C₁-8 alkyl)qaminosulfonylamino,
(aryl C₁-8 alkyl)qaminosulfonylamino C₁-6 alkyl,
C₁-6 alkylsulfonyl,
C₁-6 alkylsulfonyl C₁-6 alkyl,
30 arylsulfonyl C₁-6 alkyl,
aryl C₁-6 alkylsulfonyl,
aryl C₁-6 alkylsulfonyl C₁-6 alkyl,
C₁-6 alkylcarbonyl,
C₁-6 alkylcarbonyl C₁-6 alkyl,

- 227 -

arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
5 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,
10 (aryl)_qaminocarbonyl C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_qaminocarbonyl, or
(aryl C₁₋₈ alkyl)_qaminocarbonyl C₁₋₆ alkyl,


wherein any of the alkyl groups may be unsubstituted or substituted with
R¹³ and R¹⁴; and provided that the carbon atom to which R⁸ and R⁹
15 are attached is itself attached to no more than one heteroatom; and
provided further that the carbon atom to which R¹⁰ and R¹¹ are
attached is itself attached to no more than one heteroatom;

R¹² is selected from
20 hydrogen,
C₁₋₈ alkyl,
aryl,
aryl C₁₋₈ alkyl,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
25 aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
C₁₋₈ alkylaminocarbonylmethylene, or
C₁₋₈ dialkylaminocarbonylmethylene;


m, s and t are each independently an integer from 0 to 3;
n is an integer from 1 to 3;
30 p is an integer from 1 to 4;
q is an integer from 0 to 2;
r is an integer from 0 to 6;

and the pharmaceutically acceptable salts thereof.

20. The compound of Claim 19, wherein
Z is selected from




- 229 -




and the pharmaceutically acceptable salts thereof.

- 5 21. The compound of Claim 20, wherein Z is selected from



10


- 230 -



and the pharmaceutically acceptable salts thereof.


5

22. The compound of Claim 21 of the formula



wherein X is selected from

- 231 -



Y is selected from $-(CH_2)_r-$ or $-(CH_2)_m-NR^3-(CH_2)_t-$;

R^3 is selected from

- hydrogen,
- 5 aryl- $(CH_2)_p-$,
- C₁₋₅ alkoxy carbonyl,
- C₃₋₈ cycloalkyl,
- (aryl)_qaminocarbonyl,
- (aryl C₁₋₅ alkyl)_qaminocarbonyl,
- 10 C₁₋₈ alkyl,
- aryl C₁₋₆ alkyl,
- C₁₋₈ alkylsulfonyl,
- arylsulfonyl,
- aryl C₁₋₆ alkylsulfonyl,
- 15 C₁₋₈ alkoxy carbonyl,
- aryloxy carbonyl,
- aryl C₁₋₈ alkoxy carbonyl,
- C₁₋₈ alkyl carbonyl,
- aryl carbonyl,
- 20 aryl C₁₋₆ alkyl carbonyl,
- (C₁₋₈ alkyl)_qaminocarbonyl,
- C₁₋₆ alkylsulfonyl, or
- aryl C₁₋₆ alkyl carbonyl,
- wherein any of the alkyl groups may be unsubstituted or
- 25 substituted with R^{13} and R^{14} ;

R^4 is selected from

- hydrogen, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloheteroalkyl,
- C₃₋₈ cycloalkyl C₁₋₆ alkyl, C₃₋₈ cycloheteroalkyl C₁₋₆ alkyl,
- 30 aryl or aryl C₁₋₈ alkyl,

- 232 -

R^8 is selected from


- hydrogen,
 aryl,
 5 aryl-(CH₂)_p-,
 CH≡C-(CH₂)_s-,
 C₁₋₆ alkyl-C≡C-(CH₂)_s-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_s-,
 aryl-C≡C-(CH₂)_s-,
 10 C₁₋₆ alkylaryl-C≡C-(CH₂)_s-,
 CH₂=CH-(CH₂)_s-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_s-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_s-,
 aryl-CH=CH-(CH₂)_s-,
 15 C₁₋₆ alkylaryl-CH=CH-(CH₂)_s-,
 C₁₋₆ alkyl-SO₂-(CH₂)_s-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_s; and

r is an integer from 0 to 3;

20


and the pharmaceutically acceptable salts thereof.

23. The compound of Claim 22 of the formula




25 wherein Z is selected from

- 233 -



R^8 is selected from
hydrogen,



- 5 indolyl- $(CH_2)_p$ -,
 $CH \equiv C-(CH_2)_s$ -,
 C_{1-6} alkyl- $C \equiv C-(CH_2)_s$ -,
 C_{3-7} cycloalkyl- $C \equiv C-(CH_2)_s$ -,
aryl- $C \equiv C-(CH_2)_s$ -,
10 C_{1-6} alkylaryl- $C \equiv C-(CH_2)_s$ -,
 $CH_2=CH-(CH_2)_s$ -,
 C_{1-6} alkyl- $CH=CH-(CH_2)_s$ -,
 C_{3-7} cycloalkyl- $CH=CH-(CH_2)_s$ -,
aryl- $CH=CH-(CH_2)_s$ -,
15 C_{1-6} alkylaryl- $CH=CH-(CH_2)_s$ -,

- 234 -

C₁₋₆ alkyl-SO₂-(CH₂)_s-,
C₁₋₆ alkylaryl-SO₂-(CH₂)_s-; and

R¹² is selected from hydrogen or C₁₋₈ alkyl; and
5 s is an integer from 0 to 3;
and the pharmaceutically acceptable salts thereof.

24. The compound of Claim 23 selected from

10 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperidin-
1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperin-
1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine trifluoroacetate;

15 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;

20 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-
pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;

25 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;

30 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

- 235 -

- 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]-pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine ethyl ester;
- 5 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;
- Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- 10 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-tetrahydropyrimidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- Ethyl 2-oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- 15 2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3(S)-pyridin-3-yl- β -alanine;
- Ethyl 2-oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;
- 20 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;
- 25 Ethyl 3-(2-[2-oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino]-pyrrolidin-1-yl)-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;
- 30 3-(2-[2-Oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino]pyrrolidin-1-yl)-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;
- 3-[2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;

- 236 -

- 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-
pyridin-3-yl-propionic acid;
- 5
- 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-
ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl ester;
- 10
- 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;
- 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)-
ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine ethyl ester;
- 15
- 2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;
- 20
- 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine ethyl
ester;
- 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;
- 25
- 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-
tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-
acetylamino)-propionic acid ethyl ester;
- 30
- 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-
tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-
acetylamino)-propionic acid;
- 3-{2-(2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-
pyrrolidin-1-yl)-acetylamino}-3(S)-quinolin-3-yl-propionic acid;

3-(2-(5(S)-Ethyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl)-acetylamino)-3-(S)-quinolin-3-yl-propionic acid trifluoroacetate;

5

3-(2-{6-Methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}-2H-pyridin-1-yl)-acetylamino)-3(S)-pyridin-3-yl-propionic acid bis trifluoroacetate; or

10 3-(2-{6-Methyl-2-oxo-3-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)methyl]-amino}-2H-pyridin-1-yl)-acetylamino)-3(S)-pyridin-3-yl-propionic acid ethyl ester;

and the pharmaceutically acceptable salts thereof.

15

25. The compound of Claim 24, selected from

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]piperin-1-yl-acetyl-3(S)-pyridin-3-yl-β-alanine trifluoroacetate;

20

2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

25 2-Oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;

2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl-β-alanine;

30 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl-β-alanine;

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-imidazolidin-1-yl-acetyl-3-(S)-pyridin-3-yl-β-alanine;

- 238 -

2-Oxo-3-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-yl)ethyl]-
tetrahydropyrimidin-1-yl-acetyl-3-(S)-pyridin-3-yl- β -alanine;

5 2-Oxo-3(R)-[2-(5,6,7,8-tetrahydro[1,8]naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(R)-(2-ethylindol-3-yl)- β -alanine;

10 3-(2-{2-Oxo-3(S)-[(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
amino]pyrrolidin-1-yl}-acetylamino)-3-(S)-pyridin-3-yl-propionic acid;

3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
hexahydro-(3aS, 6aS)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-
pyridin-3-yl-propionic acid; or

15 3-{2-[6-Oxo-1-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-ylmethyl)-
hexahydro-(3aR, 6aR)pyrrolo[3,4-b]pyrrol-5-yl]-acetylamino}-3-(S)-
pyridin-3-yl-propionic acid;

20 2-Oxo-5(R)-methyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine;

2-Oxo-5(S)-benzyl-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
yl)ethyl]pyrrolidin-1-yl)acetyl-3(S)-pyridin-3-yl- β -alanine;

25 5(R)-Methyl-2-oxo-3(S)-[2-(5,6,7,8-tetrahydro[1,8]-naphthyridin-2-
ylmethyl)-amino]pyrrolidin-1-yl)acetyl-3(S)-alkynyl- β -alanine; or

30 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-{2-oxo-3(S)-[2-(5,6,7,8-
tetrahydro-[1,8]naphthyridin-2-yl)-ethyl]-pyrrolidin-1-yl}-
acetylamino)-propionic acid;

and the pharmaceutically acceptable salts thereof.

- 239 -

26. A pharmaceutical composition comprising the compound of Claim 19 and a pharmaceutically acceptable carrier.

27. A pharmaceutical composition made by combining a
5 compound of Claim 19 and a pharmaceutically acceptable carrier.

28. A process for making a pharmaceutical composition comprising combining a compound of Claim 19 and a pharmaceutically acceptable carrier.
10

29. A method of eliciting a vitronectin receptor antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 26.
15

30. A method of treating or preventing a condition mediated by antagonism of a vitronectin receptor in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 26.
20

31. A method of inhibiting bone resorption in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 26.

25 32. A method of treating osteoporosis in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 26.

30 33. A method of treating tumor growth in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound of Claim 19 and one or more agents known to be cytotoxic or antiproliferative.

INTERNATIONAL SEARCH REP RT

International application No.
PCT/US97/14912**A. CLASSIFICATION F SUBJECT MATTER**

IPC(6) : C07D 401/06; A61K 31/415, 31/435

US CL : 514/300; 546/122

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/300; 546/122

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

APS SEARCH 1975 -TO DATE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS COMPUTER SEARCH 1966-TO DATE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,260,277 A (MCKENZIE) 09 NOVEMBER 1993, SEE ENTIRE DOCUMENT.	1-33
A	US 5,534,524 A (BONEWALD ET AL) 09 JULY 1996, SEE ENTIRE DOCUMENT.	1-33
X, P — Y, P	US 5,668,159 A (JIN ET AL) 16 SEPTEMBER 1997, SEE ENTIRE DOCUMENT, ESPECIALLY COLUMN 3.	1-5, 8-19 AND 26-33 1-5, 8-19, AND 26-33

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

24 SEPTEMBER 1997

Date of mailing of the international search report

15 OCT 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ZINNA N. DAVIS

Telephone No. (703) 308-1235